





**THREE NEW SPECIES OF *GEKKO*  
AND REMARKS ON *GEKKO HOKOUENSIS*  
(LACERTIFORMES, GEKKONIDAE)**

**Kai-ya Zhou<sup>1</sup>, Yue-zhen Liu<sup>1</sup>, and De-jun Li<sup>2</sup>**

<sup>1</sup> Department of Biology, Nanjing Normal College

<sup>2</sup> Department of Biology, Zunyi Medical College

In: *Acta Zootaxonomica Sinica* 7(4): pp.438-446+pls.1-2,  
published in October 1982.

Translated by

Hidetoshi Ota<sup>3</sup>, Tsutomu Hikida<sup>3</sup> & Kuang-yang Lue<sup>4</sup>

<sup>3</sup> Department of Zoology, Kyoto University

<sup>4</sup> Department of Biology, National Taiwan Normal University

**SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
No. 77**

**1989**

SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with us.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560, U.S.A. Please include a self-addressed mailing label with requests.

TRANSLATORS' NOTES

In preparing the English version from the original (in Chinese, with English summary), we attempted to make as literal a translation as possible. However, a few minor changes were made with footnotes (\* - \*\*\*\*); these footnotes follow the references. Locality names were written in Continental spellings, followed by Taiwanese spellings in parentheses at their first appearance.

We thank R.I. Crombie and G.R. Zug for their assistance and encouragement during the process of preparation of the present manuscript.

## INTRODUCTION

Among the gekkonid genera occurring in China, *Gekko* is the largest group with the widest range of distribution. Six species and two subspecies have hitherto been known for the genus from China (Stejneger, 1932; Chen, 1969). During 1975 to 1980, Department of Biology, Nanjing Normal College collected 1637 specimens of *Gekko* from Hebei (Hopei), Shanxi (Shansi), Shaanxi (Shensi), Shandong (Shantung), Henan (Honan), Jiangsu (Kiangsu), Anhui (Anhwei), Hubei (Hupeh), Sichuan (Szuchuan), Zhejiang (Chekiang), Fujian (Fukien), Jiangxi (Kiangsi), Hunan (Hunan), Guizhou (Kweichow), Yunnan (Yunnan), Guangdong (Canton), Hainan (Hainan)\*, and Guangxi (Kwangsi) Provinces. Several specimens were also collected from Guizhou Province by Department of Biology, Zunyi Medical College. While studying these specimens, three new species were discovered. On the other hand, *Gekko japonicus hokouensis* from Yanshan (Chainshan) Prefecture, Jiangxi Province, proved to represent a good species. Consequently, ten *Gekko* species\*\* are presently recognized from China as follows:

### Key to species of *Gekko* in China

1. Rostral separated from nostril ..... 2  
Rostral in contact with nostril ..... 3
2. Body relatively large, longer than 200 mm in total length; tubercles not particularly concentrated in upper margin of ear opening ..... *G. gekko*  
Body relatively small, shorter than 150 mm in total length; upper margin of ear opening with cluster of enlarged conical tubercles concentrated in high density .... *G. auriverrucosus* sp. nov.
3. Male with 24 femoral pores in each side ..... *G. kikuchii*  
Male with preanal, or preanal-femoral pores ..... 4
4. A single enlarged spur on each side of base of tail ..... 5  
Enlarged spurs, two to three, with slight variation, on each side of base of tail ..... 8
5. Webs between digits evident ..... 6  
Webs between digits very slight or absent ..... 7
6. Tubercles absent on dorsum of body; male with 7-11 preanal pores ..... *G. subpalmatus*  
Tubercles present on dorsum of body; male with 17-27 preanal pores ..... *G. chinensis*
7. Supranasals in contact; dorsal tubercles flat; head and body length reaching 80 mm .....  
..... *G. liboensis* sp. nov.  
Supranasals separated by a small scale; dorsal tubercles relatively convex; head and body length not greater than 70 mm ..... *G. hokouensis*
8. Granular scales on dorsum of body relatively large; dorsal tubercles flat, in low density .....  
..... *G. swinhonis*  
Granular scales on dorsum of body relatively small; dorsal tubercles relatively convex, in high density ..... 9
9. Dorsal surfaces of body, thigh, and shank with much enlarged tubercles .....  
..... *G. scabridus* sp. nov.  
Tubercles moderately enlarged in dorsal surfaces of body and shank, usually lacking in thigh...  
..... *G. japonicus*





Fig. 1. *Gekko auriverrucosus* sp. nov. a. Dorsal view of head; b. Ventral view of head.

***Gekko auriverrucosus* Zhou et Liu sp. nov. (Plate I: 2, Fig. 1)**

Holotype—Male (NNC 80275), collected from Hejin (Hojin) Prefecture, Shanxi Province (alt. 459 m), on 19 August 1980. Allotype: female (NNC 80243), collection date and locality as for holotype. Paratypes: 33 males and 41 females, collected from Hejin, Yongji (Engtsi), and Linyi (Linyi) Prefecture, Shanxi Province. Collector: Xin-rong Xu. Type specimens are deposited in Department of Biology, Nanjing Normal University.

Diagnosis—Rostral separated from nostril; upper margin of ear opening with cluster of enlarged conical tubercles gathering in high density; tubercles uniformly scattered in temporal and occipital regions, neck, and dorsal surfaces of body, base of tail and limbs; male with 8-11 preanal pores.

Description—Snout about twice as long as eye diameter, distinctly longer than distance between eye and ear opening; diameter of ear opening 0.9-1.5 mm, about 30-44% of eye diameter; rostral twice as broad as high, angulated at midpoint dorsally, separated from nostril; nostril surrounded by first supralabial, supranasal, and two small scales; supranasals moderately enlarged, slightly longer than broad, separated from each other by a minute scale, or in contact with each other medially; supralabials 9-11; infralabials 9-11; mental pentagonal; chin shields forming several rows of transverse arches; first row normally comprising five shields, each slightly longer than broad, median three largest; scales following chin shields and reaching gular region uniform, granular (Fig. 1).

About 12 scales between nostril and eye; interorbital scales about 25; upper margin of ear opening with tubercle cluster comprising about six enlarged conical tubercles gathering in high density; around jaw angle and preotic region also with enlarged conical tubercles; tubercles uniformly scattered among dorsal granular scales, from temporal and occipital regions to base of tail, in 16-20 irregular rows at midbody; dorsal surfaces of forelimbs covered with small tubercles; on dorsal surfaces of hindlimbs, tubercles scattered among granular scales; scales granular in gular region, imbricate in the other part of ventral surface of body; webs between digits rudimentary; underneath dilated portions of toes with lamellae, 6-8 on toe I, 6-8 on toe II, 6-8 on toe III, 6-8 on toes IV, and 7-9 on toe V; male with 8-11, mostly 8-9 preanal pores.

Table 1. Measurements (in mm) of specimens of *Gekko auriverrucosus*.

| Specimens                      | Total length                   | Eye diameter | Diameter of ear opening | Snout length | Head length | Axilla groin length | Fore-limb length | Hind-limb length |
|--------------------------------|--------------------------------|--------------|-------------------------|--------------|-------------|---------------------|------------------|------------------|
| <b>Holotype</b><br>(NNC 80275) | 125.5 (62+63.5)                | 3.2          | 1.2                     | 6.7          | 15          | 28                  | 18.5             | 24.5             |
| <b>Allotype</b><br>(NNC 80243) | 135.5 (65.5+70)                | 3.3          | 1.3                     | 7.3          | 16          | 30.5                | 19               | 27               |
| <b>Paratypes</b>               |                                |              |                         |              |             |                     |                  |                  |
| 11 males<br>from Hejin         | 119 (59+60)-<br>130 (63+67)    | 3-<br>3.5    | 1-<br>1.3               | 6.3-<br>7    | 14-<br>16   | 25-<br>29.5         | 16-<br>19        | 24-<br>25        |
| 16 females<br>from Hejin       | 117 (56+61)-<br>133 (65+68)*** | 3-<br>3.6    | 1-<br>1.5               | 6.4-<br>7.5  | 14-<br>17   | 27-<br>33           | 16.5-<br>20      | 23-<br>28.5      |

Tail slightly compressed, with two or three enlarged spurs in each side at base; dorsum of tail covered with tubercles of various sizes; annular grooves in about every sixth to eighth tubercle; venter of tail with a longitudinal row of laterally elongated shields.

Dorsal ground color of preserved specimen pale gray; a brown bar from nostril through eye and ear to shoulder; top of head with brown markings; dorsal surfaces of neck and body with 5-6 transverse brown bands; dorsum of tail with 9-13 transverse brown bands; posterior edge of transverse bands in body and tail darkly edged; dorsal surfaces of four limbs also with transverse brown bands; venter of body light reddish yellow.

This new species might be easily misidentified as *G. japonicus*. However, the latter species has a rostral entering the nostril, and lacks a cluster of tubercles in the upper margin of the ear opening. Thus, *G. japonicus* is actually distinct from the present new species.

In the natural habitat, the density of *G. auriverrucosus* is very high. It prefers to perch on high portions of walls, and occasionally appears on artificially lighted areas to search for prey. In June and July, the present species has its reproductive season. Juveniles collected between 19 and 22 August had already reached 31-32.5 mm in head and body length. All adult females collected on the same date from the same locality with the above juveniles possessed no mature eggs. About 1/6 of the total sample had parasitic mites, especially in high density on digits.

***Gekko liboensis* Zhou et Li sp. nov.** (Plate II: 1, Fig. 2)

Holotype—Female (TMC 791669), Chengguan (Chengkwan), Libo (Libo) Prefecture, Guizhou Province (alt. 430 m), on 5 July 1979, by Zhi-lu Zhao. This specimen is deposited in Department of Biology, Zunyi Medical College.




Fig. 2. *Gekko liboensis* sp. nov. a Dorsal view of head; b. Ventral view of head.

**Diagnosis**—Head and body length exceeding 80 mm; supranasals large, in contact with each other medially; flat, cycloid tubercles among dorsal granular scales, forming about 10 irregular longitudinal rows; webs evident between fingers I, II and III, very slight between fingers III, IV and V.

**Description**—Holotype very large, head and body length nearly 85 mm; snout 1.8 times as long as eye diameter, distinctly longer than distance between eye and ear opening; diameter of ear opening 2 mm, about 40% of eye diameter; rostral broader than deep, upper margin slightly concaved at midpoint; nostril surrounded by rostral, first supralabial, supranasal, and two small scales; supranasals large, in contact with each other medially; supralabials 12; infralabials 11; mental triangular; median pair of chin shields twice as long as broad, posteriorly entered by a pair of small polygonal chin shields (Fig. 2).

About 18 scales between nostril and eye; interorbital scales about 40; flat, cycloid tubercles uniformly scattered in low density on dorsum, from parietal and occipital regions to base of tail, forming about 10 irregular rows on body; fore- and hindlimbs without tubercles; ventral surface posterior to neck covered with imbricate scales; 10 enlarged scales in preanal region.

Underneath dilated portions of fingers covered with lamellae, eight on finger I, eight on finger II, nine on finger III, nine on finger IV, and eight or nine on finger V; rudimentary webs evident

Table 2. Measurements (in mm) of a specimen of *Gekko liboensis*.

| Specimens   | Total length    | Eye diameter | Diameter of ear opening | Snout length | Head length | Axilla groin length | Fore-limb length | Hind-limb length |
|---|-----------------|--------------|-------------------------|--------------|-------------|---------------------|------------------|------------------|
| <b>Holotype</b><br>(TMC 791669)<br>(regenerated tail) | 121.8 (84.8+37) | 5            | 2                       | 9.2          | 21          | 37                  | 25.2             | 35               |



between fingers I, II and III, faintly between fingers III, IV and V; margins of webs attaching to proximal one third of toes; hindlimb much developed, its length 95% of axilla to groin length; underneath dilated portions of toes covered with lamellae, eight on toe I, seven or eight on toe II, eight on toe III, nine on toe IV, and nine on toe V; rudimentary webs evident between toes I, II, III and IV; a single large spur on each side of base of tail; tail regenerated, very short.

Dorsal ground color in preservative grayish tan; a brown bar running along lower margin of eye, almost reaching to ear opening posteriorly; dorsal surfaces of neck and body with nine transverse brown bands; dorsal surfaces of limbs also with transverse brown bands; venter of body pale reddish yellow.

This new species greatly resembles *G. hokouensis*. However, the latter has supranasals separated from each other, and conical dorsal tubercles. Moreover, the head and body length of *G. hokouensis* is shorter than 70 mm.

*G. liboensis* is rarely observed at Chengguan, Libo Prefecture.

### *Gekko hokouensis* Pope

*Gekko japonicus hokouensis* Pope, 1928, Amer. Mus. Novitates 325: 1-2 (Yanshan Prefecture, Jiangxi Province)

Pope (1928) regarded this form as a subspecies of *G. japonicus*, and stated that *G. j. hokouensis* differs from the nominal subspecies only in the number of cloacal spurs; he noted that the former has a single spur on each side of the base of tail, whereas the latter has two or three spurs. While investigating a large series of specimens, we found that *hokouensis* has a relatively large spur, measuring about 2.2-3.1 mm for the male and 1.3-2.0 mm for the female in maximum diameter. Although the spur is more or less grooved and incompletely divided in a few males and most females, the outline of the single spur remains apparent in all animals (Plate II: 4-9). On the other hand, *japonicus* possesses two or three smaller spurs below three larger spurs. The size of each spur is relatively small, and the maximum diameter of the largest spur measured 1.2-1.5 mm in the male and 0.6-1.0 mm in the female (Plate II:10-11). Differences are recognizable between *hokouensis* and *japonicus* also in the condition of dorsal tubercles as follows. In *hokouensis*, tubercles are absent on the four limbs, and relatively few around the middle of the body. On the other hand, in *japonicus*, the dorsal surface of the forearm and shank is covered with tubercles, and the tubercles around the middle of the body are in relatively high density (Plate II:2-3, Table 3).

Table 3. Comparison of dorsal tubercles in *Gekko hokouensis* and *G. japonicus*.

| Species                 | N   | Localities                    | Occipital and neck | Body | Upper arm | Forearm | Thigh | Shank |
|-------------------------|-----|-------------------------------|--------------------|------|-----------|---------|-------|-------|
| <i>Gekko hokouensis</i> | 271 | 17 locations in six provinces | - / +              | +    | -         | -       | -     | -     |
| <i>Gekko japonicus</i>  | 747 | 50 locations in 12 provinces  | +                  | ++   | -         | +       | - / + | +     |




Fig. 3. Distributions of *Gekko hokouensis* and *Gekko japonicus*. Ranges of the former and the latter are outlined by broken and complete lines, respectively. Cross-marks indicate localities of specimens of *G. hokouensis* cited in the present study\*\*\*\*.

On the basis of the above characteristics, we identified 271 specimens out of 1018 of *G. japonicus* (sensu lato) as *hokouensis*, and the remaining 747 as *japonicus* (sensu stricto). The former specimens were collected from 17 prefectures of six provinces, and the latter from 50 prefectures or cities of 12 provinces. The sampling localities of *hokouensis* are scattered within the range of *japonicus* (Fig. 3).

Conditions of natural habitats also differ between these two gekkonids; while *hokouensis* has its habitat in montane environments, *japonicus* is widely distributed in cities and villages of the plain regions. In Jiujiang (Kiukiang) City, for example, *japonicus* is found in urban area, whereas *hokouensis* in Lushan-haihui (Lushan-haihui), Bailudong (Bailudong) and Guling (Kuling). In Yixing (Ising) Prefecture, *japonicus* and *hokouensis* occur in the lowland and montane areas, respectively. The mutual displacement between *hokouensis* (a mountain dwelling species) and *japonicus* (a plain dwelling species) around a sympatric area much resembles the displacement between *Eremias brenchleyii* and *E. argus*. However, there are some areas where both of these gekkonids are collected from the same point at the same time. For example, of the 25 specimens obtained within a city of Chong'an (Chungan) Prefecture, on 26-27 June 1978, 15 specimens were identified as *japonicus* and the remainder as *hokouensis*. No intermediate forms were found among the above specimens.

Based on the morphological distinctiveness, sympatric occurrence, and ecological differences, we remove *hokouensis* from the subspecific status of *G. japonicus*, and regard it as a distinct species *Gekko hokouensis* Pope.

Within China, *G. hokouensis* is distributed in Yixing and Lishui (Lishui), Jiangsu Province, Tonglu (Tonglu) (Stejneger, 1932), Longquan (Longchuan), Beiyandangshan (Beiyantangshan), Zhejiang Province, Pucheng (Pucheng), Chong'an, Wuyishan (Wuyishan), Shaowu (Shaowu), Youxi (Yousi), Fujian Province, Taiwan Province (Maki, 1923), Jinzhai (Ginshai; Xuzhou Normal University), Huangshan (Huangshan) and Taiping (Taiping; Chengdu Institute of Biology), Anhui Province, Yanshan, Ninggang (Ningkwang), Lushan, Jiangxi Province, Yizhang (Ichang; Zhengdu Institute of Biology), and Jiangyong (Kiangyong)<sup>1</sup>, Hunan Province. In Taiwan, *G. japonicus* has been reported from several localities (Chen, 1969). Maki (1923) described specimens of "*G. japonicus*" from Taiwan as possessing a single process in each side of the base of tail, and limbs covered only with granular scales. These characteristics are identical with those of *G. hokouensis*. Of the specimens of "*G. japonicus*" collected from Japan, some animals such as the one from Nagasaki (USNM 13563; Stejneger, 1907) were reported to have three spurs in each side at the base of the tail, and tubercles on the dorsum of the shank, whereas others such as specimens from Yamagawa (USNM 31821 and 31822; Stejneger, 1907) and Fukuoka (Okada, 1936: fig. 1) possess a single spur and lack tubercles on limbs. Nakamura and Uéno (1963) noted that *G. japonicus* has normally a single but occasionally two or three tubercles, and that some animals possess tubercles on limbs. In October 1981, one of us examined specimens under the care of Dr. Shun-ichi Uéno at National Science Museum, Tokyo. Of the 21 specimens there, 12 from Tokara Is. (0231, 02287, 02293-95), Tokunoshima I. (0924), Yoronjima I. (0524, 02288-90, 02292) and Iriomotejima I. (0475) were identified as *G. hokouensis*, and the other nine from Tokyo (0236, 02286, 02297-98), Kyoto (02302) and Tsushima I. (0038, 0851, 0853, 02296) as *G. japonicus*. These results indicate that "*G. japonicus*" in Japan actually includes both *G. hokouensis* and *G. japonicus* (*sensu stricto*).

***Gekko scabridus* Liu et Zhou sp. nov. (Plate I: 1, Fig. 4)**

*Gekko* sp. Hu Shu-chin, Djao Er-mie and Liu Cheng-chao, 1973, Acta Zoologica Sinica 19(2): 155, from Guiyang (Kweiyang)

Holotype—Male (NNC 80122), Yongren (Yongzen) Prefecture, Yunnan Province (alt. 1531 m), on 4 Aug. 1980. Allotype: female (NNC 80143), collection date and locality as for holotype. Paratypes: 16 males and 33 females collected from Yongren Prefecture, Yunnan Province, and Miyi (Miyi) Prefecture, Sichuan Province. Collector: Xin-rong Xu. Type specimens are deposited in Department of Biology, Nanjing Normal University.

Diagnosis—Tubercles covering dorsal surfaces of body and hindlimbs much enlarged; male with 10-15 preanal pores.

Description—Eye relatively large, its diameter longer than half, as long as 51.4-57.1%, of snout length; snout slightly longer than distance between eye and ear opening; diameter of ear opening 0.9-1.5 mm, about 23-41% of eye diameter; rostral rectangular, its breadth less than twice of height; in a few specimens, upper margin of rostral slightly concave dorsally at mid point; nostril surrounded by rostral, first supralabial, supranasal, and two small scales; supranasals moderately enlarged, slightly broader than long, separated from each other normally by a single scale, but in

<sup>1</sup> All the locality data without citations of authors or institution are based on the specimens deposited in Department of Biology, Nanjin Normal College.






Fig. 4. *Gekko scabridus* sp. nov. a. Dorsal view of head; b. Ventral view of head.

some specimens, by two scales, or in contact with each other medially; supralabials 9-11; infralabials 9-11; mental pentagonal; chin shields longer than broad, median pair largest, outer pair relatively small, continuously graded to small granules through three to four rows of small hexagonal scales (Fig. 4). About 12 scales between nostril and eye; interorbital scales about 30; tubercles scattered among dorsal granular scales in high density, from frontal, parietal, temporal and occipital regions of head to base of tail, in 17-21 irregular rows around middle of body, those on dorsum of body extremely enlarged; limbs covered with granular scales dorsally, tubercles present on limbs except for upper arms; tubercles on hindlimbs distinctly enlarged like those in dorsum of body; venter of body covered with imbricate scales; interdigital webbings rudimentary; underneath dilated portions of digits with lamellae, 6-9 on toe I, 6-9 on toe II, 7-9 on toe III, 7-9 on toe IV, and 7-10 on toe V; male with 10-15, mostly 12 or 13, preanal pores.

Tail slightly compressed, with two or three enlarged spurs on each side at base; only one specimen (NNC 80166) of 51 examined had a single enlarged spur on both sides; dorsum of tail covered with granular scales; annular grooves in about every seventh to ninth row of granules; grooves in proximal one third of tail margined by six to eight enlarged tubercles posteriorly, such tubercles gradually disappearing in remaining portion of tail; venter of tail covered with imbricate scales, median scales enlarged and irregularly arranged, paired or not paired, in distal two-thirds to four-fifths of tail.

Dorsal ground color of preserved specimen pale brown; two brown bars from nostril through eye to temporal region; dorsal surfaces of head, body and limbs with irregular brown spots and reticulations; 7-9 transverse bars on neck and body; dorsum of tail with 10-14 transverse brown bars; venter of body light reddish yellow.

*G. scabridus* closely resembles *G. japonicus*. However, these species differ from each other as follows. In *G. japonicus*, dorsal tubercles on the body and shanks are distinctly smaller than those in *G. scabridus*, and tubercles are normally lacking on thighs. Moreover, male *G. japonicus* normally has only 4-8 preanal pores.

In the natural habitat, *G. scabridus* occurs in very high densities, and is observed equally in lighted and dark areas of walls. Of the specimens examined, a few animals possessed parasitic

Table 4. Measurements (in mm) of specimens of *Gekko scabridus*.

| Specimens                      | Total length     | Eye diameter | Diameter of ear opening | Snout length | Head length | Axilla-groin length | Fore-limb length | Hind-limb length |
|--------------------------------|------------------|--------------|-------------------------|--------------|-------------|---------------------|------------------|------------------|
| <b>Holotype</b><br>(NNC 80122) | 112.5 (57.5+55)  | 3.7          | 1.3                     | 7            | 15.7        | 26                  | 17.5             | 26               |
| <b>Allotype</b><br>(NNC 80143) | 120.5 (58+62.5)  | 3.8          | 1.4                     | 7            | 15.7        | 27                  | 19               | 24.5             |
| <b>Paratypes</b>               |                  |              |                         |              |             |                     |                  |                  |
| 11 males                       | 114 (57+57)-     | 3.7-         | 1.2-                    | 7-           | 15.4-       | 25.5-               | 17.5-            | 25-              |
| Yongren                        | 138 (64+74)      | 4.2          | 1.5                     | 7.5          | 17          | 30                  | 20               | 27               |
| 12 females                     | 116.5 (56.5+60)- | 3.5-         | 1.1-                    | 6.3-         | 14.5-       | 25-                 | 16.5-            | 24-              |
| Yongren                        | 140.5 (63.5+77)  | 4            | 1.5                     | 7.3          | 17          | 30                  | 19               | 27               |

mites. Juveniles collected in the beginning of August had reached 28-33 mm in head and body length. All adult females collected in the same date from the same locality with the above juveniles had no mature follicles.

#### REFERENCES

- Chen, J.T.F. 1969. A Synopsis of the Vertebrates of Taiwan. Commercial Press; Taipei. (in Chinese)
- Hu, S.-C., E.-M. Djao, and C.-C. Liu. 1973. A survey of amphibians and reptiles in Kweichow Province, including a herpetofaunal analysis. *Acta Zoologica Sinica* 19(2): 149-178. (in Chinese with English summary)
- Maki, M. 1923. On the gekkonid lizards of Taiwan. *Zool. Mag.* 35: 193-202. (in Japanese)
- Nakamura, K., and S.-I. Uéno. 1963. Japanese Reptiles and Amphibians in Colour. Hoikusha; Osaka. (in Japanese)
- Okada, Y. 1936. Studies on the lizards of Japan. Contribution I. Gekkonidae. *Sci. Rep. Tokyo Bunrika Daigaku*, B 2(42): 233-289.
- Pope, C.H. 1928. Four new snakes and a new lizard from South China. *Amer. Mus. Novitates* 325: 1-4.
- . 1929. Notes on reptiles from Fukien and other Chinese Provinces. *Bull. Amer. Mus. Nat. Hist.* 58: 335-487.
- . 1935. The Reptiles of China. *Nat. Hist. Central Asia*. Vol. 10. Amer. Mus. Natur. Hist., New York.
- Stejneger, L. 1907. Herpetology of Japan and adjacent territory. *Bull. U.S. Natl. Mus.* 58(1): 1-575.
- . 1932. The Chinese lizards of the genus *Gekko*. *Proc. U.S. Natl. Mus.* 82(3): 1-8.
- Wermuth, H. 1965. Liste der rezenten Amphibien und Reptilien, Gekkonidae, Pygopodidae, Xantusiidae. Walter de Gruyter & Co.; Berlin.

**TRANSLATORS' FOOTNOTES**


\* In the original, Hainan Island was included in Guandong Province. However, this island was removed from the latter to form an independent province, Hainan Province, by itself in 1987.

\*\* The original states "10 *Gekko* species and one subspecies", but it lists only 10 species in the key.

\*\*\* Table 1 in the original contains "113 (65+68)", but this must be a typographical error.

\*\*\*\* The figure legend in the original states that the cross-marks in Fig. 3 indicate localities where *G. japonicus* was collected. But it is evident, from the content of the text, that those marks actually represent sampling localities of *G. hokouensis*.





1. *Gekko scabridus* Liu et Zhou, sp. nov. Lateral view of head (above), and dorsal view (below).
2. *Gekko auriverrucosus* Zhou et Liu, sp. nov. Lateral view of head (above), and dorsal view (below).



1. *Gekko liboensis* Zhou et Li, sp. nov. Dorsal view.
2. *Gekko hokouensis*. Dorsal view of base of tail and hind limbs.
3. *Gekko japonicus*. Dorsal view of base of tail and hind limbs.
- 4-9. *Gekko hokouensis*. Enlarged spurs on right side of base of tail in males (4 and 5) and females (6 to 9). Spurs in 4 and 6 have no shallow furrows, whereas those in the remainders (5, 7, 8, and 9) show more or less developed furrows incompletely dividing the spurs.
- 10-11. *Gekko japonicus*. Spurs on right side of base of tail in a male (10) and a female (11).







6  
07

INDEX TO THE BIOGRAPHIES  
OF  
HERPETOLOGISTS & ICHTHYOLOGISTS  
COPEIA 1913-1988



C. J. McCOY

Section of Amphibians and Reptiles  
Carnegie Museum of Natural History

SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
NO. 78

1989

SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with us.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560, U.S.A. Please include a self-addressed mailing label with requests.

Let us now praise famous men, and our fathers that begat us. All these were honored in their generations, and were the glory of their times (Ecclesiasticus 44:1).

## INTRODUCTION

The history of a science consists not only of the chronology of development of knowledge and ideas, but equally of the life stories of men and women who contribute to that science. For ichthyology and herpetology a significant part of this human record is contained in seventy-five years of the journal Copeia. Unfortunately this rich source of historical and biographical information has been largely inaccessible, as most of the death notices, obituaries, and biographical sketches in Copeia appeared in the "Editorial Notes and News" section of the journal, and are not included in the journal's annual index. This compilation will provide ready access to biographies in Copeia and, hopefully, stimulate an appreciation for the personalities who fostered modern ichthyology and herpetology.

The index consists of three lists: the biographical articles arranged alphabetically by name of subject and keyed to the bibliography by number, an annual list of biographies, and a bibliography. The content of articles is coded as follows: "N" is a death notice only; "B" is a biography; "P" indicates a portrait is included; "L" identifies a biography with a bibliography. The biographies range in length from a few words to several pages. Although the distinction is arbitrary in some cases, the "B" category is reserved for articles that have substantial biographical content. Three major articles in Copeia (Dymond, 1964, (1):2-33; Myers, 1964, (1):34-41; Hubbs, 1964, (1):42-60) are not indexed. Although they contain biographical information, these articles are primarily historical reviews.

Much of the credit for documenting the lives of ichthyologists and herpetologists in Copeia belongs to two of our late colleagues, incidentally also my "scientific grandfathers." The first of these, Carl L. Hubbs, had an unusually strong sense of the importance of the historical perspective. Perhaps this was because his career spanned critical years in the transition to the modern era, as did that of his mentor David Starr Jordan. Hubbs was by far the most prolific contributor of biographical articles to Copeia (24 signed articles and dozens of unsigned notes). Moreover, as first ichthyological editor (1930-1937) of the "new series" of Copeia, he pioneered regular incorporation of biographical information in "Editorial Notes and News." George S. Myers is second to Hubbs in number of biographies published in Copeia, but his contributions are models of sympathetic and insightful biography. To these two, indeed to all who have contributed to this unique record, we owe our gratitude. I thank Alan E. Leviton for (indirectly) suggesting this project, and M. Therése Giles for enduring patience.

BIOGRAPHIES

Alcock, A. W.:30  
 Allen, E. J.:98  
 Allen, W. E.:131  
 Allen, William Ray:169  
 Allis, Jr., Edward Phelps:132  
 Alm, Gunnar:327  
 d'Ancona, Umberto:208  
 Anderson, Jr., James Donald:419  
 Anderson, Paul:201  
 Andersson, Lars Gabriel:149  
 Andrews, Roy Chapman:188  
 Angel, Fernand:143, 307  
 Artedi, Peter:352  
 Atkinson, N. J.:25  
 Awerinzew, Sergius:114  
 Ayers, Howard:32

Babbitt, Lewis H.:235  
 Babcock, Harold Lester:350  
 Babcock, John Pease:48  
 Bacon, Daniel:35  
 Baranov, Fedor Ilich:373  
 Barbour, Thomas:289  
 Barnard, K. H.:209  
 Barnhart, Percy S.:157  
 Bass, Jr., John F.:91  
 Bauman, Aden C.:134  
 Beaufort, Lieven Ferdinand de:264  
 Bennett, George W.:237  
 Bensley, B. A.:34  
 Berezovsky, A. I.:115  
 Berg, Leo Semenovich:147, 388, 389  
 Bertin, Léon:172  
 Bigelow, Henry Bryant:285  
 Bishop, Sherman C.:150, 305  
 Blair, William Franklin:341  
 Blanchard, Frank Nelson:376  
 Blegvad, Harald:155  
 Böhlke, James E.:374  
 Borodin, Nicholas A.:70  
 Boulenger, E. G.:122  
 Boulenger, George Albert:66, 395  
 Bragg, Arthur Norris:263  
 Breder, Jr., C. M.:252  
 Breyer-de Rooy, P. J.:415  
 Brimley, Clement Samuel:281  
 Bromund, Fred:108

Brumwell, Malcolm J.:97  
Bruun, Anton Frederik:197, 323  
Buen, Fernando de:324  
Burt, Charles E.:405

Cagle, Fred Ray:407  
Cameron, A. T.:135  
Camp, Charles L.:417  
Carr, Jr., Archie F.:291  
Chabanaud, Paul:177  
Chernov, Sergius Alexandrovich:203, 348  
Clark, Howard Walton:95, 96  
Clark, Hubert Lyman:129  
Clay, William Marion:277  
Clemens, Lucy Wright Smith:55  
Clemens, Wilbert A.:210  
Cobb, John N.:5, 380  
Cochran, Doris Mable:301  
Cockerell, T. D. A.:136  
Conant, Isabelle:236  
Cook, Fannye A.:206  
Cooper, Gerald Paul:256  
Cope, Edward Drinker:269, 298, 357  
Cowles, Raymond Bridgman:266  
Crawford, Sr., Donald R.:195  
Creaser, Charles W.:302  
Crimmins, Martin Lalor:168  
Cunningham, J. T.:42  
Curtis, Brian:191  
Cyren, Otto:128  
Czopek, Juliusz:403

Dahlgren, Ulric:123  
Daniel, John Franklin:99  
Davies, David H.:329  
Davis, D. Dwight:211  
De Haas, C. O. J.:141  
Ditmars, Raymond L.:375  
Donoso-Barros, Roberto:368  
Duncker, Georg:166  
Dunn, Emmett Reid:171, 378  
Dybowski, Benedikt:6  
Dymond, John Richardson:212, 385

Eager, Grace:137  
Eaton, Theodore H.:412  
Eddy, Brayton:148  
Eddy, Samuel:414  
Ege, Vilhelm:325  
Ehrenbaum, Ernst:133  
Ehrenreich, Alfred:16  
Eigenmann, Rosa Smith:127



Einarsson, Hermann:218  
 Ellis, Max M.:164  
 Embody, George Charles:74, 80  
 Engeling, Gus A.:159  
 Erwin, Richard P.:21  
 Evermann, Barton Warren:22, 310  
 Ewing, H. E.:152

Fassett, Harry C.:311  
 Flower, Stanley Smyth:121, 396  
 Forbes, Stephen Alfred:7  
 Fowler, Henry Weed:276  
 Fox, Jr., Wade:282  
 Frizzell, Don L.:231

Gage, Simon Henry:109  
 Gaige, Frederick MacMahon:254  
 Gaige, Helen Thompson:254  
 Giltay, Louis Pierre:59  
 Ginsburg, Isaac:360  
 Glauert, L.:202  
 Gloyd, Howard Kay:271  
 Goff, Carlos Clyde:73  
 Goin, Coleman J.:273  
 Goodnoh, Jr., Clifford A.:116  
 Gordon, Myron:178, 253  
 Gowanloch, James Nelson:161  
 Greenbank, John T.:240  
 Greenberg, Bernard S.:214  
 Grey, Marion:399  
 Grinnell, Joseph:82, 88  
 Grodziński, Zygmunt:402  
 Gudger, E. W.:173, 364  
 Günther, Klaus:290

Haas, Georg:418  
 Hamilton, Rodgers D.:145  
 Hankinson, Thomas L.:45  
 Hanko, Bela:184  
 Hansen, Donald F.:238  
 Harrington, Jr., Robert W.:251  
 Hart, John L.:386  
 Hartweg, Norman E.:204  
 Hass, Robert L.:222  
 Hay, Oliver Perry:11  
 Hay, William Perry:126  
 Henderson, Junius:67  
 Herald, Earl Stannard:361  
 Herre, Albert W.:198  
 Hildebrand, Samuel Frederick:139, 315, 319, 379, 383, 384  
 Hine, James S.:12  
 Hollister, Gloria:261

Hoover, Earl E.:72  
 Hora, Sunder Lal:392  
 Howard, John K.:213  
 Hoyt, Southgate Y.:153  
 Hubbs, Carl Leavitt:342, 353, 365, 391  
 Hyde, Jesse Earl:52

Ihering, Rodolpho von:421  
 Ishikawa, Chiyomatsu:40

Jacot, Arthur Paul:81  
 Jenkins, Oliver Peebles:41  
 Johansen, A. C.:18  
 Johnson, Jr., J. E.:167, 377  
 Johnson, Richard M.:349  
 Johnstone, James:26  
 Jordan, David Starr:15, 292, 296, 387  
 Jordan, Eric Knight:306  
 Juday, Chauncey:106

Kampen, P. N. van:69  
 Kelly, Howard A.:100  
 Kendall, William Converse:75, 79  
 Kishinouye, Kamakichi:8  
 Klauber, Laurence Monroe:390  
 Klugh, A. Brooker:19  
 Knight, A. P.:46  
 Koumans, Frederick P.:265  
 Kuehne, Robert Andrew:257

Lagler, Karl F.:355  
 Lahille, Fernando:94  
 Landreth, Hobart:233  
 Lantz, Louis A.:366  
 Lataste, Fernand:36  
 Limbaugh, Conrad:322  
 Lindberg, G. U.:2  
 Loftus-Hills, Jasper J.:1  
 Logier, Eugene Bernard Shelley:278  
 Longley, William Harding:53  
 Lönnberg, Einar:320  
 Lowe, John N.:71  
 Lübbert, Hans:160  
 Lutz, Adolpho:93  
 Lutz, Bertha:345  
 Lutz, Gualter Adolpho:223

Maki, Moichiro:183  
 Mann, William M.:189, 194  
 March, Douglas D. H.:83  
 Maria, Niceforo:404  
 Martof, Bernard S.:255

Maskell, F. G.:31  
 Maslin, T. Paul:393  
 Matsubara, Kiyomatsu:331  
 Mayer, Fritz:179  
 McGregor, Richard Crittenden:56  
 McIlhenny, Edward Avery:268  
 McMurrich, James Playfair:76, 87  
 Meade, George P.:288  
 Merriman, R. Owen:39  
 Mertens, Robert:300  
 Metzelaar, Jan:4  
 Michałowski, Jerzy:219  
 Miller, Frances Voorhees Hubbs:270  
 Miller, Richard B.:185  
 Miller, Richard D.:180  
 M'Intosh, W. C.:20  
 Miranda Ribeiro, Alipio de:84, 356  
 Miranda Ribeiro, Paulo de:215  
 Mitchell, F. J.:224  
 Mohr, Erna:332  
 Morgan, Stanley Stewart:111  
 Mosauer, Walter:57  
 Mukerji, Dev Dev:60  
 Müller, Lorenz:313  
 Myers, George Sprague:272

Nakamura, Hiroshi:413  
 Needham, Paul R.:207  
 Nellesose, (Captain):119  
 Nelson, Edward William:37  
 Nichols, John Treadwell:297  
 Nikolsky, G. V.:398  
 Noble, G. K.:92  
 Norman, John R.:107, 411

Okada, Yaichiro:339  
 Okkelberg, Peter O.:192  
 O'Malley, Henry:47  
 Ooster, John van:217  
 Osburn, Raymond C.:170  
 Ovchynnyk, Michael M.:225

Pack, Herbert J.:9  
 Parker, H. W.:303  
 Patten, William:23  
 Perkins, C. B.:416  
 Peters, James A.:274  
 Peyer, Bernhard:200  
 Pfaff, J. R.:181  
 Pflueger, Al:326  
 Pietschmann, Viktor:174  
 Plate, Ludwig:68

Pope, Clifford H.:343  
 Proctor, Joan:351

Raney, Charlotte F.:244  
 Ravenel, William de Chastignier:33  
 Redeke, H. C.:120  
 Reed, Hugh D.:61  
 Reed, Roger J.:242  
 Reese, Albert Moore:216  
 Regan, Charles Tate:101, 410  
 Reid, Earl Desmond:382  
 Reighard, Jacob:340  
 Richardson, Sally Leonard:275  
 Rivas, Luis Rene:293  
 Rivero, Luis Howell:246  
 Romer, Alfred Sherwood:283  
 Rosen, Donn Eric:362  
 Roughley, T. C.:193  
 Roule, Louis:102  
 Rust, Hans Theodor:146  
 Ruthven, Alexander Grant:226

Schachter, D.:228  
 Schäferna, Karel:151  
 Schindler, Otto:321  
 Schmidt, Frank J. W.:44  
 Schmidt, Johannes:27  
 Schmidt, Karl Patterson:175, 280, 369  
 Schmidt, Peter J.:156  
 Schrenkeisen, Ray:50  
 Schroeder, William C.:284  
 Schulte, H. von W.:28  
 Schultz, Leonard Peter:400  
 Scofield, Norman B.:182  
 Scortecci, Giuseppe:409  
 Scott, Will:62  
 Seale, Alvin:314  
 Sette, Oscar Elton:232  
 Shannon, Frederick Albert:394  
 Shaw, Charles E.:229  
 Shetter, David S.:334  
 Shiraishi, Yoshikazu:335  
 Shufeldt, Robert Wilson:38  
 Silvester, C. F.:10  
 Sincock, Jr., Edwin H.:112  
 Slevin, Joseph Richard:354  
 Smith, E. Victor:89  
 Smith, George Milton:154  
 Smith, Hugh McCormick:260, 316, 381, 401  
 Smith, Mrs. Hugh M.:124  
 Smith, James Leonard Brierley:330  
 Smith, Margaret Mary:312

Smith, Philip Wayne:267  
 Snedigar, Robert:205  
 Snyder, John Otterbein:103, 317  
 Soldatov, V. K.:117  
 Starks, Chloe Leslie:358  
 Starks, Edwin Chapin:24  
 Starrett, William C.:230  
 Steinbach, José:13  
 Steinitz, Heinz:347  
 Stejneger, Leonhard:258, 259, 294, 295  
 Stensiö, Erik:363  
 Stephens, Frank:63  
 Stevenson, John Alexander:49  
 Steyn, William J.:221  
 Storer, Tracy I.:417  
 Storey, Margaret Hamilton:190, 359  
 Stoye, Frederick Hans:338  
 Strecker, John Kern:420  
 Stuart, Laurence Cooper:247, 287  
 Sumner, Francis B.:318  
 Surbeck, Georg:130  
 Surber, Thaddeus:142  
 Svetovidov, A. N.:3

Tanaka, Shigeho:336  
 Tåning, A. Vedel:176  
 Tanzer, Ernest C.:227  
 Taranetz, A. J.:113, 118, 125  
 Taylor, Edward Harrison:239, 286  
 Tchernavin, Vladimir V.:140  
 Tee-Van, John:250  
 Templeton, James R.:234  
 Terentjev, Paul V.:279  
 Thienemann, August:187  
 Thompson, William Francis:328  
 Tinkle, Donald W.:346  
 Titcomb, John W.:17  
 Townsend, Charles Haskins:105  
 Trewavas, Ethelwynn:243  
 Troemner, J. Louis:158  
 Turner, Clarence Lester:333  
 Twitty, Victor Chandler:344

Urich, F. W.:64

Vanderbilt, George:196  
 Villadolid, Deogracias V.:337  
 Vinciguerra, Decio:43  
 Viosca, Jr., Percy:199  
 Vladyskov, Vadim Dimitrij:372  
 Volsøe, Helge:220



Wagler, Erich:162  
 Wakiya, Yojiro:85  
 Walford, Lionel A.:406  
 Walker, Charles F.:241, 408  
 Wall, Frank:144, 397  
 Walters, Vladimir:262  
 Ward, Jack A.:245  
 Watson, David:304  
 Weber, Max:54  
 Weed, Alfred Cleveland:165  
 Wehrle, Richard White:58  
 Welander, Arthur Donovan:370  
 Welch, Paul S.:186  
 Weller, Alfred:163  
 Weller, W. D.:14  
 Welter, Wilfred:90  
 Werner, Franz:78  
 White, Alfred McLaren:51  
 Whitley, Gilbert P.:367  
 Wiley, Grace Olive:138  
 Williamson, E. B.:29  
 Wilson, Henry van Peters:77, 86  
 Wolterstorff, Willy:65, 299  
 Woodall, Harold:104  
 Woodward, Arthur Smith:110  
 Wright, Albert Hazen:309  
 Wright, Anna A.:308  
 Wu Hsien-Wen:249, 422  
  
 Zaret, Thomas M.:248  
 Zhu Yuangding:371

ANNUAL LIST OF BIOGRAPHIES

- 1926- Jordan, Eric Knight (306).
- 1930- Cobb, John N. (5, 380); Dybowsky, Benedikt (6); Forbes, Stephen Alfred (7); Hay, Oliver Perry (11); Hine, James S. (12); Jordan, David Starr (292, 296, 387); Kishinouye, Kamakichi (8); Metzelaar, Jan (4); Pack, Herbert J. (9); Silvester, C. F. (10); Steinbach, José (13).
- 1931- Ehrenreich, Alfred (16); Jordan, David Starr (15); Proctor, Joan (351); Stejneger, Leonhard (258, 294); Weller, W. D. (14).
- 1932- Erwin, Richard P. (21); Evermann, Barton Warren (22, 310); Johansen, A. C. (18); Klugh, A. Brooker (19); M'Intosh, W. C. (20); Patten, William (23); Titcomb, John W. (17).
- 1933- Alcock, A. W. (30); Atkinson, N. J. (25); Ayers, Howard

- (32); Johnstone, James (26); Maskell, F. G. (31); Ravenel, William de Chastignier (33); Schmidt, Johannes (27); Schulte, H. von W. (28); Starks, Edwin Chapin (24); Strecker, John Kern (420); Williamson, E. B. (29).
- 1934- Bacon, Daniel (35); Bensley, B. A. (34); Lataste, Fernand (36); Merriman, R. Owen (39); Nelson, Edward William (37); Shufeldt, Robert Wilson (38).
- 1935- Cunningham, J. T. (42); Hankinson, Thomas L. (45); Ishikawa, Chiyomatsu (40); Jenkins, Oliver Peebles (41); Knight, A. P. (46); Schmidt, Frank J. W. (44); Vinciguerra, Decio (43).
- 1936- Babcock, John Pease (48); O'Malley, Henry (47); Schrenkeisen, Ray (50); Stevenson, John Alexander (49); White, Alfred McLaren (51).
- 1937- Blanchard, Frank Nelson (376); Boulenger, George Albert (66); Clemens, Lucy Wright Smith (55); Giltay, Louis Pierre (59); Henderson, Junius (67); Hyde, Jesse Earl (52); Kampen, P. N. van (69); Longley, William Harding (53); McGregor, Richard Crittenden (56); Mosauer, Walter (57); Mukerji, Dev Dev (60); Plate, Ludwig (68); Reed, Hugh D. (61); Scott, Will (62); Stephens, Frank (63); Urich, F. W. (64); Watson, David (304); Weber, Max (54); Wehrle, Richard White (58); Wolterstorff, Willy (65).
- 1938- Artedi, Peter (352); Berg, Leo S. (388); Borodin, Nicholas A. (70); Boulenger, George Albert (395); Lowe, John N. (71).
- 1939- Embury, George Charles (74, 80); Goff, Carlos Clyde (73); Grinnell, Joseph (82, 88); Hoover, Earl E. (72); Ihering, Rodolpho von (421); Jacot, Arthur Paul (81); Kendall, William Converse (75, 79); March, Douglas D. H. (83); McMurrich, James Playfair (76, 87); Miranda Ribeiro, Alipio de (84, 356); Smith, E. Victor (89); Wakiya, Yojiro (85); Werner, Franz (78); Wilson, Henry van Peters (77, 86).
- 1940- Bass, Jr., John F. (91); Cope, Edward Drinker (269, 357); Lutz, Adolpho (93); Noble, G. K. (92); Welter, Wilfred (90).
- 1941- Clark, Howard Walton (95, 96); Lahille, Fernando (94); Smith, Hugh McCormick (260, 316, 381, 401).
- 1942- Brumwell, Malcolm J. (97); Ditmars, Raymond L. (375); Reighard, Jacob (340).
- 1943- Allen, E. J. (98); Daniel, John Franklin (99); Kelly, Howard A. (100); Regan, Charles Tate (101, 410); Roule,

- Louis (102); Snyder, John Otterbein (103, 317); Stejneger, Leonhard (259, 295); Woodall, Harold (104).
- 1944- Bromund, Fred (108); Gage, Simon Henry (109); Juday, Chauncey (106); Norman, John R. (107, 411); Townsend, Charles Haskins (105); Woodward, Arthur Smith (110).
- 1945- Awerinzew, Sergius (114); Berezovsky, A. I. (115); Goodnoh, Jr., Clifford A. (116); Morgan, Stanley Stewart (111); Reid, Earl Desmond (382); Sincock, Jr., Edwin H. (112); Soldatov, V. K. (117); Sumner, Francis B. (318); Taranetz, A. J. (113, 118).
- 1946- Barbour, Thomas (289); Boulenger, E. G. (122); Brimley, Clement Samuel (281); Dahlgren, Ulric (123); Flower, Stanley Smyth (121, 396); Nellemose, (Captain) (119); Redeke, H. C. (120); Smith, Mrs. Hugh M. (124); Taranetz, A. (125).
- 1947- Allen, W. E. (131); Allis, Jr., Edward Phelps (132); Clark, Hubert Lyman (129); Cyren, Otto (128); Eigenmann, Rosa Smith (127); Hay, William Perry (126); Surbeck, Georg (130); Wolterstorff, Willy (299).
- 1948- Bauman, Aden C. (134); Cameron, A. T. (135); Cockerell, T. D. A. (136); Eager, Grace (137); Ehrenbaum, Ernst (133); Wiley, Grace Olive (138).
- 1949- Hildebrand, Samuel Frederick (139); Tchernavin, Vladimir V. (140).
- 1950- Angel, Fernand (143); De Haas, C. O. J. (141); Hamilton, Rodgers D. (145); Hildebrand, Samuel F. (315, 319, 379, 383, 384); Lönnberg, Einar (320); McIlhenny, Edward Avery (268); Rust, Hans Theodor (146); Surber, Thaddeus (142); Wall, Frank (144).
- 1951- Andersson, Lars Gabriel (149); Angel, Fernand (307); Berg, Leo Semenovich (147, 389); Bishop, Sherman C. (150); Blegvad, Harald (155); Eddy, Brayton (148); Ewing, H. E. (152); Hoyt, Southgate Y. (153); Schäferna, Karel (151); Schmidt, Peter J. (156); Smith, George Milton (154); Wall, Frank (397).
- 1952- Barnhart, Percy S. (157); Bishop, Sherman C. (305); Engeling, Gus A. (159); Gowanloch, James Nelson (161); Lübbert, Hans (160); Starks, Chloe Leslie (358); Troemner, J. Louis (158); Wagler, Erich (162).
- 1953- Babcock, Harold Lester (350); Ellis, Max M. (164); Müller, Lorenz (313); Weller, Alfred (163).

- 1954- Duncker, Georg (166); Fassett, Harry C. (311); Gudger, E. W. (364); Johnson, Jr., J. E. (167, 377); Lantz, Louis A. (366); Weed, Alfred Cleveland (165).
- 1955- Allen, William Ray (169); Crimmins, Martin Lalor (168); Perkins, C. B. (416).
- 1956- Bertin, Léon (172); Dunn, Emmett Reid (171); Gudger, E. W. (173); Hora, Sunder Lal (392); Osburn, Raymond C. (170).
- 1957- Dunn, Emmett Reid (378); Pietschmann, Viktor (174); Schmidt, Karl Patterson (175).
- 1959- Chabanaud, Paul (177); Gordon, Myron (178, 253); Maki, Moichiro (183); Mayer, Fritz (179); Miller, Richard D. (180); Nichols, John Treadwell (297); Pfaff, J. R. (181); Schmidt, Karl Patterson (280, 369); Scofield, Norman B. (182); Seale, Alvin (314); Slevin, Joseph Richard (354); Tåning, A. Vedel (176).
- 1960- Andrews, Roy Chapman (188); Hanko, Bela (184); Limbaugh, Conrad (322); Mann, William M. (189); Miller, Richard B. (185); Schindler, Otto (321); Storey, Margaret Hamilton (190); Thienemann, August (187); Welch, Paul S. (186).
- 1961- Crawford, Sr., Donald R. (195); Curtis, Brian (191); Mann, William M. (194); Okkelberg, Peter O. (192); Roughley, T. C. (193); Storey, Margaret Hamilton (359); Vanderbilt, George (196).
- 1962- Bruun, Anton Frederik (197, 323); Buen, Fernando de (324); Herre, Albert W. (198); Viosca, Jr., Percy (199).
- 1963- Alm, Gunnar (327); Anderson, Paul (201); Burt, Charles E. (405); Cope, Edward Drinker (298); Ege, Vilhelm (325); Glauert, L. (202); Peyer, Bernhard (200); Pflueger, Al (326).
- 1964- Breyer-de Rooy, P. J. (415); Chernov, Sergius Alexandrovich (203, 348); Cook, Fannye A. (206); Grey, Marion (399); Hartweg, Norman E. (204); Needham, Paul R. (207); Snedigar, Robert (205).
- 1965- d'Ancona, Umberto (208); Barnard, K. H. (209); Clemens, Wilbert A. (210); Creaser, Charles W. (302); Davies, David H. (329); Davis, D. Dwight (211); Dymond, John Richardson (212, 385); Fox, Jr., Wade (282); Greenberg, Bernard S. (214); Howard, John K. (213); Miranda Ribeiro, Paulo de (215); Shannon, Frederick Albert (394); Thompson, William Francis (328); Wright, Anna A. (308).
- 1966- Baranov, Fedor Ilich (373); Fowler, Henry Weed (276);

- Ooster, John van (217); Reese, Albert Moore (216).
- 1967- Einarsson, Hermann (218); Michałowski, Jerzy (219); Twitty, Victor Chandler (344).
- 1968- Bigelow, Henry Bryant (285); Cochran, Doris Mable (301); Smith, James Leonard Brierley (330); Tee-Van, John (250); Volsøe, Helge (220).
- 1969- Beaufort, Lieven Ferdinand de (264); Bragg, Arthur Norris (263); Cagle, Fred Ray (407); Hass, Robert L. (222); Klauber, Laurence Monroe (390); Matsubara, Kiyomatsu (331); Mohr, Erna (332); Parker, H. W. (303); Steyn, William J. (221); Turner, Clarence Lester (333).
- 1970- Lutz, Gualter Adolpho (223); Mitchell, F. J. (224).
- 1971- Ovchynnyk, Michael M. (225); Ruthven, Alexander Grant (226); Schachter, D. (228); Shetter, David S. (334); Tanzer, Ernest C. (227); Terentjev, Paul V. (279); Wright, Albert Hazen (309).
- 1972- Frizzell, Don L. (231); Nakamura, Hiroshi (413); Sette, Oscar Elton (232); Shaw, Charles E. (229); Starrett, William C. (230); Steinitz, Heinz (347).
- 1973- Babbitt, Lewis H. (235); Eddy, Samuel (414); Landreth, Hobart (233); Peters, James A. (274); Shiraishi, Yoshikazu (335); Templeton, James R. (234).
- 1974- Hart, John L. (386); Herald, Earl Stannard (361); Hubbs, Carl Leavitt (342, 365, 391); Loftus-Hills, Jasper J. (1); Pope, Clifford H. (343); Romer, Alfred Sherwood (283); Scortecci, Giuseppe (409).
- 1975- Tanaka, Shigeho (336); Whitley, Gilbert P. (367).
- 1976- Donoso-Barros, Roberto (368); Ginsburg, Isaac (360); Günther, Klaus (290); Meade, George P. (288); Mertens, Robert (300); Villadolid, Deogracias V. (337).
- 1977- Anderson, Jr., James Donald (419); Bennett, George W. (237); Conant, Isabelle (236); Cowles, Raymond Bridgman (266); Gaige, Frederick MacMahon (254); Gaige, Helen Thompson (254); Harrington, Jr., Robert W. (251); Koumans, Frederick P. (265); Lindberg, G. U. (2); Lutz, Bertha (345); Okada, Yaichiro (339); Schroeder, William C. (284); Stoye, Frederick Hans (338).
- 1978- Camp, Charles L. (417); Greenbank, John T. (240); Hansen, Donald F. (238); Nikolsky, G. V. (398); Storer, Tracy I. (417); Taylor, Edward Harrison (239, 286).



- 1979- Gloyd, Howard Kay (271); Hubbs, Carl L. (353); Martof, Bernard S. (255); Reed, Roger J. (242); Walford, Lionel A. (406); Walker, Charles F. (241, 408).
- 1980- Logier, Eugene Bernard Shelley (278); Maria, Niceforo (404); Tinkle, Donald W. (346); Trewavas, Ethelwynn (243).
- 1981- Raney, Charlotte F. (244).
- 1982- Eaton, Theodore H. (412); Haas, Georg (418); Johnson, Richard M. (349); Welander, Arthur Donovan (370).
- 1983- Böhlke, James E. (374); Grodziński, Zygmunt (402); Rivero, Luis Howell (246); Stuart, Laurence Cooper (247, 287); Ward, Jack A. (245).
- 1984- Clay, William Marion (277); Cooper, Gerald Paul (256); Maslin, T. Paul (393); Zaret, Thomas M. (248).
- 1985- Blair, William Franklin (341); Kuehne, Robert Andrew (257); Lagler, Karl F. (355); Wu Hsien-Wen (249).
- 1986- Breder, Jr., C. M. (252); Goin, Coleman J. (273); Myers, George Sprague (272); Richardson, Sally Leonard (275); Stensiö, Erik (363); Vladykov, Vadim Dimitrij (372); Wu Hsien-Wen (422).
- 1987- Carr, Jr., Archie F. (291); Czopek, Juliusz (403); Rivas, Luis Rene (293); Rosen, Donn Eric (362); Schultz, Leonard Peter (400); Smith, Philip Wayne (267); Svetovidov, A. N. (3); Zhu Yuangding (371).
- 1988- Hollister, Gloria (261); Miller, Frances Voorhees Hubbs (270); Smith, Margaret Mary (312); Walters, Vladimir (262).

#### BIBLIOGRAPHY

1. Alexander, Richard D. 1974. Jasper J. Loftus-Hills. (3):812. N
2. Andriashev, A., and Z. Krasjukova. 1977. G. U. Lindberg, 1894-1976. (3):612-613. N, B, P
3. \_\_\_\_, E. A. Dorofeyeva, and N. V. Parin. 1987. A. N. Svetovidov, 1903-1986. (1):272-273. N, B, P
4. Anon. 1930. Jan Metzelaar, 1821-1929. (1):23-24. N, B, P
5. \_\_\_\_. 1930. John N. Cobb. (1):24. N

6. \_\_\_\_\_. 1930. Benedikt Dybowsky. (1):24. N
7. \_\_\_\_\_. 1930. Stephen Alfred Forbes. (1):24. N
8. \_\_\_\_\_. 1930. Kamakichi Kishinouye. (1):24. N
9. \_\_\_\_\_. 1930. Herbert J. Pack. (1):24. N
10. \_\_\_\_\_. 1930. C. F. Silvester. (1):24. N
11. \_\_\_\_\_. 1930. Oliver Perry Hay. (4):161. N, B
12. \_\_\_\_\_. 1930. James S. Hine. (4):162. N
13. \_\_\_\_\_. 1930. José Steinbach. (4):162. N
14. \_\_\_\_\_. 1931. W. "D." Weller. (2):72. N
15. \_\_\_\_\_. 1931. David Starr Jordan. (3):146. N
16. \_\_\_\_\_. 1931. Alfred Ehrenreich. (4):166. N
17. \_\_\_\_\_. 1932. John W. Titcomb. (1):44. N
18. \_\_\_\_\_. 1932. A. C. Johansen. (2):112. N
19. \_\_\_\_\_. 1932. A. Brooker Klugh. (2):112. N
20. \_\_\_\_\_. 1932. W. C. M'Intosh. (2):112. N
21. \_\_\_\_\_. 1932. Richard P. Erwin (3):160. N
22. \_\_\_\_\_. 1932. Barton Warren Evermann. (3):160. N
23. \_\_\_\_\_. 1932. William Patten. (4):185. N
24. \_\_\_\_\_. 1933. Edwin Chapin Starks. (1):47-48. N, B, P
25. \_\_\_\_\_. 1933. N. J. Atkinson. (1):48. N
26. \_\_\_\_\_. 1933. James Johnstone. (1):48. N
27. \_\_\_\_\_. 1933. Johannes Schmidt. (1):48. N, B
28. \_\_\_\_\_. 1933. H. von W. Schulte. (1):48. N
29. \_\_\_\_\_. 1933. E. B. Williamson. (1):48. N
30. \_\_\_\_\_. 1933. A. W. Alcock. (3):156. N
31. \_\_\_\_\_. 1933. F. G. Maskell. (3):156. N

32. \_\_\_\_\_. 1933. Howard Ayers. (4):228. N
33. \_\_\_\_\_. 1933. William de Chastignier Ravenel. (4):228. N
34. \_\_\_\_\_. 1934. B. A. Bensley. (1):60. N, B
35. \_\_\_\_\_. 1934. Daniel Bacon. (2):104. N
36. \_\_\_\_\_. 1934. Fernand Lataste. (2):104. N
37. \_\_\_\_\_. 1934. Edward William Nelson. (2):104. N, B
38. \_\_\_\_\_. 1934. Robert Wilson Shufeldt. (2):104. N
39. \_\_\_\_\_. 1934. R. Owen Merriman. (3):144. N
40. \_\_\_\_\_. 1935. Chiyomatsu Ishikawa. (1):48. N
41. \_\_\_\_\_. 1935. Oliver Peebles Jenkins. (1):48. N
42. \_\_\_\_\_. 1935. J. T. Cunningham. (2):112. N
43. \_\_\_\_\_. 1935. Decio Vinciguerra. (2):112. N
44. \_\_\_\_\_. 1935. Frank J. W. Schmidt. (3):159. N
45. \_\_\_\_\_. 1935. Thomas L. Hankinson. (4):202. N
46. \_\_\_\_\_. 1935. A. P. Knight. (4):202. N
47. \_\_\_\_\_. 1936. Henry O'Malley. (2):132. N, B
48. \_\_\_\_\_. 1936. John Pease Babcock. (3):184. N
49. \_\_\_\_\_. 1936. John Alexander Stevenson. (3):184. N
50. \_\_\_\_\_. 1936. Ray Schrenkeisen. (4):244. N
51. \_\_\_\_\_. 1936. Alfred McLaren White. (4):244. N
52. \_\_\_\_\_. 1937. Jesse Earl Hyde. (1):80. N
53. \_\_\_\_\_. 1937. William Harding Longley. (1):80. N
54. \_\_\_\_\_. 1937. Max Weber. (1):80. N
55. \_\_\_\_\_. 1937. Lucy Wright Smith Clemens. (2):148. N
56. \_\_\_\_\_. 1937. Richard Crittenden McGregor. (2):148. N
57. \_\_\_\_\_. 1937. Walter Mosauer. (2):148. N

58. \_\_\_\_\_. 1937. Richard White Wehrle. (2):148. N
59. \_\_\_\_\_. 1937. Louis Pierre Giltay. (3):200. N
60. \_\_\_\_\_. 1937. Dev Dev Mukerji. (3):200. N
61. \_\_\_\_\_. 1937. Hugh D. Reed. (3):200. N
62. \_\_\_\_\_. 1937. Will Scott. (3):200. N
63. \_\_\_\_\_. 1937. Frank Stephens. (3):200. N
64. \_\_\_\_\_. 1937. F. W. Urich. (3):200. N
65. \_\_\_\_\_. 1937. Willy Wolterstorff. (4):243. B, P
66. \_\_\_\_\_. 1937. George Albert Boulenger. (4):244. N
67. \_\_\_\_\_. 1937. Junius Henderson. (4):244. N
68. \_\_\_\_\_. 1937. Ludwig Plate. (4):244. N
69. \_\_\_\_\_. 1937. P. N. van Kampen. (4):244. N
70. \_\_\_\_\_. 1938. Nicholas A. Borodin. (1):56. N, B
71. \_\_\_\_\_. 1938. John N. Lowe. (4):210. N
72. \_\_\_\_\_. 1939. Earl E. Hoover. (1):63. N, B
73. \_\_\_\_\_. 1939. Carlos Clyde Goff. (1):63-64. N, B
74. \_\_\_\_\_. 1939. George Charles Embody. (1):64. N
75. \_\_\_\_\_. 1939. William Converse Kendall. (1):64. N
76. \_\_\_\_\_. 1939. James Playfair McMurrich. (1):64. N
77. \_\_\_\_\_. 1939. Henry Van Peters Wilson. (1):64. N
78. \_\_\_\_\_. 1939. Franz Werner. (2):122. N
79. \_\_\_\_\_. 1939. William Converse Kendall. (2):122-123. B
80. \_\_\_\_\_. 1939. George C. Embody. (2):123. B
81. \_\_\_\_\_. 1939. Arthur Paul Jacot. (2):123. N
82. \_\_\_\_\_. 1939. Joseph Grinnell. (2):124. N
83. \_\_\_\_\_. 1939. Douglas D. H. March. (2):124. N

84. \_\_\_\_\_. 1939. Alipio de Miranda Ribeiro. (2):124. N
85. \_\_\_\_\_. 1939. Yojiro Wakiya. (2):124. N, B
86. \_\_\_\_\_. 1939. Henry Van Peters Wilson. (2):124. B
87. \_\_\_\_\_. 1939. James Playfair McMurrich. (3):183. B
88. \_\_\_\_\_. 1939. Joseph Grinnell. (3):184. B
89. \_\_\_\_\_. 1939. E. Victor Smith. (4):244. N
90. \_\_\_\_\_. 1940. Wilfred Welter. (1):60. N
91. \_\_\_\_\_. 1940. John F. Bass, Jr. (2):148. N
92. \_\_\_\_\_. 1940. G. K. Noble. (4):274-275. N, B, P
93. \_\_\_\_\_. 1940. Adolpho Lutz. (4):275-276. N, B, P
94. \_\_\_\_\_. 1941. Fernando Lahille. (2):124. N
95. \_\_\_\_\_. 1941. Howard Walton Clark. (3):192. N
96. \_\_\_\_\_. 1941. Howard Walton Clark. (4):278-279. B, P
97. \_\_\_\_\_. 1942. Malcolm J. Brumwell. (1):62. N
98. \_\_\_\_\_. 1943. E. J. Allen. (1):68. N
99. \_\_\_\_\_. 1943. John Franklin Daniel. (1):68. N
100. \_\_\_\_\_. 1943. Howard A. Kelly. (1):68. N
101. \_\_\_\_\_. 1943. C. Tate Regan. (1):68. N
102. \_\_\_\_\_. 1943. Louis Roule. (1):68. N
103. \_\_\_\_\_. 1943. John Otterbein Snyder. (3):204. N
104. \_\_\_\_\_. 1943. Harold Woodall. (3):204. N
105. \_\_\_\_\_. 1944. Charles Haskins Townsend. (1):68. N
106. \_\_\_\_\_. 1944. Chauncey Juday. (1):68. N
107. \_\_\_\_\_. 1944. John R. Norman. (3):196. N
108. \_\_\_\_\_. 1944. Fred Bromund. (4):266. N
109. \_\_\_\_\_. 1944. Simon Henry Gage. (4):266. N

110. \_\_\_\_\_. 1944. Arthur Smith Woodward. (4):266. N
111. \_\_\_\_\_. 1945. Stanley Stewart Morgan. (1):59. N
112. \_\_\_\_\_. 1945. Edwin H. Sincock, Jr. (2):127. N
113. \_\_\_\_\_. 1945. A. Taranets. (3):182. N
114. \_\_\_\_\_. 1945. Sergius Awerinzew. (4):239. N (but see 1947  
(1):73.)
115. \_\_\_\_\_. 1945. A. I. Berezovsky. (4):239. N
116. \_\_\_\_\_. 1945. Clifford A. Goodnoh, Jr. (4):239. N
117. \_\_\_\_\_. 1945. V. K. Soldatov. (4):239. N
118. \_\_\_\_\_. 1945. A. J. Taranetz. (4):239. N
119. \_\_\_\_\_. 1946. Captain Nellemose. (1):54. N
120. \_\_\_\_\_. 1946. H. C. Redeke. (1):55. N
121. \_\_\_\_\_. 1946. Stanley Smythe Flower. (1):56. N
122. \_\_\_\_\_. 1946. E. G. Boulenger. (3):182. N
123. \_\_\_\_\_. 1946. Ulric Dahlgren. (3):183. N
124. \_\_\_\_\_. 1946. Mrs. Hugh M. Smith. (3):183. N
125. \_\_\_\_\_. 1946. A. Taranetz. (3):184. N
126. \_\_\_\_\_. 1947. William Perry Hay. (1):75. N
127. \_\_\_\_\_. 1947. Rosa Smith Eigenmann. (1):75-76. N, B
128. \_\_\_\_\_. 1947. Otto Cyren. (1):76. N
129. \_\_\_\_\_. 1947. Hubert Lyman Clark. (3):215. N
130. \_\_\_\_\_. 1947. Georg Surbeck. (3):216. N
131. \_\_\_\_\_. 1947. W. E. Allen. (4):294. N
132. \_\_\_\_\_. 1947. Edward Phelps Allis, Jr. (4):295. N
133. \_\_\_\_\_. 1948. Ernst Ehrenbaum. (1):75. N
134. \_\_\_\_\_. 1948. Aden C. Bauman. (1):76. N
135. \_\_\_\_\_. 1948. A. T. Cameron. (1):76. N



136. \_\_\_\_\_. 1948. T. D. A. Cockerell. (1):76. N
137. \_\_\_\_\_. 1948. Grace Eager. (1):76. N
138. \_\_\_\_\_. 1948. Grace Olive Wiley. (3):231. N
139. \_\_\_\_\_. 1949. Samuel F. Hildebrand. (1):87. N
140. \_\_\_\_\_. 1949. Vladimir V. Tchernavin. (3):236. N
141. \_\_\_\_\_. 1950. C. O. (P.) J. De Haas. (1):71. N
142. \_\_\_\_\_. 1950. Thaddeus Surber. (1):71. N
143. \_\_\_\_\_. 1950. Fernand Angel. (3):252. N
144. \_\_\_\_\_. 1950. Colonel Frank Wall. (3):252. N
145. \_\_\_\_\_. 1950. Rodgers D. Hamilton. (4):329. N
146. \_\_\_\_\_. 1950. Hans Theodor Rust. (4):329. N
147. \_\_\_\_\_. 1951. Leo S. Berg. (1):112. N
148. \_\_\_\_\_. 1951. Brayton Eddy. (1):112. N
149. \_\_\_\_\_. 1951. Lars Gabriel Andersson. (2):184. N
150. \_\_\_\_\_. 1951. Sherman C. Bishop. (2):184. N
151. \_\_\_\_\_. 1951. Karel Schäferna. (2):184. N
152. \_\_\_\_\_. 1951. H. E. Ewing. (3):256. N
153. \_\_\_\_\_. 1951. Southgate Y. Hoyt. (3):256. N
154. \_\_\_\_\_. 1951. George Milton Smith. (3):256. N
155. \_\_\_\_\_. 1951. Harald Blegvad. (4):324. N
156. \_\_\_\_\_. 1951. Peter J. Schmidt. (4):324. N
157. \_\_\_\_\_. 1952. Percy S. Barnhart. (1):58. N
158. \_\_\_\_\_. 1952. J. Louis Troemner. (1):58. N
159. \_\_\_\_\_. 1952. Gus A. Engeling. (1):59. N
160. \_\_\_\_\_. 1952. Hans Lübbert. (2):125. N
161. \_\_\_\_\_. 1952. James Nelson Gowanloch. (3):221. N

162. \_\_\_\_\_. 1952. Erich Wagler. (4):289. N
163. \_\_\_\_\_. 1953. Alfred Weller. (2):132. N
164. \_\_\_\_\_. 1953. Max M. Ellis. (4):254. N
165. \_\_\_\_\_. 1954. Alfred Cleveland Weed, 1881-1953. (2):163-164. N,  
B, L
166. \_\_\_\_\_. 1954. Georg Duncker. (2):168. N
167. \_\_\_\_\_. 1954. J. E. Johnson, Jr. (3):248. N
168. \_\_\_\_\_. 1955. Colonel Martin Lalor Crimmins. (2):158. N
169. \_\_\_\_\_. 1955. William Ray Allen. (4):325. N
170. \_\_\_\_\_. 1956. Raymond C. Osburn. (1):74. N
171. \_\_\_\_\_. 1956. Emmett Reid Dunn. (2):137-138. N
172. \_\_\_\_\_. 1956. Léon Bertin. (2):138. N
173. \_\_\_\_\_. 1956. E. W. Gudger. (2):138. N
174. \_\_\_\_\_. 1957. Viktor Pietschmann. (4):331. N
175. \_\_\_\_\_. 1957. K. P. Schmidt. (4):331. N
176. \_\_\_\_\_. 1959. A. Vedel Tåning. (1):86. N
177. \_\_\_\_\_. 1959. Paul Chabanaud. (2):184. N
178. \_\_\_\_\_. 1959. Myron Gordon. (2):184-185. N
179. \_\_\_\_\_. 1959. Fritz Mayer. (2):185. N
180. \_\_\_\_\_. 1959. Richard D. Miller. (2):185. N
181. \_\_\_\_\_. 1959. J. R. Pfaff. (2):185. N
182. \_\_\_\_\_. 1959. Norman B. Scofield. (2):185. N
183. \_\_\_\_\_. 1959. Moichiro Maki. (3):272. N
184. \_\_\_\_\_. 1960. Bela Hanko. (2):164. N
185. \_\_\_\_\_. 1960. Richard B. Miller. (2):164. N
186. \_\_\_\_\_. 1960. Paul S. Welch. (2):164. N

187. \_\_\_\_\_. 1960. August Thienemann. (4):391. N
188. \_\_\_\_\_. 1960. Roy Chapman Andrews. (4):395. N
189. \_\_\_\_\_. 1960. William M. Mann. (4):395. N
190. \_\_\_\_\_. 1960. Margaret H. Storey. (4):395. N
191. \_\_\_\_\_. 1961. Dr. Brian Curtis. (2):260. N
192. \_\_\_\_\_. 1961. Dr. Peter O. Okkelberg. (2):260. N
193. \_\_\_\_\_. 1961. Dr. T. C. Roughley. (2):261. N
194. \_\_\_\_\_. 1961. William M. Mann. (3):368 N
195. \_\_\_\_\_. 1961. Donald R. Crawford, Sr. (4):519. N
196. \_\_\_\_\_. 1961. George Vanderbilt. (4):519. N
197. \_\_\_\_\_. 1962. A. F. Bruun. (1):251. N
198. \_\_\_\_\_. 1962. Albert W. Herre. (1):251. N
199. \_\_\_\_\_. 1962. Percy Viosca, Jr. (1):251. N
200. \_\_\_\_\_. 1963. Bernhard Peyer. (1):223. N
201. \_\_\_\_\_. 1963. Paul Anderson. (2):472. N, B
202. \_\_\_\_\_. 1963. Dr. L. Glauert. (4):723. N
203. \_\_\_\_\_. 1964. Dr. S. A. Chernov. (1):255. N
204. \_\_\_\_\_. 1964. Norman E. Hartweg. (1):256. N
205. \_\_\_\_\_. 1964. Robert Snedigar. (1):256. N, B
206. \_\_\_\_\_. 1964. Fannye A. Cook. (3):598. N
207. \_\_\_\_\_. 1964. Paul R. Needham. (3):598. N
208. \_\_\_\_\_. 1965. Umberto d'Ancona. (1):124. N
209. \_\_\_\_\_. 1965. Dr. K. H. Barnard. (1):124. N
210. \_\_\_\_\_. 1965. Wilbert A. Clemens. (1):124. N
211. \_\_\_\_\_. 1965. D. Dwight Davis. (1):124. N, B
212. \_\_\_\_\_. 1965. John Richardson Dymond. (1):124. N

213. \_\_\_\_\_. 1965. Colonel John K. Howard. (2):262. N, B
214. \_\_\_\_\_. 1965. Bernard S. Greenberg. (3):399. N, B
215. \_\_\_\_\_. 1965. Paulo de Miranda Ribeiro. (4):543. N
216. \_\_\_\_\_. 1966. Albert Moore Reese. (2):380. N
217. \_\_\_\_\_. 1966. John Van Ooster. (2):381-382. N
218. \_\_\_\_\_. 1967. Dr. Hermann Einarsson. (2):503. N
219. \_\_\_\_\_. 1967. Jerzy Michałowski. (2):503. N
220. \_\_\_\_\_. 1968. Dr. Helge Volsøe. (3):663. N
221. \_\_\_\_\_. 1969. Dr. William J. Steyn. (2):422-423. N
222. \_\_\_\_\_. 1969. Dr. Robert L. Hass. (2):423. N
223. \_\_\_\_\_. 1970. Gualter Adolpho Lutz. (3):598. N
224. \_\_\_\_\_. 1970. F. J. Mitchell. (3):598. N
225. \_\_\_\_\_. 1971. Michael M. Ovchynnyk. (1):194. N
226. \_\_\_\_\_. 1971. Alexander Grant Ruthven (1882-1971). (3):587. N
227. \_\_\_\_\_. 1971. Tanzer Memorial Fund (Ernest C. Tanzer). (3):588.  
N
228. \_\_\_\_\_. 1971. Madam D. Schachter. (4):768. N
229. \_\_\_\_\_. 1972. Charles E. Shaw. (1):206. N
230. \_\_\_\_\_. 1972. William C. Starrett. (1):206. N
231. \_\_\_\_\_. 1972. Don L. Frizzell. (4):906. N
232. \_\_\_\_\_. 1972. Oscar Elton Sette. (4):906. N
233. \_\_\_\_\_. 1973. Hobart Landreth. (3):639. N
234. \_\_\_\_\_. 1973. James R. Templeton. (3):639. N
235. \_\_\_\_\_. 1973. Lewis H. Babbitt. (4):831. N
236. \_\_\_\_\_. 1977. Isabelle Conant. (1):209-210. N, B
237. \_\_\_\_\_. 1977. George W. Bennett. (3):614. N
238. \_\_\_\_\_. 1978. Donald F. Hansen. (2):380. N

239. \_\_\_\_\_. 1978. Edward Harrison Taylor. (3):557. N
240. \_\_\_\_\_. 1978. John T. Greenbank. (4):740. N
241. \_\_\_\_\_. 1979. Charles F. Walker. (2):382. N
242. \_\_\_\_\_. 1979. Roger J. Reed. (4):757. N
243. \_\_\_\_\_. 1980. Ethelwynn Trewavas. (4):952. B
244. \_\_\_\_\_. 1981. Charlotte F. Raney. (3):749. N
245. \_\_\_\_\_. 1983. Jack A. Ward. (1):285. N
246. \_\_\_\_\_. 1983. Luis Howell Rivero. (2):582-583. N
247. \_\_\_\_\_. 1983. Lawrence C. Stuart. (3):853-854. N
248. \_\_\_\_\_. 1984. Thomas M. Zaret. (4):1042. N
249. \_\_\_\_\_. 1985. Wu Hsien-Wen. (4):1087. N
250. Atz, James W. 1968. John Tee-Van, July 6, 1897-November 5, 1967. (3)660-661. N, B
251. \_\_\_\_\_. 1977. Robert W. Harrington, Jr., 1911-1975. (1):204-205. N, B, P
252. \_\_\_\_\_. 1986. C. M. Breder, Jr., 1897-1983. (3):853-856. N, B, P
253. \_\_\_\_\_, and Donn E. Rosen. 1959. Myron Gordon, 1899-1959. (4):352-354. B, P
254. Bailey, Joseph R. 1977. Helen Thompson Gaige; Frederick MacMahon Gaige, 1890-1976. (3):609-611. N, B, P
255. \_\_\_\_\_. 1979. Bernard S. Martof. (2):380-381. N, B
256. Bailey, Reeve M. 1984. Gerald Paul Cooper. (4):1042-1043. N, B, P
257. Barbour, Roger W. 1985. Robert Andrew Kuehne, 1927-1984. (2):528-529. N, B, P
258. Barbour, Thomas. 1931. To Leonhard Stejneger. (3):73. P
259. \_\_\_\_\_. 1943. Leonhard Stejneger. (1):1. N, P
260. Bartsch, Paul. 1941. Dr. Hugh M. Smith, director of the Philippine Cruise of the "Albatross." (4):209-215. B

261. Berra, Tim M. 1988. Gloria Hollister, 1900-1988. (4):1113. N, B, P
262. Berry, Fred. 1988. Vladimir Walters, 1927-1987. (4):1112-1113. N, B, P
263. Black, Jeffrey Howard, and Charles C. Carpenter. 1969. Arthur Norris Bragg, 18 December 1897-27 August 1968. (2):419-420. N, B, P
264. Boeseman, M. 1969. Lieven Ferdinand de Beaufort, 1879-1968. (1):221-222. N, B, P
265. \_\_\_\_\_. 1977. Frederick P. Koumans, 1905-1977. (4):802. N, B, P
266. Brattstrom, Bayard H. 1977. Raymond Bridgman Cowles, 1896-1975. (3):611-612. N, B, P
267. Burr, Brooks M., and Lawrence M. Page. 1987. Philip Wayne Smith, 1921-1986. (3):839-840. N, B, P
268. Cagle, Fred R. 1950. Edward Avery McIlhenny. (3):245-246. N, B
269. Case, E. C. 1940. Cope- The man. (2):61-65. B, P
270. Chernoff, Barry. 1988. Frances Voorhees Hubbs Miller, 1919-1987. (2):520-523. N, B, P
271. Clay, William M. 1979. Howard Kay Gloyd (1902-1978). (1):187-189. N, B, P
272. Cohen, Daniel M., and Stanley H. Weitzman. 1986. George Sprague Myers, 1905-1985. (3):851-853. N, B, P
273. Colbert, Edwin H. 1986. Coleman J. Goin, 1911-1986. (4):1041-1043. N, B, P
274. Collette, Bruce B. 1973. James A. Peters. (2):388-390. N, B, P
275. \_\_\_\_\_. 1986. Sally Leonard Richardson, 1944-1986. (4):1043-1045. N, B, P
276. Conant, Roger. 1966. Henry Weed Fowler, 1878-1965. (3):628-629. N, B, P
277. \_\_\_\_\_. 1984. William Marion Clay, 1906-1983. (2):563. N, B
278. Crossman, E. J. 1980. Eugene Bernard Shelley Logier. (3):572-574. N, B, P

279. Darevsky, Ilya S., and Leo I. Khosatzky. 1971. Paul V. Terentjev (1903-1970). (2):382-384. N, B, P
280. Davis, D. Dwight. 1959. Karl Patterson Schmidt, 1890-1957. (3):188-192. B, P
281. Davis, Harry T. 1946. Clement Samuel Brimley. (3):182. N, B
282. Dessauer, H. C. 1965. Wade Fox, Jr., 1920-1964. (1):123. N, B, P
283. Dick, Myvanwy M. 1974. Alfred Sherwood Romer, 1884-1973. (1):293-294. N, B
284. \_\_\_\_\_. 1977. William C. Schroeder. 1894-1977. (2):412-413. N, B
285. \_\_\_\_\_, and William C. Schroeder. 1968. Henry Bryant Bigelow, 1879-1967. (3):657-659. N, B, P
286. Duellman, William E. 1978. Edward Harrison Taylor, 1889-1978. (4):737-738. N, B, P
287. \_\_\_\_\_. 1983. Laurence Cooper Stuart. (4):1116-1118. B, P
288. Dundee, Harold A. 1976. George P. Meade. (1):220-221. N, B
289. Dunn, Emmett R. 1946. Thomas Barbour, 1884-1946. (1):1-3. N, B, P
290. Ebeling, A. W. 1976. Klaus Günther, 1907-1975. (4):845. N, B
291. Ehrenfeld, David. 1987. Archie F. Carr, Jr., 1909-1987. (4):1087-1089. N, B, P
292. Evermann, Barton Warren. 1930. David Starr Jordan, the man. (4):93-105. B, P
293. \_\_\_\_\_. Fink, William L. 1987. Luis Rene Rivas, 1916-1986. (1):269-271. N, B, P
294. Fisher, Albert Kendrick. 1931. Leonhard Stejneger. (3):74-83. B, P
295. \_\_\_\_\_. 1943. Leonhard Stejneger. (3):137-141. B
296. Fowler, Henry W. 1930. David Starr Jordan and ichthyology in the Pacific. (4):108-109. B
297. \_\_\_\_\_. 1959. John Treadwell Nichols, 1883-1958. (1):83. N, B

298. \_\_\_\_\_. 1963. Cope in retrospect. (1):195-198. B
299. Freytag, Gunther. 1947. Dr. Willy Wolterstorff. (3):215. N, B
300. Gans, Carl. 1976. Robert Mertens, 1894-1975. (2):420. N, B
301. Goin, Coleman J. 1968. Doris Mable Cochran, May 18, 1898-May 22, 1968. (3):661-662. N, B, P
302. Gorbman, Aubrey. 1965. Charles W. Creaser. (3):398-399. N, B
303. Grandison, Alice G. C. 1969. H. W. Parker. (2):416-417. N, B, P
304. Gregory, W. K. 1937. David Watson. (3):197. B, P
305. Grobman, Arnold B. 1952. Sherman C. Bishop, 1887-1951. (3):127-128. B, P
306. Guerard, Albert. 1926. Eric Knight Jordan, 1903-1926. (152):suppl. N, B
307. Guibe, J. 1951. Fernand Angel, 1881-1950. (1):1-2. B, P
308. Hamilton, Jr., W. J. 1965. Mrs. Anna A. Wright. (1):124. N, B
309. \_\_\_\_\_. 1971. Albert Hazen Wright, August 15, 1879-July 4, 1970. (2):381-382. N, B, P
310. Hanna, G Dallas. 1932. Barton Warren Evermann, 1853-1932. (4):161-162. B, P
311. \_\_\_\_\_. 1954. Harry C. Fassett. (2):167. N
312. Heemstra, Phillip C., and Michael N. Bruton. 1988. Margaret Mary Smith, 1916-1987. (2):523-525. N, B, P
313. Hellmich, Walter. 1953. Lorenz Müller, 1868-1953. (3):133-134. N, B, P
314. Herre, Albert W. 1959. Alvin Seale, 1871-1958. (1):85-86. N, B
315. Higgins, Elmer. 1950. Samuel F. Hildebrand as a government scientist. (1):8-11. B
316. Hildebrand, Samuel F. 1941. Hugh McCormick Smith and the Bureau of Fisheries. (4):216-220. B, P
317. Hubbs, Carl L. 1943. John O. Snyder. (4):265-266. B, P



318. \_\_\_\_\_. 1945. Francis B. Sumner, 1874-1945. (4):183-184. N, B,  
P
319. \_\_\_\_\_. 1950. Samuel F. Hildebrand, ichthyologist and  
herpetologist. (1):12-14. B, P
320. \_\_\_\_\_. 1950. Einar Lönnberg. (1):70. N
321. \_\_\_\_\_. 1960. Dr. Otto Schindler. (3):269. N, B
322. \_\_\_\_\_. 1960. Conrad Limbaugh. (3):269-270. N, B, P
323. \_\_\_\_\_. 1962. Anton Frederick Bruun. (2):481-482. B, P
324. \_\_\_\_\_. 1962. Fernando de Buen. (4):875-876. N, B, P
325. \_\_\_\_\_. 1963. Vilhelm Ege. (1):220-222. N, B, P, L
326. \_\_\_\_\_. 1963. Al Pflueger. (1):223. N
327. \_\_\_\_\_. 1963. Gunnar Alm. (1):224. N
328. \_\_\_\_\_. 1965. William Francis Thompson. (4):542. N, B
329. \_\_\_\_\_. 1965. David H. Davies. (4):543. N, B
330. \_\_\_\_\_. 1968. James Leonard Brierley Smith, 1897-1968. (3):659-  
660. N, B, P
331. \_\_\_\_\_. 1969. Kiyomatsu Matsubara, 1907-1968. (2):420-422. N,  
B, P
332. \_\_\_\_\_. 1969. Erna Mohr, July 11, 1894-September 10, 1968.  
(3):646. N, B, P
333. \_\_\_\_\_. 1969. Clarence Lester Turner, 1890-1969. (3):646-647.  
N, B
334. \_\_\_\_\_. 1971. David S. Shetter, 1910-1970. (1):194. N
335. \_\_\_\_\_. 1973. Yoshikazu Shiraishi. (2):393-394. N, B
336. \_\_\_\_\_. 1975. Shigeho Tanaka, 1878-1974. (4):792. N, B, P
337. \_\_\_\_\_. 1976. Deogracias V. Villadolid, 1896-1976. (3):617. N
338. \_\_\_\_\_. 1977. Frederick Hans Stoye, 1887-1977. (3):615-616. N,  
B, P
339. \_\_\_\_\_, and Katsuzo Kuronuma. 1977. Yaichiro Okada, 1892-1976.  
(1):206-207. N, B, P, L

340. \_\_\_\_, and Karl F. Lagler. 1942. Jacob Reighard, 1861-1942.  
(1):63-64. N, B, P
341. Hubbs, Clark. 1985. William Franklin Blair, 1912-1984.  
(2):529-531. N, B, P
342. Hubbs, Laura C. 1974. List of students whose PhD degree was  
under the chairmanship of Carl L. Hubbs through 1974.  
(3):609-610.
343. Inger, Robert F. 1974. Clifford H. Pope. (4):1012. N, B
344. Kelly, Douglas E. 1967. Victor Chandler Twitty, November 5,  
1901-March 22, 1967. (4):887-888. N, B, P
345. Kennedy, J. P. 1977. Bertha Lutz, 1874-1976. (1):208-209. N,  
B
346. Kluge, Arnold. 1980. Donald W. Tinkle (1930-1980). (3):572.  
N, B, P
347. Kuehne, Robert A. 1972. Dr. Heinz Steinitz, 1909-1971.  
(3):609-610. N, B, P
348. Leviton, Alan E. 1964. Sergius Alexandrovich Chernov, 1903-  
1964. (2):466. B, P
349. Liner, Ernest A. and Allan H. Chaney. 1982. Richard M.  
Johnson. (4):987. N, B
350. Loveridge, A. 1953. Harold Lester Babcock, 1886-1953.  
(3):134-135. N, B
351. Mann, William M. 1931. Joan Proctor. (4):166. N, B
352. Merriman, Daniel. 1938. Peter Artedi-Systematist and  
ichthyologist. (1):33-39. B
353. Miller, Frances H., Clark Hubbs, and Earl L. Hubbs. 1979.  
Carl L. Hubbs. (4):756. N, P
354. Miller R. C. 1959. Joseph Richard Slevin, 1881-1957. (1):84-  
85. N, B, P
355. Miller, Robert Rush. 1985. Karl F. Lagler, 1912-1985.  
(4):1090-1091. N, B, P
356. Myers, George S. 1939. Alipio de Miranda Ribeiro. (3):184. B
357. \_\_\_\_. 1940. Cope as an ichthyologist. (2):76-78. B, P

358. \_\_\_\_\_. 1952. Chloe Leslie Starks. (2):124-125. N
359. \_\_\_\_\_. 1961. Margaret Hamilton Storey (1900-1960). (2):261-263. B, P, L
360. \_\_\_\_\_. 1976. Isaac Ginsburg. (1):217-219. N, B, P
361. \_\_\_\_\_, W. I. Follett, and William A. Gosline. 1974. Earl Stannard Herald. (1):291-292. N, B, P
362. Nelson, Gareth, James W. Atz, Klaus D. Kallman, and C. Lavett Smith. 1987. Donn Eric Rosen, 1929-1986. (2):541-546. N, B, P, L
363. \_\_\_\_\_, and Mee-Man Chang. 1986. Erik Stensiö, 1891-1984. (2):558. N, B, P
364. Nichols, John T. 1954. E. W. Gudger. (2):164-165. B
365. Norris, Kenneth S. 1974. To Carl Leavitt Hubbs, a modern pioneer naturalist on the occasion of his eightieth year. (3):581-594. B, P
366. Parker, H. W. 1954. Louis A. Lantz. (1):79. N, B
367. Paxton, John R., and Douglass F. Hoese. 1975. Gilbert P. Whitley, 1903-1975. (4):792-793. N, B, P
368. Pefaur, Jaime E. 1976. Roberto Donoso-Barros. (1):219-220. N, B, P
369. Peters, James A. 1959. A bibliography and index of Karl P. Schmidt's papers on coral snakes. (3):192-196. L
370. Pietsch, Theodore W. 1982. Arthur Donovan Welander (1908-1982). (3):737. N, B
371. Qingwen, Meng. 1987. Zhu Yuangding (Yuanting T. Chu), 1896-1986. (3):838-839. N, B, P
372. Renaud, Claude B. 1986. Vadim Dimitrij Vladykov. (2):559-560. N, B, P
373. Ricker, W. E. 1966. Fedor Ilich Baranov. (1):147. N
374. Robins, C. Richard. 1983. James E. Böhlke. (3):850-851. N, B, P
375. Ruthven, Alexander G. 1942. Raymond L. Ditmars. (2):131. N, B
376. Schmidt, Karl P. 1937. Frank Nelson Blanchard, 1888-1937.

- (3):149-150. N, B, P
377. \_\_\_\_\_. 1954. J. E. Johnson, Jr.- An appreciation. (3):247-248.  
B
378. \_\_\_\_\_. 1957. Emmett Reid Dunn, 1894-1956. (2):74-77. B, P
379. Schmitt, Waldo L. 1950. Samuel Frederick Hildebrand. (1):1.  
P
380. Schultz, Leonard P. 1930. John N. Cobb, Dean of the College  
of Fisheries. (2):50-51. B
381. \_\_\_\_\_. 1941. Hugh McCormick Smith. (4):194-209. N, B, P, L
382. \_\_\_\_\_. 1945. Earl Desmond Reid Retires. (2):128. B
383. \_\_\_\_\_. 1950. Samuel Frederick Hildebrand. (1):2-7. B, P
384. \_\_\_\_\_, and Ann Shepard Green. 1950. Bibliography of Samuel F.  
Hildebrand. (1):15-18. L
385. Scott, W. B. 1965. John Richardson Dymond, 1887-1965.  
(3):396-398. B, P
386. \_\_\_\_\_. 1974. John L. Hart, 1904-1973. (2):579. N, B
387. Seale, Alvin. 1930. To David Starr Jordan. (4):106-107. B, P
388. Shapovalov, Leo. 1938. Leo S. Berg. (1):55. B, P
389. \_\_\_\_\_. 1951. Leo Semenovich Berg, 1876-1950. (3):185-186. B, P
390. Shaw, Charles E. 1969. Laurence Monroe Klauber, 1883-1968.  
(2):417-419. N, B, P
391. Shor, Elizabeth N. 1974. Selected bibliography of Carl  
Leavitt Hubbs from 1915 to 1974. (3):594-609. L
392. Silas, E. G. 1956. Sunder Lal Hora. (2):134-136. N, B, P
393. Smith, Hobart M. 1984. T. Paul Maslin, 1909-1984. (3):806-  
807. N, B, P
394. \_\_\_\_\_, and Findlay E. Russell. 1965. Frederick Albert Shannon,  
1921-1965. (4):541-542. N, B, P
395. Smith, Malcolm. 1938. George Albert Boulenger, 1858-1937.  
(1):1-3. B, P
396. \_\_\_\_\_. 1946. Stanley Smyth Flower, 1871-1946. (4):185-187. B,  
P

397. \_\_\_\_\_. 1951. Frank Wall, 1868-1950. (2):113-114. B, P
398. Soin, S. G., and M. V. Mina. 1978. G. V. Nikolsky (1910-1977). (2):379. N, B, P
399. Sonoda, Pearl M., and Robert F. Inger. 1964. Marion Grey. (3):598. N, B
400. Springer, Victor G. 1987. Leonard Peter Schultz, 1901-1986. (1):271-272. N, B, P
401. Stejneger, Leonhard. 1941. To Hugh McCormick Smith. (4):193. P
402. Szarski, Henryk. 1983. Zygmunt Grodziński. (3):851-852. N, B, P
403. \_\_\_\_\_. 1987. Juliusz Czopek, 1922-1986. (3):839. N
404. Tamsitt, J. R. 1980. Brother Niceforo Maria. (4):952-953. N, B
405. Tanner, Wilmer W. 1963. Charles E. Burt. (4):722-723. N, B, P
406. Thompson, Paul E. 1979. Lionel A. Walford. (3):564. N
407. Tinkle, Donald W. 1969. Fred Ray Cagle, 1915-1968. (1):220. N, B, P
408. \_\_\_\_\_. 1979. Charles F. Walker. (3):563-564. B, P
409. Tortonese, Enrico. 1974. Prof. Giuseppe Scortecchi, 1898-1973. (1):294. N, B
410. Trewavas, Ethelwynn. 1943. Charles Tate Regan. (3):202-204. B, P
411. \_\_\_\_\_. 1944. John R. Norman. (4):265-266. B, P
412. Trueb, Linda. 1982. Theodore H. Eaton. (1):245. N, B
413. Ueyanagi, Shoji. 1972. Dr. Hiroshi Nakamura, 1906-1971. (1):204-205. N, B, P
414. Underhill, James. 1973. Samuel Eddy. (2):392-393. N, B, P
415. Van der Feen-van Benthem Jutting, W. S. S. 1964. Mrs. Dr. P. J. Breyer-de Rooy. (3):598. N, B

416. Wade, Wallace B. 1955. C. B. Perkins, 1889-1955. (4):323-324. N, B
417. Wake, David B. 1978. Tracy I. Storer and Charles L. Camp. (1):196-197. B
418. Werner, Yehudah L. 1982. Georg Haas, 1905-1981. (2):491-493. N, B, P
419. Wilhoft, Daniel C., and Keith A. Hawthorne. 1977. James Donald Anderson, Jr. (2):412. N, B
420. Williams, Walter J. 1933. John Kern Strecker. (1):48. N, B
421. Wright, Stillman. 1939. Rodolpho von Ihering. (4):244. B
422. Yiyu, Chen. 1986. Wu Hsien-wen, 1900-1985. (2):557-558. B, P














A KEY AND CHECKLIST  
TO THE  
NEOTROPICAL SNAKE GENUS LIOPHIS  
WITH COUNTRY LISTS AND MAPS



JAMES R. DIXON

Department of Wildlife and Fisheries Sciences  
Texas A&M University



SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
NO. 79

1989

SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with us.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560, U.S.A. Please include a self-addressed mailing label with requests.

## INTRODUCTION

The genus Liophis currently contains 35 species and 62 recognized subspecies. About 137 names have been proposed, attesting to extensive qualitative and quantitative variation between and among the species of Liophis. The purpose of the checklist, keys, and country lists is to offer museum curators and herpetologists a means to identify the species and subspecies of one of the most commonly encountered xenodontine snake genera of the Neotropics. Species lists are constructed from specimens personally examined and verified by me. The keys resulted from an analysis of variation of 5,198 specimens of all recognized populations.

Some species have been recognized since 1758 and others as recently as 1987. Some species are represented in collections by only a few specimens (L. atraventer, L. problematicus), by several hundred (most species), or over a thousand (L. miliaris, L. poecilogyrus). Taxonomic data are generally adequate for most taxa.

The checklist contains the accepted name and its author, followed by primary synonymies, their author(s), date, page number, and type locality. The primary synonymies are followed by the author, date, and page number of the first proper usage of the epithet, if necessary for clarification. Species are arranged alphabetically. Subspecies are also arranged alphabetically under each species, except for the nominate race, which is placed first. Synonymies for subspecies follow the accepted name, arranged in alphabetical order, followed by the author(s) name(s). A statement of distribution follows the synonymy of each species and subspecies, along with a citation denoting a publication with a distribution map.



KEY TO SPECIES OF LIOPHIS

- |      |  |                      |
|------|--|----------------------|
| 1.a  | Posterior dorsal scale rows at least two less than at midbody .....  | 5                    |
| b    | Dorsal scale rows without reduction .....  | 2                    |
| 2.a  | Dorsal scale rows 15-15-15 .....   | 3                    |
| b    | Dorsal scale rows 17-17-17 .....   | 4                    |
| 3.a  | Three supralabials entering orbit .....  | <u>andinus</u>       |
| b    | Two supralabials entering orbit. <u>reginae</u> (=oligolepis of others)  |                      |
| 4.a  | Eight supralabials (rarely seven); dorsum olive green with or without reddish vertebral stripe and small dorsolateral black spots .....      | <u>jaegeri</u>       |
| b    | Seven supralabials; dorsum tan or brown with darker blotches and four blackish posterior lines, and a black edged, white labial stripe ..... | <u>williamsi</u>     |
| 5.a  | Nineteen midbody dorsal scale rows .....   | 29                   |
| b    | Seventeen midbody dorsal scale rows .....  | 6                    |
| 6.a  | Dorsal scale rows 17-17-15 .....   | 7                    |
| b    | Dorsal scale rows 17-17-13 .....   | <u>flavifrenatus</u> |
| 7.a  | Seven supralabials .....   | 8                    |
| b    | Eight supralabials .....   | 9                    |
| 8.a  | Light dorsal bands not widened laterally into broad triangles; $\leq 17$ maxillary teeth .....   | <u>breviceps</u>     |
| b    | Light dorsal bands distinctly widened laterally into broad triangles; $\geq 20$ maxillary teeth .....  | <u>longiventris</u>  |
| 9.a  | Venter with checkered pattern of black and red or yellow .....   | 10                   |
| b    | Venter never checkered with black and red or yellow, but occasionally with black marks on lateral edges of ventrals .....                    | 19                   |
| 10.a | Black lateral posterior stripe present and extends onto tail .....   | 11                   |
| b    | Not as above .....   | 13                   |
| 11.a | Apical scale pit present .....   | 12                   |
| b    | No scale pit .....   | <u>taeniurus</u>     |
| 12.a | Posterior lateral black stripe occurs as an edge between 3rd and 4th scale rows .....  | <u>reginae</u>       |
| b    | Posterior lateral black stripe occurs over most of 3rd, one-third to all of 4th and occasionally on 5th scale row .....                      | <u>epinephelus</u>   |
| 13.a | One preocular .....  | 14                   |
| b    | Two preoculars .....   | <u>festae</u>        |



|      |   |                  |
|------|---|------------------|
| 14.a | Ventrals $\leq 179$ .....   | 15               |
| b    | Ventrals $\geq 186$ ..... <u>frenatus</u>   |                  |
| 15.a | Apical scale pit present, tail length $\geq 26\%$ of total length .....   | 16               |
| b    | Scale pit absent, tail length $\leq 23\%$ of total length .....   | 17               |
| 16.a | Subcaudals $\geq 74$ ; $\geq 26$ maxillary teeth ..... <u>juliae</u>  |                  |
| b    | Subcaudals $\leq 72$ ; $\leq 27$ maxillary teeth ..... <u>epinephelus</u>   |                  |
| 17.a | Banded dorsal pattern, occasionally a combination bands and reticulations; ventrals and subcaudals always checkered with black and red or yellow .....  | 18               |
| b    | Salt and pepper dorsal pattern, obscure in Amazon populations; venter light with some dark edging in all populations except Amazon, where large black checkered marks are present; Amazon population usually with dusky throats and white subcaudals, subcaudals of other populations white or dark edged ..... |                  |
|      | ..... <u>miliaris</u>   |                  |
| 18.a | Light body bands widened laterally into broad triangles .....   |                  |
|      | ..... <u>longiventris</u>   |                  |
| b    | Light body bands of equal width .....   | <u>cobellus</u>  |
| 19.a | Lateral edge of ventrals unmarked .....   | 20               |
| b    | Lateral edge of ventrals marked with black .....  | 25               |
| 20.a | Dorsum with or without posterior lateral black stripes; $\geq 13$ maxillary teeth .....   | 21               |
| b    | Dorsum with three dark lines from head to tail; $\leq 13$ maxillary teeth ..... <u>paucidens</u>  |                  |
| 21.a | Posterior lateral black stripe present on body and tail .....   | 22               |
| b    | Stripe absent .....   | 23               |
| 22.a | Nine infralabials; $\leq 21$ maxillary teeth; tail length about 22% of total length .....   | <u>melanotus</u> |
| b    | Ten infralabials; $\geq 21$ maxillary teeth; tail length about 26% of total length .....  | <u>reginae</u>   |
| 23.a | Subcaudals $\geq 74$ ; dorsum uniform color or variable, but never green or olive green .....   | 24               |
| b    | Subcaudals $\leq 73$ ; dorsum olive green or green, with or without reddish dorsal stripe and small dorsolateral black spots .....  | <u>jaegeri</u>   |
| 24.a | Ventrals $\leq 167$ .....   | <u>juliae</u>    |
| b    | Ventrals $\geq 182$ .....   | <u>perfuscus</u> |

|      |  |                      |
|------|--|----------------------|
| 25.a | Dorsum without dark stripes .....  | 26                   |
| b    | Dorsum with dark stripes .....   | 28                   |
| 26.a | Tail length $\geq 26\%$ of total length .....  | 27                   |
| b    | Tail length $\leq 23\%$ of total length ..... <u>miliaris</u>  |                      |
| 27.a | Subcaudals $\leq 93$ .....   | <u>ornatus</u>       |
| b    | Subcaudals $\geq 96$ .....   | <u>cursor</u>        |
| 28.a | Dorsum with three dark and two yellow stripes, yellow stripes begin on snout .....   | <u>flavifrenatus</u> |
| b    | Dorsum with three to five dark stripes, yellow stripes absent .....  | <u>triscalis</u>     |
| 29.a | Dorsal scale rows 19-19-17 .....   | 30                   |
| b    | Dorsal scale rows 19-19-15 or 19-19-13 .....   | 39                   |
| 30.a | Dorsum uniform green or olive green, occasionally with an ill-defined reddish brown mid-dorsal stripe .....  | 31                   |
| b    | Dorsum variable, but never uniform or with an ill-defined reddish mid-dorsal stripe .....  | 33                   |
| 31.a | Ventrals $\leq 159$ .....  | 32                   |
| b    | Ventrals $\geq 169$ .....  | <u>viridis</u>       |
| 32.a | Venter rose or red; $\geq 62$ subcaudals; $\geq 25$ maxillary teeth .....  | <u>maryellenae</u>   |
| b    | Venter black; $\leq 56$ subcaudals; $\leq 24$ maxillary teeth .....  | <u>atraventer</u>    |
| 33.a | One preocular .....  | 34                   |
| b    | Two preoculars .....   | <u>festae</u>        |
| 34.a | Dorsum uniform brown or with two to five dark lines on a tan to light brown ground color .....   | 35                   |
| b    | Dorsum gray or olive, with or without dark irregular marks scattered throughout; or dorsum brown to dark brown with large blotches or bands anteriorly ..... | 37                   |
| 35.a | Dorsum tan or light brown with three to five distinct dark stripes, the median stripe beginning on the snout or the nape .....                               | 36                   |
| b    | Dorsum uniform brown, with or without dark freckles and a dark ventrolateral line .....  | <u>sagittifer</u>    |

- 36.a Dorsum tan or light brown with three distinct black lines, the median line beginning on the snout,  $\geq 159$  ventrals;  $\geq 77$  subcaudals; tail/total length ratio (males)  $\geq .24$  ..... lineatus
- b Dorsum light brown with a broad dark median stripe, narrow dark dorsolateral stripes, and dark ventrolateral stripes; median stripe begins on the nape; ventrals 133, subcaudals 36, tail/total length ratio .185 (known only from the male holotype) ..... problematicus
- 37.a Dorsum gray or tan with darker blotches, no black dorsolateral line posteriorly; faint to moderately distinct dorsolateral light line posteriorly; dorsal surface of head gray, tan or brown with or without a whitish U,V,X, or Y mark on the parietals with exterior black edging, the mark may extend anteriorly to the internasals; with or without one apical scale pit ..... 38
- b Dorsum dark brown or chocolate brown, banded or blotched anteriorly with contrasting dorsolateral black and white lines posteriorly; no apical scale pit ..... taeniurus
- 38.a Dorsal surface of frontal and parietals of head gray, brown, or olive, with or without a few darker flecks; no apical scale pit present (characters from type description) ..... carajasensis
- b Dorsal surface of frontal and parietal scales of head gray, brownish or live, with a whitish U, V, X, or Y mark with exterior black edging, mark may extend onto the internasals; one apical scale pit present ..... almadensis
- 39.a Dorsal scale rows 19-19-15 ..... 40
- b Dorsal scale rows 19-19-13 ..... perfuscus
- 40.a Dorsum bright green; head occasionally blue, with or without dark chevrons dorsally and/or laterally ..... 41
- b Dorsum variable but never green ..... 42
- 41.a Ventrals  $\leq 168$  ..... typhlus
- b Ventrals  $\geq 187$  ..... guentheri
- 42.a Dorsum with black and/or yellow longitudinal stripes ..... 43
- b Dorsum variable but without stripes ..... 46

- 43.a Dorsum blotched, ground color olive or brown, with darker blotches above and below narrow yellow paravertebral lines ..... 45
- b Dorsum striped, ground color tan or light brown, with three black stripes, or three black and two yellow stripes, the median stripe beginning on the snout ..... 44
- 44.a Dorsum with three dark and two narrow yellow stripes, yellow stripes and median dark stripe begin on snout ..... meridionalis
- b Dorsum with three wide black stripes, median black stripe begins on snout ..... dilepis
- 45.a Ventrals  $\leq 166$ ; maxillary teeth 13-17 ..... anomalus
- b Ventrals  $\geq 176$ ; maxillary teeth 18-20 ..... vanzolinii
- 46.a Dorsum uniform or variable, but never as below ..... 47
- b Dorsum tan or yellowish, with four rows of rounded black, reddish, or reddish brown spots, lateral row of spots smaller than paravertebral row ..... sagittifer (see 35)
- 47.a Dorsum blotched or with squarish spots ..... 48
- b Dorsum uniform brown to chocolate brown ..... perfuscus
- 48.a Head gray to black, occasionally with light or dark markings, but never red, no middorsal reddish stripe, and generally without rounded black, intercalary spots ..... 49
- b Head black, spotted with red; dorsum with large paravertebral black blotches on olive ground color, rounded lateral black intercalary spots, broad diffuse median ruddy stripe from head to tail, frequently invaded by dorsolateral black spots ..... elegantissimus

- 49.a Ground color usually gray or tan with a broad whitish dorsolateral line from midbody to tail on scale rows 4, 5 and 6, or 5 and 6; body with small squarish blotches slightly to moderately darker than ground color; venter checkered with black and some shade of pink or red maxillary teeth 19-25 (mode 21-22); diameter of eye 56-78% ( $\bar{x}$  = 66%) of snout length ..... Liophis sp.
- b Ground color usually brown to almost black, one subspecies with an ill-defined, posterior dorsolateral whitish stripe on scale rows 5 and 6. Head usually uniform brown or black; if brown, cephalic scales edged with black; dorsal color pattern highly variable, from salt and pepper pattern to blotches, bands, reticulations or combinations of the above and in many colors; venter from immaculate white to almost black. Maxillary teeth 13-21 (mode 17-18); diameter of eye 41-66% ( $\bar{x}$  = 54%) per cent of snout length ..... poecilogyrus

GENUS LIOPHIS Wagler, 1830:187Liophis almadensis (Wagler)

Natrix almadensis Wagler 1824:30. Type locality: Almada, Bahia, Brasil.

Natrix almada Wagler 1824:30. (as above)

Liophis conirostris Günther 1854:46. Type locality: Brasil and Bahia.

L. (iophis) verecundus Jan 863:300. Type locality: unknown.

Liophis wagleri Jan 1863:297. Type locality: Brasil and Bahia.

Liophis (Lygophis) y-graecum Peters 1882:129. Type locality: Villa de Guaratingueta, São Paulo (Brasil).

Trigonocephalus scolecomorphus Bacque 1906:116. Type locality: Asuncion, Paraguay.

Liophis almadensis Wagler 1830:188.

DISTRIBUTION: Chaco Boreal plant formation of Bolivia, all plant formations of Brazil except for Caatinga and Amazon rainforest; Paraguay except for cropland situations in central Paraguay. (published reports of this taxon from the state of Misiones, Argentina, have not been verified).

SUBSPECIES: none described; possibly one undescribed cryptic subspecies in Argentina and western Bolivia.

Liophis andinus Dixon

Liophis andinus Dixon 1983:129. Type locality: Incachaca, 2500 m, Cochabamba, Bolivia.

DISTRIBUTION: Known only from the type locality (Figure 2, p. 118, Dixon, 1983).

Liophis anomalus (Günther)

Coronella anomala Günther 1858:37. Type locality: banks of the Río Paraná.

Lygophis rutilus Cope 1862:80. Type locality: Río Tígre, a tributary of the Río Paraná, Paraguay.

Coronella pulchella Jan 1863:251. Type locality: Buenos Aires, Argentina.

Liophis anomala Amaral 1925:7.

DISTRIBUTION: Southern South America, from northwestern Argentina (Salta) east-southeast through Uruguay to Pôrto Alegre, Brasil, south to the province of Buenos Aires, Argentina.

Liophis atraventer Dixon and Thomas

Liophis atraventer Dixon and Thomas 1985:260. Type locality: Boracéia, São Paulo, Brasil.

DISTRIBUTION: Known only from the type locality (= Estação Biologica da Boracéia) (figure 4, p. 188, Dixon, 1987).

Liophis breviceps Cope

Liophis breviceps Cope 1860:252. Type locality: Suriname.

Liophis canaima Roze 1957:188. Type locality: Río Ugueto, Amazonas, Venezuela.

DISTRIBUTION: Eastern flanks of the Andes of Ecuador and Perú, eastward to Obidos, Brasil; on the north from central Colombia and northern Guyana south to Río Mamoré (Trinidad) Bolivia, and Posto Diuarum, Brasil.

SUBSPECIES: Two.

- 1.a Ventrals  $\leq 169$  ..... breviceps breviceps  
 b Ventrals  $\geq 176$  ..... breviceps canaimus

Liophis breviceps breviceps Cope 1862:252. Type locality: Suriname.

DISTRIBUTION: Forested parts of the Amazon Basin in Bolivia, Brasil, Colombia, Perú, and similar areas in French Guiana, Guyana, Suriname, and Venezuela (see figure 3, p. 154, Dixon 1983).

Liophis breviceps canaima Roze 1957:188. (see above)

DISTRIBUTION: Known only from the region of the Río Ugueto, Amazonas, Venezuela (see figure 3, p. 154, Dixon 1983).

Liophis carajasensis Cunha, Nascimento and Avila-Pires

Liophis carajasensis Cunha, Nascimento and Avila-Pires 1985:53. Type locality: Campo Rupestre d. N 1, Serra Norte, Pará, Brasil.

DISTRIBUTION: Known only from the type locality (see figure between pp. 18 & 19, Cunha, et al 1985).

Liophis cobellus (Linnaeus)

Coluber cobella Linnaeus 1758:218. Type locality: America.

Coluber serpens Daudin 1803:87. Type locality: none given.

Coluber cenchrus Daudin 1803:292. Type locality: Asia.

Liophis taeniogaster Jan 1863:292. Type locality: Brasil and South America.

Liophis trebbau Roze 1958:262. Type locality: Auyán Tepuí, Bolívar, Venezuela.

Liophis ingeri Roze 1958:303. Type locality: Chimantá Tepuí, Bolívar, Venezuela.

DISTRIBUTION: From Villavicencio, Colombia, south to Buenavista, Bolivia, and from Trinidad and Carapito, Venezuela, southeast to Bahia, Brasil.

SUBSPECIES: Four.

- 1.a Ventrals  $\geq 171$  ..... cobellus ingeri  
 b Ventrals  $\leq 164$  .....

- 2.a Ventral black bands  $\geq 45$  ..... cobellus cobellus  
 b Ventral black bands  $\leq 38$  ..... 3  
 3.a Ventrals vary from 130-147 ( $\bar{x}$  = 138.4) ..... cobellus dyticus  
 b Ventrals vary from 143-164 ( $\bar{x}$  = 151.6) ..... cobellus taeniogaster

Liophis cobellus cobellus (Linnaeus) 1758:292.

SYNONYMS: serpentinus Daudin, cenchrus Daudin.

DISTRIBUTION: Eastern Guyana Shield, exclusive of Venezuela Tepuí system; also Trinidad (see figure 4, p. 158, Dixon 1983).

Liophis cobellus dyticus Dixon 1983:159. Type locality: Monte Carmelo (= Requena), Loreto, Perú.

DISTRIBUTION: Western Amazon Basin, from Loma Linda, Colombia, south to Buenavista, Bolivia, east to Pôrto Velho, Brasil (see figure 4, p. 158, Dixon 1983).

Liophis cobellus taeniogaster Jan 1863:292. (see above)

DISTRIBUTION: Northeast Brasil, from Isla Bananal, east to central Bahia, north to Rio Amazonas (south bank) (see figure 4, p. 158, Dixon 1983).

Liophis cobellus trebbau Roze 1958:262.

SYNONYM: ingeri Roze.

DISTRIBUTION: Known only from the Chimantá and Auyán Tepuís, and km marker 144 of the El Dorado-Santa Elena highway, Bolívar, Venezuela (see figure 4, p. 158, Dixon 1983).

#### Liophis cursor (Lacépède)

Coluber cursor Lacépède 1789:96. Type locality: Martinique.

Coluber fugitivus Donndorf 1798:206. Type locality: Martinique.

Liophis putnami Cope 1862:78. Type locality: Martinique.

Liophis cursor andreoides Werner 1924:36. Type locality: Grenada.

DISTRIBUTION: Known only from the island of Martinique, West Indies (see figure 1, p. 297, Dixon 1981).

#### Liophis dilepis (Cope)

Lygophis dilepis Cope 1862:348. Type locality: Paraguay.

Liophis dilepis, Dixon 1980:7.

DISTRIBUTION: From northeast Brasil, south to southern Brasil, and Paraguay, extending northwest to southern Bolivia (see figure 1, p. 4, Michaud and Dixon 1987).



Liophis elegantissimus (Koslowsky)

Rhadinaea elegantissima Koslowsky 1895:115. Type locality: Sierra de la Ventana, Province of Buenos Aires, Argentina.

Liophis elegantissimus, Amaral 1925.

DISTRIBUTION: Known from five localities within the Sierra de la Ventana, Buenos Aires, Argentina (see figure 1, p. 566, Dixon 1985).

Liophis epinephelus Cope

Liophis epinephelus Cope 1862:78. Type locality: Truando, Colombia.

Liophis reginae albiventris Jan 1863:294. Type locality: Western Andes of Ecuador (fra Lacutunga e Guayaquil) (= Latacunga?).

Liophis reginae quadrilineata Jan 1863:295. Type locality: Ecuador.

Ophimorphus alticolus Cope 1868:102. Type locality: Quito Valley, Ecuador.

Zamensis ater Günther 1872:22. Type locality: Biscra, Algeria (in error).

Liophis fraseri Boulenger 1894:131. Type locality: Western Ecuador.

Liophis bimaculatus Cope 1899:71. Type locality: Colombia.

Liophis bipraeocularis Boulenger 1903:351. Type locality: Facatativa, Colombia.

Liophis opisthotaenia Boulenger 1908:114. Type locality: Mérida, Venezuela.

Liophis pseudocobella Peracca 1914:99. Type locality: Angelópolis, Colombia.

Liophis cobella alticolus Amaral 1931:87. Type locality: Jericó, Colombia.

Liophis taeniurus juvenalis Dunn 1937:213. Type locality: San Jose, Costa Rica.

Liophis bimaculatus lamonae Dunn 1944:486. Type locality: Sonsón, Antioquia, Colombia.

Liophis epinephelus ecuadorensis Laurent 1949:8. Type locality: Ecuador.

DISTRIBUTION: Trans-Andean South America from Venezuela to Perú; Cis-Andean South America at and above 2,200 m in Ecuador and Perú; also Central America from central Costa Rica through Panamá (see figures 1, 2, and 3, pp. 133-135, Dixon 1983).

SUBSPECIES: Eight.

- |     |  |                                   |
|-----|--|-----------------------------------|
| 1.a | Lateral black tail stripe absent, dorsal and ventral surfaces of body and tail alternately marked with red and black .....             | 2                                 |
| b   | Lateral black tail stripe present, dorsum may be banded, spotted, flecked, or almost unicolor, belly checkered with black or not ..... | 3                                 |
| 2.a | Subcaudals $\leq 54$ .....   | <u>epinephelus pseudocobellus</u> |
| b   | Subcaudals $\geq 54$ .....   | <u>epinephelus juvenalis</u>      |
| 3.a | Ventrals $\leq 165$ .....  | 4                                 |
| b   | Ventrals $\geq 165$ .....  | <u>epinephelus bimaculatus</u>    |

- 4.a Anterior half of dorsum with or without black flecks, streaks, or dark spots, never banded in adults; ventrals  $\geq 143$  ..... 5  
 b Anterior fourth of dorsum banded with black or brownish black bands, venter immaculate white, yellow or reddish; ventrals  $\leq 143$  ..... epinephelus epinephelus
- 5.a Venter immaculate white, yellow or pinkish ..... 6  
 b Venter checkered with marks of yellow and black ..... 7
- 6.a Dorsum leaf green with flecks of black, posterior dorsolateral black stripe usually absent ..... epinephelus albiventris  
 b Dorsum olive, olive brown, or grayish brown, posterior dorsolateral black stripe usually present ..... epinephelus opisthotaenius
- 7.a Ventrals from 141-156 ( $\bar{x}$  = 150), subcaudals from 51-67 ( $\bar{x}$  = 59) ..... epinephelus lamonae  
 b Ventrals from 151-164 ( $\bar{x}$  = 157), subcaudals from 61-76 ( $\bar{x}$  = 67) ..... epinephelus fraseri

Liophis epinephelus epinephelus Cope 1862.

DISTRIBUTION: Lower elevations of the mountains of western Panamá, east to Colombian lowlands, most interAndean valleys below 1,500 m, extending southward along the Colombian coast to northern Ecuador (see figures 1-3, pp. 133-135, Dixon 1983).

Liophis epinephelus albiventris Jan 1863.

SYNONYMS: alticolus Cope, ater Günther, quadrilineatus Jan

DISTRIBUTION: Western Ecuador from sealevel to 2,600 m elevation (see figure 3, p. 153, Dixon 1983).

Liophis epinephelus bimaculatus Cope 1899.

SYNONYMS: bipraeocularis Boulenger, ecuadorensis Laurent

DISTRIBUTION: High Andean slopes (2,600 - 3,300 m) of western Venezuela, central Colombia, south to northern Perú (see figures 2, 3, p. 134, 135, Dixon 1983).

Liophis epinephelus fraseri Boulenger 1894.

DISTRIBUTION: Middle elevations of the eastern and western slopes of southern Ecuador, south to central Perú (see figure 4, p. 136, Dixon 1983).

Liophis epinephelus juvenalis Dunn 1937.

DISTRIBUTION: Middle slopes of mountains from central Costa Rica to western Panamá (see figure 1, p. 133, Dixon 1983).

Liophis epinephelus lamonae Dunn 1944.

DISTRIBUTION: Andean slopes (1,500 - 2,600 m) of Colombia, southward to east-central Ecuador (see figures 2, 3, pp. 134, 135, Dixon 1983).

Liophis epinephelus opisthotaenius Boulenger 1908.

DISTRIBUTION: Mérida region of Venezuela and the Páramo de Tama region of Venezuela and Colombia (see figure 2, p. 134, Dixon 1983).

Liophis epinephelus pseudocobellus Peracca 1914.

SYNONYM: alticolus Amaral.

DISTRIBUTION: Middle elevations of central and western Andes of Colombia, south to Ecuador border (see figure 2, p. 134, Dixon 1983).

Liophis festae (Perraca)

Rhadinaea festae Peracca 1897:16. Type locality: Valley of Río Santiago, Ecuador.

Liophis festae, Amaral 1929:171.

DISTRIBUTION: From middle elevations of southern Colombia, south through Ecuador, to the middle elevations of Cis-Andean central Perú.

Liophis flavifrenatus (Cope)

Lygophis flavifrenatus Cope 1862:80. Type locality: Río Vermejo, (Bermejo region) Paraguay.

Dromicus amabilis Jan 1867: livr. 24. Type locality: Brasil.

Liophis flavifrenatus, Dixon 1980:8.

DISTRIBUTION: Southern Brasil, southward through central and southern Paraguay, northeastern Argentina, and extreme southeastern Brasil (see figure 3, p. 9, Michaud and Dixon 1987).

Liophis frenatus (Werner)

Rhadinaea frenata Werner 1909:224. Type locality: Paraguay.

Rhadinaea brazili Amaral 1923:87. Type locality: Julio Pontes, Brasil.

Liophis frenatus, Amaral 1929:45.

DISTRIBUTION: From Primavera, Paraguay, east-southeast to Guayuvira, Brasil (see figure 2, p. 154, Dixon 1983).

Liophis guentheri Peracca

Liophis guentheri Peracca 1897:11. Type locality: Caiza, Bolivia.

DISTRIBUTION: Apparently restricted to the central Chaco region of Argentina, Bolivia and Paraguay (see figure 1, p. 175, Dixon 1987).

Liophis jaegeri (Günther)

Coronella jaegeri Günther 1858:37. Type locality: Brasil.

Liophis dorsalis Peters 1863:283. Type locality: Brasil.

Aporophis coralliventris Boulenger 1894:346. Type locality: an island north of Concepcion, near San Salvador, north Paraguay.

Rhadinaea lineata Jensen 1900:105. Type locality: Taboleiro Grande, Minas Gerais, Brasil.

Liophis jaegeri, Amaral 1926:78.

DISTRIBUTION: Southeast Brasil to coastal Uruguay, and the Río Paraná Basin of Argentina, Brasil, and Paraguay (see figure 3, p. 184, Dixon 1987).

SUBSPECIES: Two.

- 1.a Subcaudals from 63-75 ( $\bar{x}$  = 68.5); tail/total length ratios from 0.214-0.268 ( $\bar{x}$  = 0.231) .... jaegeri coralliventris  
 b Subcaudals from 52-71 ( $\bar{x}$  = 60.4); tail/total length ratios from 0.193-0.248 ( $\bar{x}$  = 0.221) ..... jaegeri jaegeri

Liophis jaegeri jaegeri (Günther) 1858.

SYNONYMS: dorsalis Peters, lineata Jensen.

DISTRIBUTION: Southeastern Brasil and coastal Uruguay, east of the Río Paraná Basin (see figure 3, p. 184, Dixon 1987).

Liophis jaegeri coralliventris (Boulenger) 1894 (see above)

DISTRIBUTION: Known only from the Río Paraná Basin of Argentina, Brasil, and Paraguay (see figure 3, p. 184, Dixon 1987).

Liophis juliae (Cope)

Aporophis juliae Cope 1879:373. Type locality: Dominica.

Leimadophis mariae Barbour 1914:340. Type locality: Marie-Galante

Dromicus juliae copeae Parker 1936:232. Type locality: Guadeloupe

Liophis juliae, Dixon 1980:10.

DISTRIBUTION: West Indian islands of Dominica, Guadeloupe, and Marie-Galante.

SUBSPECIES: None recognized.

Liophis lineatus (Linnaeus)

Coluber lineatus Linnaeus 1758:221. Type locality: Asia (in error)

Coluber jaculatrix Linnaeus 1766:381. Type locality: Suriname

Coluber terlineatus Lacépède 1826:106. Type locality: none given.

Liophis lineatus, Dixon 1980:10.

SUBSPECIES: None recognized.

DISTRIBUTION: Central Panamá east through Colombia, Venezuela, Guyana, Suriname, French Guiana, to the mouth of the Rio Amazonas, Brasil (Figure 1, p. 4, Michaud and Dixon 1987). A few specimens are known from near the port cities of Guayaquil and Esmeraldas, Ecuador, and are probably accidental introductions via the shipping trade.

Liophis longiventris Amaral

Liophis longiventris Amaral 1925:16. Type locality: none given, but probably the state of Mato Grosso, Brasil.

DISTRIBUTION: Known only from Rio Manjuro, Amazonas, Brasil, and from 12°51'S - 51°46'W, Mato Grosso, Brasil (see figure 2, p. 154, Dixon 1983).

Liophis maryellenae Dixon

Liophis maryellenae Dixon 1985:295. Type locality: Annapolis, Goiás, Brasil.

DISTRIBUTION: Central, southeast Brasil, from Annapolis on the west, to Grão Mogol, on the east; from near Barreiras on the north, to Itambe do Dentro on the south (see figure 4, p. 188, Dixon 1987).

Liophis melanotus (Shaw)

Coluber melanotus Shaw 1802:534. Type locality: Cape of Good Hope, Africa (in error).

Coluber raninus Merrem 1820:106. Type locality: none given.

Coluber vittatus Hallowell 1845:242. Type locality: within 200 miles of Caracas, Colombia (actually in Venezuela).

Liophis melanonotus Cope 1860:253. (replacement name for melanotus Shaw).

Liophis melanotus, Dixon 1980:11.

DISTRIBUTION: From both sides of the Andes in Central Colombia, northeast to northeastern Venezuela, Trinidad and Tobago.

SUBSPECIES: Possibly one, undescribed.

- 1.a Dorsolateral black stripe begins on the nape and is continuous to the tail, head more or less uniform in color, without distinct white parietal spots, maxillary teeth vary from 14-17 ( $\bar{x}$  = 15.3) ..... melanotus (subsp.)
- b Dorsolateral black stripe begins posterior to nape, interrupted anteriorly by nape bands or spots that are separated by light colored interspaces, parietals with white diagonal marks extending to the posterior corner of mouth, maxillary teeth vary from 14-21 ( $\bar{x}$  = 17.9) ..... melanotus melanotus

Liophis meridionalis (Schenkel)

Aporophis lineatus meridionalis Schenkel 1901:160. Type locality: Mte. Sociedad, Bemalcue, Paraguay.

Aporophis lineatus lativittatus Mueller 1908:74. Type locality: San Fermin (Chiquitos), Bolivia.

Liophis meridionalis, Dixon 1980:11.

DISTRIBUTION: Central Brasil and northern Bolivia south to southern Paraguay, northeastern tip of Argentina, and extreme southeastern Brasil. (figure e, p. 9, Michaud and Dixon, 1987).

Liophis miliaris (Linnaeus)

Coluber miliaris Linnaeus 1758:220. Type locality: India (in error).

Coluber merremii Wied 1821:121. Type locality: Pedro d'Alcantara, Bahia, Brasil.

C. (oluber) dictyodes Wied 1824:668. Type locality: Cabo Frio, Rio de Janeiro, Brasil.

Coluber bicolor Reuss 1834:145. Type locality: Ilheus, Bahia, Brasil.

Ablabes purpurans Dumeril, Bibron and Dumeril 1854:312. Type locality: Mana, Cayenne. (French Guiana)

Coronella australis Günther 1858:40. Type locality: Australia (in error).

Opheomorphus merremii semiaureus Cope 1862:348. Type locality: Paraguay.

Liophis cobella collaris Jan 1863:293. Type locality: South America.

Liophis reginae ornata Jan 1863:295. Type locality: Buenos Aires, Argentina.

Coronella orientalis Günther 1864:236. Type locality: Dekkan (in error).

Rhadinaea chrysostoma Cope 1868:104. Type locality: Napo or Marañon, Ecuador (= Perú).

Coronella poecilolaemus Günther 1872:19. Type locality: Upper Río Amazonas.

Opheomorphus fuscus Cope 1885:190. Type locality: São João da Monte Negro, Rio Grande do Sul, Brasil.

Rhadinaea orina Griffin 1916:195. Type locality: Sierras of Bolivia (in error).

Dromicus amazonicus Dunn 1922:219. Type locality: Santarém, Brasil.

Rhadinaea merremii natricodes Werner 1926:246. Type locality: unknown.

Liophis mossoroensis Hoge and Lima-Verde 1972:215. Type locality: Mossoro, Rio Grande do Norte, Brasil.

Liophis miliaris, Amaral 1926:78.

DISTRIBUTION: Eastern South America, from Guyana south to Buenos Aires, Argentina, with scattered records in the Amazon Basin and Cerrado of Brasil (see figure 1, p. 792, Dixon 1983; figures 5, 6, and 7, pp. 12-14, Gans 1964).

SUBSPECIES: Seven.

- 1.a Venter with large medial, contrasting yellow and black marks from near throat to the anal plate ..... miliaris chrysostomus
- b Venter without checkerboard pattern of yellow and black marks, but ventrals may be edged or have a suffusion of dark color ..... 2
- 2.a Ventrals from 163-190 ( $\bar{x}$ = 176.2); dorsum typically with dark blotches with light interspaces ..... miliaris semiaureus
- b Ventrals  $\leq 173$ ; dorsum typically unicolored or with light centers to each scale, or with dorsal and lateral dark blotches separated by dorsolateral light lines posteriorly, or with an almost black dorsum with light flecks scattered throughout ..... 3
- 3.a Subcaudals  $\geq 76$  or more, juveniles with a pair of light nuchal spots, adults uniform brown, each dorsal scale with a pale light center ..... miliaris amazonicus
- b Subcaudals  $\leq 68$ ; juvenile pattern variable, but never with a pair of light nuchal spots, adults light tan to black, with light centered scales, or with light flecks scattered throughout ..... 4
- 4.a Dorsum light to dark brown, each scale with a light center, venter obscurely marked with dark or each ventral lightly edged with black ..... 5
- b Dorsum dark brown to black with scattered flecks of white, or with both dorsal and lateral dark blotches separated posteriorly by a dorsolateral light line; venter uniform white, or marked with black ..... 6
- 5.a Juveniles with dark-edged gulars; venter marked with yellow and black. Adults with or without obscure marks on the gulars; dorsum dark brown with obscure light centers to each scale; venter with less contrasting dark and light marks ..... miliaris miliaris
- 5.b Juveniles with white gulars; venter white or ventrals edged with black laterally. Adults with white venters, but each ventral lightly edged with black laterally and black color eventually meeting at the midline posteriorly; dorsum of adults brown to dark brown with highly contrasting light centers to each scale ..... miliaris orinus

- 6.a Ventrals from 148-166 ( $\bar{x}$  = 158.7); dorsum dark brown to black with many scattered white flecks; edges of ventrals and subcaudals heavily marked with black from midbody to the tip of the tail  
 ..... miliaris mossoroensis
- b Ventrals from 135-156 ( $\bar{x}$  = 146.5); dorsum with a series of dark brown to black dorsal and lateral blotches on a light tan, greenish or brown ground color, with a dorsolateral light line separating the dorsal and lateral blotches posteriorly. Occasionally, the dorsum is uniform in color with a light center to each scale; ventrals and subcaudals immaculate white or cream ..... miliaris merremii

Liophis miliaris miliaris (Linnaeus) 1758.

SYNONYMS: purpurans Dumeril, Bibron and Dumeril; orientalis Günther; collaris Jan.

DISTRIBUTION: Guyana, Suriname, and French Guiana (see figure 1, p. 792, Dixon 1983).

Liophis miliaris amazonicus (Dunn) 1922.

DISTRIBUTION: Santarém, Brasil, south to Rio Iténez, Beni, Bolivia, east and south to Mato Grosso, Brasil (see figure 1, p. 792, Dixon 1983).

Liophis miliaris chrysostomus (Cope) 1868.

SYNONYM: poecilolaemus Günther.

DISTRIBUTION: Rainforests of Brasil, Colombia, Ecuador and Perú (see figure 1, p. 792, Dixon 1983).

Liophis miliaris merremii (Wied) 1821.

SYNONYMS: australis Günther; bicolor Reuss; dictyodes Wied.

DISTRIBUTION: Recife, Pernambuco, Brasil, south-southwest to Rio de Janeiro (principally the Brazilian Atlantic rainforest).

Liophis miliaris mossoroensis Hoge and Lima-Verge 1972.

DISTRIBUTION: Northeastern Brasil, primarily Caatinga and dry Cerrado (see figure 1, p. 792, Dixon 1983).

Liophis miliaris orinus (Griffin) 1914.

SYNONYM: natricoides Werner.

DISTRIBUTION: Southeastern Brasil, from southern Minas Gerais, south through the states of São Paulo, Paraná, Santa Catarina, to the northern one-third of Rio Grande do Sul (see figure 1, p. 792, Dixon 1983).



Liophis miliaris semiaureus (Cope) 1862.

SYNONYMS: fuscus Cope; ornata Jan.

DISTRIBUTION: Paraguay, west and south of Iguazú Falls; northeastern Argentina; southern and eastern Uruguay; southern one-half of the Brazilian state of Rio Grande do Sul (see figure 1, p. 792, Dixon 1983).

Liophis ornatus (Garman)

Dromicus ornatus Garman 1887:281. Type locality: Saint Lucia, West Indies.

Dromicus giganteus Jan 1863:67. Type locality: unknown (see Dixon, 1981, concerning the nature of this senior synonym).

Leimadophis boulengeri Barbour 1914:339 (replacement name for ornatus Garman)

Liophis ornatus, Dixon 1981:13.

DISTRIBUTION: Saint Lucia and the satellite island of Maria (see figure 1, p. 792, Dixon 1983).

Liophis paucidens (Hoge)

Lygophis paucidens Hoge 1953:253. Type locality: Mato Verde, Mato Grosso, Brasil.

Liophis paucidens, Dixon 1980:13.

DISTRIBUTION: Known only from east-central Brasil (see figure 3, p. 9, Michaud and Dixon 1987).

Liophis perfuscus Cope

Liophis perfuscus Cope 1862:77. Type locality: Barbados.

Liophis rufus Jan 1863:91. Type locality: unknown.

DISTRIBUTION: Known only from the West Indian island of Barbados (see figure 1, p. 297, Dixon 1981).

Liophis poecilogyrus (Wied)

Coluber poecilogyrus Wied 1825:371. Type locality: Barra de Jucú, Rio Espirito Santo, Brasil.

Coluber m-nigrum Raddi 1820:338. Type locality: Rio de Janeiro, Brasil.

Natrix G. forsteri Wagler 1824:16. Type locality: Bahia, Brasil.

Coluber doliatus Wied 1825:368. Type locality: Barra de Jucú, Rio Espirito Santo, Brasil.

X. (enodon) schotti Schlegel 1837:91. Type locality: South America.

Liophis merremii sublineatus Cope 1860:252. Type locality: Buenos Aires, Argentina.

Opheomorphus doliatus caesius Cope 1862:348. Type locality: Santa Fé, Paraguay.

Liophis subfasciatus Cope 1862:77. Type locality: Paraguay.

Liophis ornatissima Jan 1863:53. Type locality: Paraná (Brasil?).

Liophis typhlus gastrostictus Jan 1863:53. Type locality: Fernambuco (= Pernambuco, Brasil).

- Liophis reginae viridicyanea Jan and Sordelli 1866:18(2)91). Type locality: Paraná, Brasil.
- Liophis cobella flaviventris Jan and Sordelli 1866:16(5)92). Type locality: South America.
- Rhadinaea dichroa Werner 1899:115. Type locality: Argentina.
- Rhadinaea praeornata Werner 1909:58. Type locality: central Brasil.
- Leimadophis poecilogyrus reticulatus Parker 1931:285. Type locality: Makthlawaiya, Paraguay.
- Leimadophis poecilogyrus albadspersus Amaral 1944:78. Type locality: Piracicaba, São Paulo, Brasil.
- Leimadophis poecilogyrus amazonicus Amaral 1944:81. Type locality: probably Pará, Brasil (but not stated as such).
- Leimadophis poecilogyrus franciscanus Amaral 1944:80. Type locality: Pirapora, Minas Gerais, Brasil.
- Leimadophis poecilogyrus intermedius Amaral 1944:81. Type locality: Goiás, Brasil.
- Leimadophis poecilogyrus montanus Amaral 1944:79. Type locality: Piquete, São Paulo, Brasil.
- Leimadophis poecilogyrus pictostriatus Amaral 1944:77. Type locality: Sao Lourenço, Brasil.
- Leimadophis poecilogyrus pineticola Amaral 1944:78. Type locality: central Paraná, Brasil.
- Leimadophis poecilogyrus platensis Amaral 1944:77. Type locality: La Plata, Argentina.
- Leimadophis poecilogyrus xerophilus Amaral 1944:81. Type locality: probably Ceara, Brasil (but not stated as such).
- Leimadophis poecilogyrus lancinii Hoge, Romano and Cordeiro 1978:77. (replacement name for L. p. amazonicus Amaral).
- Liophis poecilogyrus, Dixon 1980:13.

DISTRIBUTION: Much of eastern South America, from Venezuela (?) east and south through Brasil to central Bolivia, southeast into northeastern Argentina.

SUBSPECIES: None to possibly nine. I do not recognize subspecies of this taxon because of the mosaic nature of the variation examined thus far. Note that there are three senior synonyms for the name poecilogyrus. Any use of a senior synonym would upset the stability of the long, continued use of poecilogyrus in the literature, and I recommend none be used.

#### Liophis problematicus Myers

Liophis problematicus Myers 1986:2. Type locality: San Juan, Río Tambopata; Sandia Province, 14°13'S - 69°10'W, 1,520 m, Puno, Perú.

DISTRIBUTION: Known only from type locality.

#### Liophis reginae (Linnaeus)

- Coluber reginae Linnaeus 1758:219. Type locality: India (in error).
- Coluber violaceus Lacépède 1789:116. Type locality: none given.
- Coluber graphicus Shaw 1802:474. Type locality: America.
- Natrix semilineata Wagler 1824:33. Type locality: Rio Solimões, Brasil.

Liophis oligolepis Boulenger 1905:455. Type locality: Igapé-Assu, Pará, Brasil.

Leimadophis reginae macrosoma Amaral 1935:238. Type locality: Canna Brava, Goiás, Brasil.

Leimadophis reginae maculicauda Hoge 1954:241. Type locality: none given.

Leimadophis zweifeli Roze 1959. Type locality: Rancho Grande, Aragua, Venezuela.

Liophis reginae, Dixon 1980:24.

DISTRIBUTION: Cis-Andean South America, from Colombia to northern Argentina; also Trinidad and Tobago (see figure 2, p. 118, Dixon 1983).

SUBSPECIES: Four.

- |     |  |                             |
|-----|--|-----------------------------|
| 1.a | Dorsum with small black and yellow spots; black lateral caudal stripe faint or absent .....  | 2                           |
| b   | Dorsum greenish, olive, or grayish, never with small yellow and black spots; black lateral caudal stripe always present and distinct ..... | 3                           |
| 2.a | Subcaudals average 80 (69-88) .....  | <u>reginae zweifeli</u>     |
| b   | Subcaudals average 65 (55-78) .....  | <u>reginae semilineatus</u> |
| 3.a | Dorsal scale rows one and two pale colored, in contrast to dorsal coloration .....   | 4                           |
| b   | Dorsal scale rows one and two similar in color to rest of body .....   | 5                           |
| 4.a | Dorsum with dense pale and dark paravertebral flecking; subcaudals average 74 (63-80) .....  | <u>reginae reginae</u>      |
| b   | Dorsum without pale and dark paravertebral flecking; subcaudals average 67 (63-80) .....   | <u>reginae semilineatus</u> |
| 5.a | Subcaudals with ventrolateral black spots, flecks, or smudges; subcaudals average 81 (75-91) ....  | <u>reginae macrosomus</u>   |
| b   | Subcaudals immaculate; subcaudals average 70 (55-81) .....   | <u>reginae semilineatus</u> |

Liophis reginae reginae (Linnaeus) 1758.

SYNONYMS: violaceus Lacépède; graphicus Shaw.

DISTRIBUTION: Guyana, Suriname, and French Guiana (see figure 2, p. 118, Dixon 1983).

Liophis reginae macrosomus (Amaral) 1935.

SYNONYM: maculicaudus Hoge.

DISTRIBUTION: Chaco and Cerrado of Argentina, Bolivia, Brasil, and Paraguay (see figure 2, p. 118, Dixon 1983).

Liophis reginae semilineatus (Wagler) 1824.

SYNONYM: oligolepis Boulenger.

DISTRIBUTION: Forested Amazon Basin of Venezuela, Colombia, Ecuador, Perú, Bolivia, and Brasil; also Atlantic rainforest of Brasil (see figure 2, p. 118, Dixon 1983).

Liophis reginae zweifeli (Roze) 1959.

DISTRIBUTION: Montane rainforests of Venezuela and Trinidad (see figure 2, p. 118, Dixon 1983).

Liophis sagittifer (Jan)

L. (iopeltis) sagittifer Jan 1863:82. Type locality: Mendoza, Argentina.

Liophis pulcher Steindachner 1867:267. Type locality: Chile (in error).

Rhadinaea modesta Koslowsky 1896:453. Type locality: Salta, Argentina.

Liophis trifasciatus Werner 1899:114. Type locality: Paraguay.

Zamensis argentinus Bréthès 1917:93. Type locality: La Banda, Santiago del Estero, Argentina.

Liophis sagittifer, Dixon 1980:15.

DISTRIBUTION: Foothills of the Andes of Bolivia and Argentina, south to Chubut, Argentina, north and east into the Monte and Chaco of Argentina and Paraguay. (see figure 2, p. 391, Dixon and Thomas 1982).

SUBSPECIES: Two.

- 1.a Dorsum uniform brownish gray, olive gray, or with obscure undulating middorsal dark line and some indication of a lateral dark line bordering the upper edge of scale row three; 81% of population with 19-19-17 scale rows ..... sagittifer modestus
- b Dorsum with large to median sized paravertebral reddish black to black blotches from nape to tail; often with secondary row of intercalary black blotches laterally, and occasionally a median series of dorsal blotches anteriorly; 100% of population with 19-19-15 scale rows ..... sagittifer sagittifer

Liophis sagittifer sagittifer (Jan) 1863.

SYNONYMS: pulcher Steindachner; argentinus Bréthès.

DISTRIBUTION: Monte of Patagonia, from Tucuman, south to the state of Chubut, Argentina. (see figure 2, p. 391, Dixon and Thomas 1982).

Liophis sagittifer modestus (Koslowsky) 1896.

SYNONYM: trifasciatus Werner.

DISTRIBUTION: Chaco-Bonariesian Plain of Argentina, Bolivia, and Paraguay. (see figure 2, p. 391, Dixon and Thomas 1982).

Liophis taeniurus Tschudi

Liophis taeniurus Tschudi 1845:164. Type locality: Perú, in der heissen waldregion.

DISTRIBUTION: Middle and upper elevations of the Andes in southern Ecuador, south through Perú to the Cochabamba region of Bolivia.

SUBSPECIES: None described; possibly two or more suggested from recent analysis of data from additional material.

Liophis triscalis (Linnaeus)

Coluber triscalis Linnaeus 1758:224. Type locality: India (in error).

Coluber corallinus Linnaeus 1758:223. Type locality: Asia (in error).

Liophis triscalis, Boulenger 1894:129.

DISTRIBUTION: Known only from the Leeward Island of Curaçao. (see figure 1, p. 297, Dixon 1981).

Liophis typhlus (Linnaeus)

Coluber typhlus Linnaeus 1758:218. Type locality: India (in error).

Xenodon isolepis Cope 1870:155. Type locality: Pebas, Ecuador (= Perú)

Opheomorphus brachyurus Cope 1887:57. Type locality: Chupada, Mato Grosso, Brasil.

Liophis elaeoides Griffin 1916:187. Type locality: Prov. del Sara, Bolivia.

Liophis macrops Werner 1925:57. Type locality: Paramaribo, Suriname.

Liophis typhlus, Dixon 1980:16.

DISTRIBUTION: Rainforests of the Guiana Shield and Amazon Basin, also the Chaco and Cerrado of Bolivia, Brasil, and Paraguay. (see figure 1, p. 175, Dixon 1987).

SUBSPECIES: Three.

1.a Ventrals 133-163 ( $\bar{x}$  = 147.3); juveniles and adults  
with dark paravertebral chevron marks ..... typhlus typhlus

b Ventrals 158-172 ( $\bar{x}$  = 163.5); juveniles and adults  
without dark chevron marks ..... 2

2.a Subcaudals 40-49 ( $\bar{x}$  = 44.4); tail/total length ratios  
0.140-0.160 ( $\bar{x}$  = 0.149) ..... typhlus brachyurus

b Subcaudals 49-56 ( $\bar{x}$  = 52.0); tail/total length ratios  
0.160-0.200 ( $m$  = 0.171) ..... typhlus elaeoides

Liophis typhlus typhlus (Linnaeus) 1758.

SYNONYMS: isolepis Cope; macrops Werner.

DISTRIBUTION: Rainforests of Guiana Shield, and Amazon Basin (see figure 1, p. 175, Dixon 1987).

Liophis typhlus brachyurus (Cope) 1887.

DISTRIBUTION: Deciduous mesophytic forests of southeastern Brasil, and the Campo Cerrado forests of east-central Brasil (see figure 1, p. 175, Dixon 1987).

Liophis typhlus elaeoides Griffin 1916.

DISTRIBUTION: Mesic Chaco forests of southeastern Bolivia; northern Paraguay, and western Mato Grosso, Brasil (see figure 1, p. 175, Dixon 1987).

Liophis vanzolinii Dixon

Liophis vanzolinii Dixon 1985:567. Type locality: Achiras, Cordoba, Argentina.

DISTRIBUTION: Known only from three localities in the western part of the Argentine state of Cordoba (see figure 1, p. 566, Dixon 1985).

Liophis viridis Günther

Liophis viridis Günther 1862:58. Type locality: Pernambuco, Brasil  
Liophis typhlus prasina Jan and Sordelli 1866:18(4)(3). Type locality: Brasil.

DISTRIBUTION: The Caatinga, Agreste, and Atlantic rainforests of Brasil. (see figure 2, p. 181, Dixon 1987).

SUBSPECIES: Two.

- 1.a Ventrals 169-188 ( $\bar{x}$  = 177); reduction site ventrals  
98-116 ( $\bar{x}$  = 106.6) ..... viridis viridis  
b Ventrals 181-202 ( $\bar{x}$  = 189.8); reduction site ventrals  
102-123 ( $\bar{x}$  = 114.6) ..... viridis prasinus

Liophis viridis viridis Günther 1862.

DISTRIBUTION: The agreste and Atlantic rainforests of Brasil, from Recife to Salvador (see figure 2, p. 181, Dixon 1987).

Liophis viridis praesinus Jan and Sordelli 1866.

DISTRIBUTION: The Caatinga forest of Brasil (see figure 2, p. 181, Dixon 1987).

Liophis williamsi (Roze)

Urotheca williamsi Roze 1958:1. Type locality: El Junquito, D.F., Venezuela.  
Liophis williamsi, Dixon 1980:17.

DISTRIBUTION: Cloud Forests of the coastal Andes of Venezuela (see figure 2, p. 118, Dixon 1983).

## COUNTRY LISTS

(containing taxa of Liophis)

## CENTRAL AMERICA

## COSTA RICA

Liophis epinephelus juvenalis

## PANAMÁ

Liophis epinephelus juvenalis  
Liophis epinephelus epinephelus  
Liophis lineatus

## SOUTH AMERICA

## ARGENTINA

Liophis sp. nov.  
Liophis anomalus  
Liophis elegantissimus  
Liophis flavifrenatus  
Liophis guentheri  
Liophis jaegeri coralliventris  
Liophis miliaris semiaureus  
Liophis meridionalis  
Liophis poecilogyrus  
Liophis reginae macrosomus  
Liophis sagittifer modestus  
Liophis sagittifer sagittifer  
Liophis vanzolinii

## BOLIVIA

Liophis almadensis sub-sp.  
Liophis almadensis almadensis  
Liophis andinus  
Liophis breviceps breviceps  
Liophis cobella dyticus  
Liophis dilepis  
Liophis guentheri  
Liophis meridionalis  
Liophis miliaris amazonicus  
Liophis poecilogyrus  
Liophis reginae macrosomus  
Liophis reginae semilineatus  
Liophis sagittifer modestus  
Liophis taeniurus  
Liophis typhlus elaeoides  
Liophis typhlus typhlus

## BRASIL

Liophis atraventer  
Liophis almadensis  
Liophis anomalus  
Liophis breviceps breviceps  
Liophis carajasensis  
Liophis cobellus dyticus  
Liophis cobellus taeniogaster  
Liophis dilepis  
Liophis flavifrenatus  
Liophis frenatus  
Liophis jaegeri jaegeri  
Liophis jaegeri coralliventris  
Liophis lineatus  
Liophis longiventris  
Liophis maryellenae  
Liophis meridionalis  
Liophis miliaris amazonicus  
Liophis miliaris chrysostomus  
Liophis miliaris merremii  
Liophis miliaris mossoroensis  
Liophis miliaris orinus  
Liophis miliaris semiaureus  
Liophis paucidens  
Liophis poecilogyrus  
Liophis reginae macrosomus  
Liophis reginae semilineatus  
Liophis typhlus brachyurus  
Liophis typhlus elaeoides  
Liophis typhlus typhlus  
Liophis viridis prasinus  
Liophis viridis viridis

## COLOMBIA

Liophis breviceps breviceps  
Liophis cobellus dyticus  
Liophis epinephelus bimaculatus  
Liophis epinephelus epinephelus  
Liophis epinephelus lamonae  
Liophis epinephelus opisthotaenius  
Liophis epinephelus pseudocobellus  
Liophis festae  
Liophis lineatus  
Liophis melanotus melanotus  
Liophis miliaris chrysostomus  
Liophis reginae semilineatus  
Liophis typhlus typhlus



## ECUADOR

Liophis breviceps breviceps  
Liophis cobellus dyticus  
Liophis epinephelus albiventris  
Liophis epinephelus bimaculatus  
Liophis epinephelus epinephelus  
Liophis epinephelus fraseri  
Liophis epinephelus lamonae  
Liophis festae  
Liophis lineatus  
Liophis miliaris chrysostomus  
Liophis reginae semilineatus  
Liophis taeniurus  
Liophis typhlus typhlus

## FRENCH GUIANA

Liophis breviceps breviceps  
Liophis cobellus cobellus  
Liophis lineatus  
Liophis miliaris miliaris  
Liophis poecilogyrus  
Liophis reginae reginae  
Liophis typhlus typhlus

## GUYANA

Liophis breviceps breviceps  
Liophis cobellus cobellus  
Liophis lineatus  
Liophis miliaris miliaris  
Liophis poecilogyrus  
Liophis reginae reginae  
Liophis typhlus typhlus

## PARAGUAY

Liophis almadensis  
Liophis dilepis  
Liophis flavifrenatus  
Liophis frenatus  
Liophis quentheri  
Liophis jaegeri coralliventris  
Liophis longiventris  
Liophis meridionalis  
Liophis miliaris semiaureus  
Liophis poecilogyrus  
Liophis reginae macrosomus  
Liophis sagittifer modestus  
Liophis typhlus elaeoides

## PERÚ

Liophis breviceps breviceps  
Liophis cobellus dyticus  
Liophis epinephelus fraseri  
Liophis festae  
Liophis miliaris chrysostomus  
Liophis problematicus  
Liophis miliaris chrysostomus  
Liophis reginae semilineatus  
Liophis taeniurus  
Liophis typhlus typhlus

## SURINAME

Liophis breviceps breviceps  
Liophis cobellus cobellus  
Liophis lineatus  
Liophis miliaris miliaris  
Liophis poecilogyrus  
Liophis reginae reginae  
Liophis typhlus typhlus

## URUGUAY

Liophis anomalus  
Liophis jaegeri jaegeri  
Liophis miliaris semiaureus  
Liophis poecilogyrus

## VENEZUELA

Liophis breviceps breviceps  
Liophis breviceps canaimus  
Liophis cobellus cobellus  
Liophis cobellus trebbau  
Liophis epinephelus opisthotaenius  
Liophis lineatus  
Liophis melanotus melanotus  
Liophis melanotus subspecies  
Liophis poecilogyrus (?)  
Liophis reginae semilineatus  
Liophis reginae zweifeli  
Liophis typhlus typhlus  
Liophis williamsi

## CARIBBEAN ISLANDS

## BARBADOS

Liophis perfuscus

## CARIBBEAN ISLANDS (Continued)

## CURACAO

Liophis triscalis

## DOMINICA

Liophis juliae

## GUADELOUPE

Liophis juliae

## MARIE-GALANTE

Liophis juliae

## MARTINIQUE

Liophis cursor

## SAINT LUCIA

Liophis ornatus

## TOBAGO

Liophis melanotus


Liophis reginae


## TRINIDAD


Liophis cobellus cobellus

Liophis melanotus subsp.


Liophis reginae zweifeli































66

64

62

Montserrat

Guadeloupe

Desirade

*L. juliae*

Marie-Galante

Dominica

*L. cursor*

Martinique

*L. ornatus*

Santa Lucia

*L. perfuscus*

Barbados

*L. triscalis*

Curacao

San Vicente

The Grenadines

Grenada

*L. reginae*

Tobago

*L. cobellus*

*L. melanotus*

Trinidad

VENEZUELA

16

16

14

14

12

12


10

10

66

64

62











c  
T

A SURVEY OF OFFICIALLY REJECTED  
NOMINAL HERPETOLOGICAL TAXA  
AND  
THEIR ALLOCATIONS

HOBART M. SMITH & DAVID CHISZAR

Departments of Environmental, Population and Organismic  
Biology (HMS) & Psychology (DC)  
University of Colorado



SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
NO. 80

1989

SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with the Division of Amphibians and Reptiles.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions on preparation and submission. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington DC 20560 USA. Please include a self-addressed mailing label with requests.

## INTRODUCTION

All herpetologists undertaking taxonomic work should utilize the recent summary (Melville and Smith, 1987) of all names and works placed through 1985 on the Official Lists and Indices of the International Commission on Zoological Nomenclature (available, together with a mimeographed supplement covering 1986-1988, from the International Trust for Zoological Nomenclature, c/o British Museum (Natural History), Cromwell Road, London SW7 5BD, England (£60 or \$110), or from the American Association for Zoological Nomenclature, c/o NHB Stop 163, National Museum of Natural History, Washington, DC 20560 U.S.A. (\$110 or \$100 to members of A.A.Z.N.)).

Although a mandatory reference for much taxonomic work, Melville and Smith's compilation contains no indication of the major group of animals to which individual rejected names belong, and no index to these for any such group. Conserved names are allocated to animal group, and are listed in an index for each group, but rejected names are not so treated. Our goal here is thus to provide a list of all rejected names of amphibians and reptiles, inasmuch as those names, as well as available ones, need to be dealt with in synoptic taxonomic works.

Our list has been drawn exclusively from Melville and Smith (1987), except for names appearing in more recent Opinions, through no. 1517, 1988. Opinion numbers follow only those entries published after 1985. Although assuredly the survey is complete for Opinions 1369 et seq., since we have examined all of them, no attempt has been made to examine all of the first 1368 Opinions, since they are covered in Melville and Smith's work. However, our recognition of suppressed herpetological names depended upon familiarity with them or the cited works, since the major group to which any rejected name pertains was never given (and was not in earlier Indexes). Hence some names may inadvertently have been missed.

A total of 12 family-group names, 87 genus-group names and 101 species-group names is included in the following survey, alphabetically arranged in each group. Appended cross-indexes arrange all 200 names in accordance with the nine orders of amphibians and reptiles to which they belong.

Acknowledgments. We are much indebted to Drs. Kraig Adler and Jeremy D.D. Smith, who kindly reviewed the ms and offered numerous suggestions for its improvement.

## FAMILY-GROUP NAMES

**AGAMOIDEA** Fitzinger, 1826, Neue Classif. Rept.: 11, 17 (an incorrect original spelling for Agamidae Fitzinger, 1826, as

corrected; non-existent nomenclaturally; type genus Agama Daudin, 1802).

**AMBYSTOMIDAE** Hallowell, 1856, Proc. Acad. Nat. Sci. Philadelphia 8:11 (an incorrect original spelling for Ambystomatidae Hallowell, 1856, as corrected; non-existent nomenclaturally; type genus Ambystoma Tschudi, 1838).

**CAECILIIDAE** Rafinesque-Schmaltz, 1814, Specc. Sci. Giorn. Encicl. Sicilia 2:104 (an incorrect original spelling of Caeciliidae Rafinesque-Schmaltz, 1814, as corrected; non-existent nomenclaturally; type genus Caecilia Linnaeus, 1758). Op. 1462.

**CECILINIA** Rafinesque-Schmaltz, 1814, Specc. Giorn. Encicl. Sicilia 2:104 (an incorrect original spelling of Caeciliidae Rafinesque-Schmaltz, 1814, as corrected; non-existent nomenclaturally; type genus Caecilia Linnaeus, 1758). Op. 1462.

**HATTERIIDAE** Cope, 1864, Proc Acad. Nat. Sci. Philadelphia 16:227 (suppressed in priority but not homonymy contexts, making it a jr. synonym of Sphenodontidae Cope, 1870, the earliest available family-group name for its family; type genus Hatteria Gray, 1842, a jr. synonym of Sphenodon Gray, 1831).

**LEIOPELMIDAE** Turbot, 1942, Trans. Roy. Soc. New Zealand 71:247 (a jr. synonym of Leiopelmatidae Mivart, 1869; type genus Leiopelma Fitzinger, 1861).

**LIOPELMATINA** Mivart, 1869, Proc. Zool. Soc. London 1869: 291 (an incorrect original spelling of Leiopelmatidae Mivart, 1869, as corrected; non-existent nomenclaturally; type genus Leiopelma Fitzinger, 1861).

**LIOPELMIDAE** Noble, Am. Mus. Novitates (132): 9(a jr. synonym of Leiopelmatidae Mivart, 1869, as corrected; type genus Leiopelma Fitzinger, 1861).

**RHYNCHOCEPHALIDAE** Hoffmann, 1881, Bronn's Klass. Ordn. Thierr. 6(3):1065 (invalid because name of its type genus, Rhynchocephalus Owen, 1845, is a jr. homonym of Rhynchocephalus Fischer von Waldheim, 1806, a dipterous insect; Owen's name is also a jr. synonym of Sphenodon Gray, 1831, and Hoffmann's name is a jr. synonym of Sphenodontidae Cope, 1870).

**STELLIONIDAE** Bell, 1825, Zool. J. 1:457 (ruled invalid because its type genus, Stellio Daudin, 1802, was drawn from Stellio Laurenti, 1786, a nomen dubium because its type species, S. saxatilis, is unidentifiable; neither Laurenti name has been rejected, however, and Stellio has commonly been regarded as an invalid sr. synonym of Agama Daudin, 1802; the family name is likewise an invalid sr. synonym of Agamidae Gray, 1827).

**STELLIONIDAE** Gray, 1825, Ann Philos. (2) 10:196 (invalid as a jr. synonym of Stellionidae Bell, and also for the same reason that Bell's name is invalid).

**TUPINAMBIDAE** Gray, 1825, Ann. Philos., (2) 10:199 (unavailable and non-existent nomenclaturally because the name of the type genus, Tupinambis Daudin, 1802, was not regarded as valid when the family-group name was established; hence the name Tupinambidae cannot replace the currently accepted name Teiidae Gray, 1827, as it would otherwise do).

#### GENUS-GROUP NAMES

Acrodytes Fitzinger, 1843, Syst. Rept.: 30 (suppressed in priority but not in homonymy contexts, in favor of Phrynohyas Fitzinger, 1843:30, both having the same type species, Hyla venulosa Laurenti, 1768).

Ahaetulla Gray, 1825, Ann. Philos. (2)10:208 (invalid as a jr. homonym of Ahaetulla Link 1807, type species Ahaetulla mycterizans Link, 1807; a jr. synonym of the earlier Leptophis Bell, 1825, having the same type species, Coluber ahaetulla Linnaeus, 1758).

Amblystoma Agassiz, 1846, Nomencl. Zool. 6(Rept.): 2 (an unjustified emendation and jr. synonym of Ambystoma Tschudi, 1838). The date 1846, given in Opinion 649, is not correct. Vanzolini (1977:64) has shown that the probable year of publication of fascicle 6, containing reptiles and amphibians, was 1844, and certainly not later than 1845.

Amplyclephalus Kuhl and van Hasselt, 1822, Algemeene Konsten Letterbode, Haarlem 1: 101 (rejected in priority but not in homonymy contexts to protect Pareas Wagler, 1830, and Cemophora Cope, 1860, since the two species most logically assignable as type of Amplyclephalus (no species were originally included in the genus) belong one to Pareas, one to Cemophora, as now interpreted; since application of the rules of the Code was not suspended, Coluber coccineus Blumenbach, 1788, has to be accepted as the type species of Amplyclephalus, which is thus now a jr. synonym of Cemophora, of which Blumenbach's name is the type species).

Anaides Baird, 1851, Heck's Icon. Encyclop. Sci. 2:256 (an erroneous original spelling of Aneides Baird, 1851:257, hence non-existent nomenclaturally; also a jr. homonym of Anaides Westwood, 1842, for a beetle). As pointed out by Smith and Smith (1973:11) Heck's vol. 2 was not published in 1849 as

often cited, although as stated on the verso of the title page its publication was authorized in 1849.

Anodon Smith, 1829, Zool. J. 4:143 (although this name is antedated by Anodon Oken, 1815, a mollusc, the latter work has been rejected for nomenclatural purposes (Opinion 417), hence Anodon Smith is available; since its type species, Coluber scaber Linnaeus, 1758, is the same as that of Dasypeltis Wagler, 1830, Smith's name was suppressed in priority but not in homonymy contexts, thereby becoming a jr. synonym of Wagler's name).

Asthenognathus Bocourt, 1884, Bull. Sci. Soc. Philomath. Paris (7)8:149 (a jr. homonym of Asthenognathus Stimpson, 1858, a crustacean; the name is a jr. synonym of Sibon Fitzinger, 1826, since its type species, Petalognathus multifasciatus Jan, 1884, is a jr. synonym of Sibon d. dimidiata (Günther, 1872)).

Autodax Boulenger, 1887, Ann. Mag. Nat. Hist. (5)19:67 (a jr. synonym of Aneides Baird, 1851, having the same type species, Salamandra lugubris Hallowell, 1849; proposed as a substitute for Anaides Baird, 1851 (q.v.)).

Axolot Bonaparte, 1831, Giorn. Arcad. Sci. Lett. Arti 49:77 (suppressed in priority but not in homonymy contexts to protect Ambystoma Tschudi, 1838, of which it is now a jr. synonym, since its type species, Siren pisciformis Shaw, 1802, is a jr. synonym of Ambystoma mexicanum (Shaw, 1798)).

Axolotus Jarocki, 1822, Zoologia 3:179 (suppressed in priority but not in homonymy contexts to protect Ambystoma Tschudi, 1838, of which it is now a jr. synonym, since its type species, Siren pisciformis Shaw, 1802, is a jr. synonym of Ambystoma mexicanum (Shaw, 1798)).

Berus Oken, 1816, Lehrb. Naturgeschichte 3:234 (Oken's work has been rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited as a synonym of anything).

Brachypus Fitzinger, 1826, Neue Classif. Rept.:20,50 (a jr. homonym of Brachypus Meyer, 1814, a bird; a sr. but invalid synonym of Bachia Gray, 1845, since its type species is Brachypus cuvieri Fitzinger, 1826, now Bachia cuvieri (Fitzinger, 1826)).

Centrocerus Fitzinger, 1843, Syst. Rept.:18,86 (a jr. homonym of Centrocerus Swainson, 1832, a bird, hence not valid; a jr. synonym of Uromastix Merrem, 1820, since its type species is

Uromastix griseus Cuvier, 1827, a jr. synonym of Uromastix hardwickii Gray, 1827).

Chemelys Rafinesque-Schmaltz, 1832, Atl. J. and Friend of Knowledge 1:64 (suppressed in priority but not in homonymy contexts to protect Rhinoclemmys Fitzinger, 1835, of which it is now a jr. synonym, since its type species, Testudo verrucosa Walbaum, 1782, is a jr. synonym of Rhinoclemmys punctularia (Daudin, 1802)).

Cobra Laurenti, 1768, Specimen Medicum:103 (suppressed in priority but not homonymy contexts to protect Bitis Gray, 1842, of which it is now a jr. synonym, since its type species, Coluber atropos Linnaeus, 1758, now Bitis atropos (Linnaeus) is congeneric with Vipera (Echidna) arietans Merrem, 1820, the type species of Bitis and a conserved List name.)

Constrictor Laurenti, 1768, Specimen Medicum:106 (a jr. synonym of Boa Linnaeus, 1758, having the same type species, Boa constrictor Linnaeus, 1758, of which Constrictor formosissimus Laurenti, 1768, the type species of Constrictor, is a jr. synonym).

Cora Jan, 1863, Elenco Sist. Ofidi:74 (a jr. homonym of Cora Selys, 1853, for an odonate insect; type species Regina kirtlandii Kennicott; Clonophis Cope, 1888, with the same type species, is thus the oldest generic name for that species).

Coriudo Fleming, 1822, Phil. Zool. 2:271 (a jr. synonym of Dermochelys Blainville, 1816, having the same type species, Testudo coriacea Vandelli, 1761).

Crocodilus Bertrand, 1763, Dict. Univ. Foss. Propres Foss. Accid. 1:183 (Bertrand's work has been rejected for nomenclatural purposes (Opinion 592), and therefore names appearing therein do not exist in nomenclatural contexts; none should be assigned in synonymy to anything).

Dendraspis Fitzinger, 1843, Syst. Rept.:28 (suppressed in priority but not in homonymy contexts in order to protect Ophiophagus Günther, 1864, type species Hamadryas elaps Günther, 1858, a jr. synonym of Naja hannah Cantor, 1836, of which the type species of Dendraspis [not to be confused with Dendroaspis Schlegel, 1848], Naja bungarus Schlegel, 1837, is also a jr. synonym).

Dendrophis Boie, 1826, in Fitzinger, Neue Classif. Rept.:29 (a jr. synonym of Leptophis Bell, 1825, having the same type species, Coluber ahaetulla Linnaeus, 1758).



Dermatochelys Wagler, 1830, Natürl. Syst. Amph.:133 (a jr. synonym of Dermochelys Blainville, 1816, having the same type species, Testudo coriacea Vandelli, 1761).

Dermochelis Lesueur, 1829, in Cuvier, Règne Anim., Ed. 2, 2:14 (an incorrect subsequent spelling of Dermochelys Blainville, 1816; without nomenclatural status but commonly cited in synonymy of the latter name).

Diemichylus Cope, 1859, Proc. Acad. Nat. Sci. Philadelphia 11:128 (an incorrect subsequent spelling of Diemictylus Rafinesque, 1820; without nomenclatural status but commonly cited in synonymy of Notophthalmus Rafinesque, 1820, of which Diemictylus is a jr. synonym).

Diemyctelus Günther, 1901, Biologia Centrali-Am., Rept. Batr.:294 (an incorrect subsequent spelling of Diemictylus Rafinesque, 1820; without nomenclatural status but commonly cited in synonymy of Notophthalmus Rafinesque-Schmaltz, 1820, of which Diemictylus is a jr. synonym).

Diemyctylus Hallowell, 1856, Proc. Acad. Nat. Sci. Philadelphia 8:6-11 (an unjustified emendation of Diemictylus Rafinesque, 1820; an available but invalid jr. synonym of Notophthalmus Rafinesque, 1820, of which Diemictylus is a jr. synonym).

Discosomus Oken, 1816, Lehrb. Naturgeschichte 3:310 (Oken's work has been rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited as a synonym of anything).

Draco Oken, 1816, Lehrb. Naturgeschichte 3:273 (same fate as Discosomus, preceding).

Dracunculus Wiegmann, 1834, Herp. Mex.:14 (a jr. homonym of Dracunculus Reichard, 1759, a nematode; type species Draco lineatus Daudin, 1802; hence a jr. synonym of Draco Linnaeus, 1758).

Dryinus Merrem, 1820, Tent. Syst. Amph.:15,136 (a jr. homonym of Dryinus Latreille, 1804, an insect, and a jr. synonym of Ahaetulla Link, 1807, having the same type species, A. mycterizans Link, 1807).

Dryophis Dalman, 1823, Anat. Entomol.:7 (a jr. synonym of Ahaetulla Link, 1807, having the same type species, A. mycterizans Link, 1807).

Enhydrus Rafinesque-Schmaltz, 1815, Analyse Nature:77 (an incorrect subsequent spelling of Enhydris Latreille, 1802;

without nomenclatural status but commonly cited in synonymy of the latter name).

Epirhexis Cope, 1866, J. Acad. Nat. Sci. Philadelphia (2)6:96 (rejected for priority but not homonymy purposes (Opinion 1024), to protect Syrrhophus Cope, 1878, since its type species, Batrachyla longipes Baird, 1859, is congeneric with the type species of Syrrhophus, namely S. marnockii Cope, 1878).

Eremiophilus Fitzinger, 1843, Syst. Rept.:32 (rejected in priority but not homonymy contexts in order to protect Kassina Girard, 1853, both having the same type species, Cystignathus senegalensis Duméril and Bibron, 1841).

Hamadryas Cantor, 1836, Asiatick Res. 19:87 (a jr. homonym of Hamadryas Hübner, 1808, a lepidopteran insect, and a sr. synonym, although invalid, of Ophiophagus Günther, 1864, type species Naja elaps Schlegel, 1843, a jr. synonym of Hamadryas hannah Cantor, 1836, type species of its genus). The original description of Hamadryas Cantor did not appear, as indicated in Opinion 789, in 1838 (Cantor, 1838), but in 1836 (Cantor, 1836). The 1836 description named the sole species (hence the type species) H. hannah, whereas the 1838 description named the sole species H. ophiophagus, without mention of the earlier name H. hannah.

Hatteria Gray, 1842, Zool. Misc. (2):72 (a jr. synonym of Sphenodon Gray, 1831, having the same type, Hatteria punctata Gray, 1842).

Herpeton Oken, 1816, Lehrb. Naturgeschichte 3:282 (this work by Oken has been rejected for nomenclatural purposes, hence the name does not exist nomenclaturally and should not be cited as a synonym of anything).

Ibiba Gray, 1825, Ann. Phil. 10:209 (rejected in priority but not homonymy contexts to protect Boiga Fitzinger, 1826, since both have the same type species, Coluber irregularis Merrem). Op. 1374.

Liopelma Günther, 1868, Proc. Zool. Soc. London 36:478 (an unjustified emendation of Leiopelma Fitzinger, 1861; an available but invalid jr. synonym of the latter name).

Mabouia Cuvier, 1829, Règne Animal, Ed. 2, 2:62 (an incorrect subsequent spelling of Mabuya Fitzinger, 1826; without nomenclatural status but commonly cited in synonymy of the latter name).

- Mabouya Duméril and Bibron, 1839, Exp. Gén. 5:663, 671 (an incorrect subsequent spelling of Mabuya Fitzinger, 1826; without nomenclatural status but commonly cited in synonymy of the latter name).
- Mabuia Cuvier, 1829, Règne Animal, Ed. 2, 2:64 (an incorrect subsequent spelling of Mabuya Fitzinger, 1826; without nomenclatural status but commonly cited in synonymy of the latter name).
- Mabuya Rafinesque-Schmaltz, 1815, Analyse Nature:76 (a nomen nudum, without nomenclatural status, antedating but not displacing Mabuya Fitzinger, 1826).
- Notophthalma Gray, 1858, Proc. Zool. Soc. London 26:138 (an incorrect subsequent spelling of Notophthalmus Rafinesque, 1820; without nomenclatural status but commonly cited in synonymy of the latter name).
- Notophthalmia Gray, 1850, Cat. Batr. Grad. Brit. Mus.:22 (an incorrect subsequent spelling of Notophthalmus Rafinesque, 1820; without nomenclatural status but commonly cited in synonymy of the latter name).
- Notophthalmus Baird, 1850, J. Acad. Nat. Sci. Philadelphia (2)1(4):284 (an incorrect subsequent spelling of Notophthalmus Rafinesque, 1820; without nomenclatural status but commonly cited in synonymy of the latter name).
- Oedipus Tschudi, 1838, Mém. Soc. Sci. Nat. Neuchâtel 2:28 (invalid as a jr. homonym of Oedipus Berthold, 1827, for an orthopteran insect, itself suppressed in priority but not in homonymy contexts to restore order in nomenclature of the genus now accepted as Bolitoglossa Duméril, Bibron and Duméril, 1854, whose type species is Salamandra platydactylus Gray, 1831, the same as for Oedipus Tschudi).
- Ophidion Pomel, 1853, Cat. Méth. Vert. Foss. Loire:128 (a jr. homonym of Ophidion Linnaeus, 1758, for a fish; Ophidioniscus a substitute name, Kuhn, 1963; type species Ophidion antiquus Pomel, 1853; a fossil snake probably referable to Boidae).
- Palaeotriton Fitzinger, 1837, Ann. Wien. Mus. Naturgesch. 2:186 (rejected in priority but not homonymy contexts to protect Andrias Tschudi, 1837, type Salamandra scheuchzeri Holl, 1831; Palaeotriton type species Salamandra gigantea Meyer, 1832, a jr. synonym of S. scheuchzeri and also a jr. homonym of Salamandra gigantea Barton, 1808, a jr. synonym of Cryptobranchus alleganiensis Daudin, 1802).

- Palmatotriton Smith, 1945, Ward's Nat. Sci. Bull. 19(1):4 (ruled a nomen nudum, without nomenclatural status; now incorrectly cited as a jr. synonym of Bolitoglossa Duméril, Bibron and Duméril, 1854, based on Bolitoglossa rufescens (Cope, 1869)).
- Passerita Gray, 1825, Ann. Philos. 26:208 (a jr. synonym of Ahaetulla Link, 1807, both having the same type species, A. mycterizans Link, 1807).
- Petrodactylus Oken, 1816, Lehrb. Naturg. 3:index (this work by Oken has been rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited as a synonym of anything).
- Philhydrus Brookes, 1828, Prodr. Syn. Anim. Brookesian Mus.:16 (rejected in priority but not homonymy contexts to protect Ambystoma Tschudi, 1838; type species Siren pisciformis Shaw, 1802, a jr. synonym of Gyrinus mexicanus Shaw, 1798).
- Philodendros Fitzinger, 1843, Syst. Rept.:26 (rejected in priority but not homonymy contexts to protect Dromophis Peters, 1869, both having the same type species, Dendrophis praeornata Schlegel, 1837). Op. 1384.
- Philodendrus Agassiz, 1846, Nomencl. Zool. Index Univ.:285 (an unjustified emendation of Philodendros Fitzinger, 1843; an available but invalid jr. synonym of Dromophis Peters, 1869). Op. 1384.
- Phyllhydrus Gray, 1831, in Griffith's Cuvier, Anim. Kingd. 9, Syn. Spec.: 108 (rejected in priority but not homonymy contexts to protect Ambystoma Tschudi, 1838; type species Siren pisciformis Shaw, 1802, a jr. synonym of Gyrinus mexicanus Shaw, 1798).
- Propus Oken, 1816, Lehrb. Naturg. 3:287 (this work by Oken has been rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited as a synonym of anything).
- Proteocordylus Eichwald, 1831, Zoologia Specialis (3):165 (rejected in priority but not in homonymy contexts to protect Andrias Tschudi, 1837; type species P. diluvii Eichwald, 1831, a jr. synonym of Salamandra scheuchzeri Holl, 1831, type species of Andrias).
- Pterodactylus Oken, 1816, Lehrb. Naturg. 3:312 (this work by Oken has been rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited as a synonym of anything).

Rhinaspis Fitzinger, 1843, Syst. Rept.:25 (ruled a nomen nudum, as its type species, R. proboscideum Fitzinger, 1843, is a nomen nudum; neither name exists nomenclaturally but both are commonly cited in the synonymy of Simophis Peters, 1860, and its type species, Heterodon rhinostoma Schlegel, 1837; their synonymy includes Jan's 1863 occupation of Fitzinger's nomina nuda, as Rhinaspis proboscideus).

Rhinosimus Duméril, Bibron and Duméril, 1854, Erp. Gén. 7:991 (a jr. homonym of Rhinosimus Latreille, 1802, a genus of beetles; type species Rhinosimus guerini Duméril, Bibron and Duméril, 1854, now placed in Phimophis Cope, 1860, as its type species).

Rhinostoma Fitzinger, 1826, Neue Classif. Rept.:56,29 (rejected in priority but not homonymy contexts to protect Lystrophis Cope, 1885, type species Heterodon dorbignyi Duméril, Bibron and Duméril, of which Vipera nasua Wagler, 1830, is a jr. synonym through rejection in priority but not homonymy contexts; Rhinostoma was diagnosed but without an acceptable species, although two nomina nuda were named; subsequently Vipera nasua was designated type species).

Rhynchocephalus Owen, 1845, Trans. Geol. Soc. London (2)7:78 (a jr. synonym of Sphenodon Gray, 1831, having the same type species, Hatteria punctata Gray, 1842; also a jr. homonym of Rhynchocephalus Fischer von Waldheim, 1806, for a dipterous insect).

Scinci Oken, 1816, Lehrb. Naturg. 3:300 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited in the synonymy of anything).

Scincorum Oken, 1816, Lehrb. Naturg. 3:index (as in the preceding account for Scinci).

Siredon Wagler, 1830, Syst. Amph.:209,210 (rejected in priority but not in homonymy contexts to protect Ambystoma Tschudi, 1838; type species S. axolotl Wagler, 1830, a jr. synonym of Gyrinus mexicanus Shaw, 1898, now in Ambystoma).

Sirena Fischer von Waldheim, 1808, Zoognosia, Ed. 2:tab. iii (an unjustified emendation of Siren Linnaeus, 1766; an available but invalid jr. synonym of the latter name).

Sirene Oken, 1816, Lehrb. Naturg. 3:187 (this work by Oken has been rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited as a synonym of anything; the name is also an unjustified emendation of Siren Linnaeus, 1766, as well as a jr. homonym of

an earlier, identical emendation, Sirene Link, 1794, which is a citeable jr. synonym of Siren).

Sirenodon Wiegmann, 1832, in Wiegmann and Ruthe, Handb. Zool., Ed. 2:204 (rejected in priority but not homonymy contexts to protect Ambystoma Tschudi, 1838; type species Siredon axolotl Wagler, 1830, a jr. synonym of Gyrinus mexicanus Shaw, 1798, now assigned to Ambystoma).

Sphaenodon Gray, 1831, Zool. Misc. (1):14 (ruled an incorrect original spelling, non-existent nomenclaturally, of Sphenodon Gray, 1831, type species, Hatteria punctata Gray, 1842; emended first to Sphenodon by Gray, 1872, and thus accepted by the ICZN).

Sphalerosophis Jan, 1865, in Filippi, Note Viaggio Persia:356 (an incorrect original spelling and jr. synonym of Spalerosophis Jan, 1865; Sphalerosophis Jan is therefore non-existent nomenclaturally).

Sphargis Merrem, 1820, Tent. Syst. Amph.:19 (a jr. synonym of Dermochelys Blainville, 1816, having the same type species, Testudo coriacea Vandelli, 1761; in the case of Sphargis, through a jr. synonym of T. coriacea, S. mercurialis Merrem, 1820).

Stegoporus Wiegmann, 1832, in Wiegmann and Ruthe, Handb. Zool., Ed. 2:204 (rejected in priority but not homonymy contexts in order to protect Ambystoma Tschudi, 1838; proposed as a substitute for Siredon Wagler, 1830, a jr. synonym of Gyrinus mexicanus Shaw, 1898, now in Ambystoma).

Stellio Daudin, 1802, Hist. Nat. Rept. 4:5 (a jr. homonym of Stellio Laurenti, 1768, a nomen dubium because its type species, S. saxatilis, is unidentifiable; neither Laurenti name has been rejected officially, however, and Stellio Laurenti has commonly been regarded as an invalid sr. synonym of Agama Daudin, 1802).

Tachyophis Mertens, 1934, Arch. Naturg. (n.f.) 3:197 (invalid as a jr. homonym of Tachyophis Rochebrune, 1884, a fossil snake; type species Coluber pictus Gmelin, 1788, now placed in Dendrelaphis Boulenger, 1890, as a valid species).

Tapaiia Oken, 1816, Lehrb. Natur. 3:vi (index) (this work by Oken has been rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally, as of Oken, 1816, and should not be cited in the synonymy of anything, although commonly referred to Phrynosoma Wiegmann, 1828; also an incorrect original spelling of Tapaiia Oken, 1816, q. v.).

- Tapaja Oken, 1817, Isis von Oken 1817:1183 (rejected in priority but not in homonymy contexts to protect Phrynosoma Wiegmann, 1828, having the same type species, Lacerta orbicularis Linnaeus, 1758).
- Tapaja Oken, 1816, Lehrb. Naturg. 3:295 (this work by Oken was rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of anything).
- Tapaya Fitzinger, 1826, Neue Classif. Rept.:17 (rejected in priority but not homonymy contexts to protect Phrynosoma Wiegmann, 1828, having the same type species, Lacerta orbicularis Linnaeus, 1758).
- Tapaya Oken, 1817, Isis von Oken 1817:1183 (an incorrect original spelling of Tapaja Oken, 1817, hence non-existent nomenclaturally).
- Tapavia Gray, 1825, Ann. Philos. 26:197 (rejected in priority but not in synonymy contexts to protect Phrynosoma Wiegmann, 1828, having the same type species, Lacerta orbicularis Linnaeus, 1758).
- Thermophilus Fitzinger, 1843, Syst. Rept.:21 (rejected in priority but not homonymy contexts to protect Ichnotropis Peters, 1854, both having the same type species, Algyra capensis A. Smith, 1838, in Thermophilus via Fitzinger's Tropidosaura capensis "Duméril and Bibron," in error for A. Smith, and in Ichnotropis via I. macrolepidota Peters, 1854, a jr. synonym of A. capensis Smith). Op. 1422.
- Tortrix Oppel, 1811, Ann. Mus. Nat. Hist. Nat. Paris 16(95):377,381 (rejected as a jr. homonym of Tortrix Linnaeus, 1758, a lepidopteran insect; type species Anguis scytale Linnaeus, 1758, still valid, now referred as type species to the genus Anilius Oken, 1816, of which Tortrix Oppel is a sr. synonym, but invalid).
- Tritropis Fitzinger, 1843, Syst. Rept.:59 (rejected in priority but not in homonymy contexts to protect Chalarodon Peters, 1854, having the same type species, Tropidogaster blainvillii Duméril and Bibron, 1837).
- Tropidogaster Duméril and Bibron, 1837, Erp. Gen. 4:329 (rejected in priority but not in homonymy contexts to protect Chalarodon Peters, 1854, having synonymous type species; see blainvillii in the species-group list)

Typhlina Wagler, 1830, Nat. Syst. Amph.:196 (rejected in priority but not in homonymy contexts to protect Leptotyphlops Fitzinger, 1843, type species Typhlops nigricans Schlegel, 1839, in which genus Anguis septemstriatus Schneider, 1801, the type species of Typhlina, also belongs).

Zygnis Oken, 1816, Lehrb. Naturg. 3:284 (this work by Oken has been rejected for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of anything).

#### SPECIES-GROUP NAMES

alamose, Kinosternon, Pritchard, 1979, Encyclopedia of Turtles: 556 (rejected in both priority and homonymy contexts to protect K. alamosae Berry and Legler, 1980; a non-existent name, nomenclaturally, that should not be cited as a synonym of anything).

alleghaniensis, Abranchus, Harlan, 1825, Ann. Lyceum Nat. Hist. New York 1(18):271 (an unjustified emendation and jr. synonym of Salamandra alleghaniensis Daudin, 1803, now Cryptobranchus alleghaniensis (Daudin)).

alligator, Lacerta, Blumenbach, 1779, Handb. Naturg. 1:263 (rejected in priority but not in homonymy contexts to protect Crocodylus mississippiensis Daudin, 1801, now Alligator mississippiensis (Daudin), of which it is a jr. synonym).

areolata, Lacerta, Houttuyn, 1787, Anim. Mus. Houtt. Index: 24 (this work by Houttuyn was suppressed for nomenclatural purposes (Opinion 380), hence this name should not be cited in the synonymy of anything, as it does not exist nomenclaturally, and is in addition unidentifiable).

atratus, Coluber, Gmelin, 1788, in Linnaeus, Syst. Nat., Ed. 13, 1: 1103 (rejected in both priority and homonymy contexts, by Gmelin or any other author, prior to erection of Coluber atratus Hallowell, 1845, now Ninia atrata (Hallowell), in order to protect the latter name; previous usages do not exist nomenclaturally and should not be cited in the synonymy of anything).

besseri, Anguis, Andrzejowski, 1832, Nouv. Mém. Soc. Imp. Nat. Moscou (2)2:338, tab. 22, fig. 7, tab. 24 (rejected in priority but not in homonymy contexts to protect Otophis eryx colchica Nordmann, 1840, now Anguis fragilis colchica, of which A. besseri is now a jr. synonym).



- bibronii, Trapelus (Psammorrhhoa), Fitzinger, 1843, Syst. Rept.: 81 (rejected in both priority and homonymy contexts to protect Agama bibronii A. Duméril in Duméril and Bibron, 1851; Fitzinger's name was a sr. secondary homonym of Duméril's, having been transferred to the synonymy of Agama hispida aculeata Merrem, 1820; it does not now exist nomenclaturally).
- bilineatus, Psammophis moniliger, Peters, 1867, Monatsb. Akad. Wiss. Berlin 1867: 237 (rejected in priority but not in homonymy contexts to protect P. sibilans subtaeniata Peters, 1882, now P. subtaeniata Peters, of which bilineatus is now a jr. synonym).
- blainvillii, Tropidogaster, Duméril and Bibron, 1837, Erp. Gen. 4:300 (rejected in priority but not in homonymy contexts to protect Chalarodon madagascariensis Peters, 1854, of which blainvillii is now a jr. synonym).
- bosci, Rana, Bory de St. Vincent, 1828, Rés. Erp.: 266 (rejected in priority but not in homonymy contexts, to protect Rana esculenta perezii Seoane, 1885, now Rana perezii Seoane, of which the former is now a jr. synonym).
- caesius, Coluber, Cloquet, 1818, Dict. Sci. Nat. 11: 201 (rejected in priority but not in homonymy contexts to protect Coluber irregularis Leach, 1819, now Philothamnus irregularis (Leach), of which the former is now a jr. synonym).
- californiana, Aspidonectes, Rivers, 1889, Proc. California Acad. Sci. (2)2: 233-236 (rejected in priority but not in homonymy contexts to protect Trionyx steindachneri Siebenrock, 1906, of which the former is now a jr. synonym).
- chiametla, Coluber, Shaw, 1802, Gen. Zool. 3(2): 440 (rejected in priority but not in homonymy contexts to protect Herpetodryas margaritifera Schlegel, 1838, now Drymobius margaritifera (Schlegel), and Drymobius margaritifera fistulosus Smith, 1942, of which Shaw's name is now a jr. synonym).
- cincolor, Crotalus durissus, Notestein, 1905, 7th Rep. Michigan Acad. Sci.: 123 (ruled non-existent nomenclaturally because cited only in synonymy, of Crotalus horridus Linnaeus, 1758; presumably a lapsus for concolor, and presumably drawn from Jan, 1859, although the only source stated was "J").
- cinereous, Crotalus, Le Conte, 1852, in Hallowell, Proc. Acad. Nat. Sci. Philadelphia 5(5): 177 (rejected in priority but not in homonymy contexts to protect Crotalus atrox Baird and Girard, 1853, of which the former in now a jr. synonym).

- coerulea, Rana, Houttuyn, 1787, Anim. Mus. Houtt. Index: 19 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name does not exist nomenclaturally and is unidentifiable anyhow).
- colonorum, Agama, Daudin, 1802, Hist. Nat. Rept. 3: 336 (a jr. objective synonym of Lacerta agama Linnaeus, 1758, now A. a. agama (Linnaeus)).
- concolor, Crotalus durissus, Garman, 1883, Mem. Mus. Comp. Zool. 8: 175 (ruled non-existent nomenclaturally because cited only in synonymy, of C. horridus Linnaeus, 1758; name attributed to Jan, 1859).
- concolor, Crotalus durissus, Gloyd, 1940, Spec. Publ. Chicago Acad. Sci. 4:171 (ruled non-existent nomenclaturally because cited only in synonymy, of C. viridis decolor Klauber, 1930; name attributed to Jan, 1859).
- concolor, Crotalus durissus, Jan, 1859, Rev. Mag. Zool. (2)10: 153 (ruled non-existent nomenclaturally because a nomen nudum).
- cruciger, Bufo, Oken, 1816, Lehrb. Naturg. 3:209 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name is non-existent nomenclaturally and should not be cited in the synonymy of any species).
- cupreus, Coluber, Houttuyn, 1787, Anim. Mus. Houtt. Index: 28 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name does not exist nomenclaturally and is unidentifiable).
- diglossis, Chirotas, Saenz, 1869, An. Univ. Nac. Est. Unidos Colombia 1869: 63 (rejected in priority but not in homonymy contexts to protect Heteroclonium bicolor Cope, 1896, now Bachia bicolor (Cope), of which Saenz' name is now a jr. synonym). Op. 1482.
- doliatus, Coluber, Linnaeus, 1766, Syst. Nat., Ed. 12, 1:376 (rejected in priority but not in homonymy contexts to protect Coluber coccineus Blumenbach, 1788, now Cemophora coccinea (Blumenbach), of which doliatus is now a jr. synonym).
- dorsata, Testudo, Schoepff, 1801, Naturg. Schildk.: 158 (rejected in priority but not in homonymy contexts to protect Testudo punctularia Daudin, 1802, now Rhinoclemmys punctularia (Daudin), of which dorsata is now a jr. synonym).
- dracaena, Lacerta, Linnaeus, 1766, Syst. Nat., Ed. 12, 1: 250 (rejected in priority but not in homonymy contexts to protect

Tupinambis bengalensis Daudin, 1802, now Varanus bengalensis (Daudin), of which dracaena is now a jr. synonym).

dubia, Amphisbaena, Rathke, 1863, Abh. K.-Bayer. Akad. Wiss. München 9(1): 128 (rejected in both priority and homonymy contexts to protect Amphisbaena dubia Müller, 1924; Rathke's name is non-existent nomenclaturally but was based on A. fuliginosa Linnaeus, 1758, and, more precisely, A. f. amazonica Vanzolini, 1951).

elaphis, Coluber, Shaw, 1802, Gen. Zool. 3: 450 (rejected in priority but not in homonymy contexts to protect Coluber scalaris Schinz, 1822, now Elaphe scalaris (Schinz), of which Shaw's name is now a jr. synonym).

ereticauda, Triton, Eschscholtz, 1833, Zool. Atlas 5: 14 (rejected in priority but not in homonymy contexts to protect Salamandra lugubris Hallowell, 1849, now Aneides lugubris (Hallowell), of which Eschscholtz' name is now a jr. synonym).

erythronota, Salamandra, Rafinesque, 1818, Sci. J. 1: 25 (rejected in priority but not in homonymy contexts to protect Salamandra cinerea Green, 1818, now Plethodon cinereus (Green), of which Rafinesque's name is now a jr. synonym).

fasciata, Lacerta, Houttuyn, 1787, Anim. Mus. Houtt. Index: 24 (this work by Houttuyn has been rejected for nomenclatural purposes (Opinion 380), hence the name does not exist nomenclaturally, and besides is unidentifiable).

fasciata, Rana, Burchell, 1824, Travels Interior South Africa 2: 32 (rejected in priority but not in homonymy contexts to protect Rana grayi Smith, 1849, of which Burchell's name is now a jr. synonym; in addition, all other uses of Rana fasciata prior to that of Smith, 1849, are similarly rejected).

flava, Testudo, Lacépède, 1788, Hist. Nat. Quad. Ovip. Serpens 1, Synops. Meth.: 135, tab. 16 (rejected in priority but not in homonymy contexts to protect Cistudo blandingii Holbrook, 1838, now Emydoidea blandingii (Holbrook), of which Lacépède's name is now a jr. synonym).

flavescens, Amphisbaena, Houttuyn, 1787, Anim. Mus. Houtt. Index: 29 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name does not exist nomenclaturally; it is also unidentifiable).

foetidus, Coluber, Gùldenstedt, 1801, in Georgi, Geogr.-Phys. Naturh. Beschreib. Russ. Reich. 3(7): 1884 (rejected in priority but not in homonymy contexts to protect Pelias renardi

Christoph, 1861, now Vipera ursinii renardi, of which foetidus is now a jr. synonym).

formosissimus, Constrictor, Laurenti, 1768, Specimen Medicum...: 107 (a jr. objective synonym of Boa constrictor Linnaeus, 1758).

funnebris, Salamandra, Bory de St. Vincent, 1828, Rés. Exp.: 236 (rejected in priority but not in homonymy contexts to protect Pleurodeles waltl Michahelles, 1830, of which funnebris is now a jr. synonym).

galliwasp, Scincus, Oken, 1816, Lehrb. Naturg. 3: 299 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name is non-existent nomenclaturally and should not be cited in the synonymy of any species).

graecus, Stellio, Oken, 1816, Lehrb. Naturg. 3: 202 (as in the preceding account of galliwasp)

granulatus, Anguis, Houttuyn, 1787, Anim. Mus. Houtt. Index: 29 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name does not exist nomenclaturally; it is also unidentifiable).

grisea, Lacerta, Oken, 1816, Lehrb. Naturg. 3: 303 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited in the synonymy of any species).

indicus, Crocodylus vulgaris, Gray, 1831, Syn. Rept.: 58 (rejected in priority but not in homonymy contexts to protect Crocodylus palustris Lesson, 1831, now Crocodylus palustris (Lesson), of which Gray's name is now a jr. synonym).

italicus, Stellio, Oken, 1816, Lehrb. Naturg. 3: 204 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of any species).

jacapara, Coluber, Houttuyn, 1787, Anim. Mus. Houtt. Index: 26 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name does not exist nomenclaturally: it is also unidentifiable).

lancifer, Trigonocephalus, Oken, 1816, Lehrb. Naturg. 3: 270 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of any species).

- leberis, Coluber, Linnaeus, 1758, Syst. Nat., Ed. 10, 1:216 (rejected in priority but not in homonymy contexts to protect Coluber occipitomaculatus Storer, 1839 (now Storeria occipitomaculata (Storer), of which it is now a jr. synonym).
- lepidopus, Bipes, Oken, 1816, Lehrb. Naturg. 3: 249 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally and should not be cited as a synonym of any species).
- lucius, Crocodilus (Alligator), Cuvier, 1807, Ann. Mus. Nat. Hist. Nat. Paris 10: 28 (a jr. synonym of Crocodilus mississippiensis Daudin, 1801, now Alligator mississippiensis (Daudin)).
- lutescens, Triturus, Rafinesque-Schmaltz, 1832, Atlantic J. Friend of Knowledge 1: 121 (rejected in priority but not in homonymy contexts to protect Gyrinophilus porphyriticus duryi Weller, 1930, of which it is now a jr. synonym).
- marmorata, Amphisbaena, Houttuyn, 1787, Anim. Mus. Houtt. Index: 30 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name does not exist nomenclaturally and should not be cited in the synonymy of any species; it is also unidentifiable).
- maxima, Rana, Laurenti, 1768, Specimen Medicum...: 32 (a jr. synonym of Rana boans Linnaeus, 1758, now Hyla boans (Linnaeus)).
- melanepis, Coluber, Rafinesque-Schmaltz, 1814, Précis Découv. Trav. Semiolog.: 15 (rejected in priority but not in homonymy contexts to protect Coluber viridiflavus carbonarius Bonnaparte, 1833, now Hemorrhoids viridiflava carbonaria (Bonaparte), of which melanepis is now a jr. synonym).
- melanocercus, Drymarchon corais, Smith, 1941, J. Washington Acad. Sci. 31: 437, 434 (a jr. objective synonym of Spilotes melanurus Duméril, Bibron and Duméril, 1854, now Drymarchon corais melanurus (Duméril, Bibron and Duméril); originally proposed as a substitute for the latter name, supposed to be invalidated by its senior secondary homonym, Coluber melanurus Schlegel, 1837, via Spilotes melanurus (Schlegel) Gray, 1858).
- meleagris, Testudo, Shaw, 1793, Nat. Misc.: tab. 44 (rejected in priority but not in homonymy contexts to protect Cistudo blandingii Holbrook, 1838, now Emydoidea blandingii (Holbrook), of which Shaw's name is now a jr. synonym).

- mercurialis, Sphargis, Merrem, 1820, Tent. Syst. Amph.: 19 (a jr. objective synonym of Testudo coriacea Vandelli, 1761, now Dermochelys coriacea (Vandelli)).
- michahellesii, Podarcis, Fitzinger, 1864, in Erber, Verh. Zool.-Bot. Ges. Wien, 14: 703 (rejected in priority but not in homonymy contexts to protect Lacerta viridis trilineata Bedriaga, 1886, now L. trilineata (Bedriaga), of which Fitzinger's name is now a jr. synonym).
- mildei, Amphisbaena, Peters, 1878, Monatsb. K. Preuss. Akad. Wiss. 1878: 778-781 (rejected in priority but not in homonymy contexts to protect Amphisbaena trachura Cope, 1885, now A. darwini trachura, of which Peters' name is now a jr. synonym).
- minor, Testudo mydas, Suckow, 1798, Anfangsgr. Naturg. Thiere 3: 30 (rejected in priority but not in homonymy contexts to protect Thalassochelys (Colpochelys) kempii Garman, 1880, now Lepidochelys kempii (Garman), of which Suckow's name is now a jr. synonym).
- mississippiensis, Alligator, Gray, 1831, Syn. Rept.: 62 (an incorrect subsequent spelling of Crocodylus mississippiensis Daudin, 1801, now Alligator mississippiensis (Daudin), without nomenclatural status).
- mississippiensis, Crocodylus, Daudin, 1801, Hist. Nat. Rept. 2: 412 (an incorrect original spelling, non-existent nomenclaturally, of Crocodylus mississippiensis Daudin, 1801, now Alligator mississippiensis (Daudin)).
- mitrata, Rana, Houttuyn, 1787, Anim. Mus. Houtt. Index: 19 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name is nomenclaturally non-existent and should not be assigned to any species; in addition the name is unidentifiable).
- molochina, Psammophis, Berthold, 1846, Mitt. Nachr. G.A. Univ. K. Ges. Wiss. Göttingen Zool.Mus. Göttingen 1846: 143, 144 (rejected in priority but not in homonymy contexts to protect Philodryas nattereri Steindachner, 1870, of which Berthold's name is now a jr. synonym).
- monitor, Lacerta, Linnaeus, 1758, Syst. Nat., Ed. 10, 1:201 (rejected in priority but not in homonymy contexts to protect Stellio salvator Laurenti, 1768, now Varanus (V.) salvator (Laurenti), of which monitor is now a jr. synonym).
- multimaculata, Crotalus lugubris, Jan, 1863, Elenco Sist. Ofidi: 124 (rejected in priority but not in homonymy contexts to

protect Caudisona polysticta Cope, 1865, now Crotalus polystictus (Cope), of which Jan's name is now a jr. synonym).

nasua, Vipera, Wagler, 1830, Natürl. Syst. Amph.: 171 (rejected in priority but not in homonymy contexts to protect Heterodon dorbignyi Duméril, Bibron and Duméril, 1854, now Lystrophis dorbignyi (Duméril, Bibron and Duméril), of which nasua is now a jr. synonym).

neocaesariensis, Proteus, Green, 1818, J. Acad. Nat. Sci. Philadelphia, 1:358 (rejected in priority but not in homonymy contexts to protect Salamandra tigrina Green, 1825, now Ambystoma t. tigrinum (Green), of which Green's name, of 1818, is now a jr. synonym).

niger, Scytale, Daudin, 1803, Hist. Nat. Gén. Partic. Rept.: 342 (based on what is now known as Heterodon platirhinos Latreille, 1801, but rejected in both priority and homonymy contexts, to protect Scytale neuwiedii nigrum Duméril, Bibron and Duméril, 1854, now Pseudoboa nigra (Duméril, Bibron and Duméril); Daudin's name is now non-existent nomenclaturally).

nigricollis, Coluber, Dwigubskij, 1832, Opyt Estestv. Istorii 3:26 (rejected in priority but not in homonymy contexts to protect Coronella modesta Martin, 1838, now Eirenis modesta (Martin), of which Dwigubskij's name is now a jr. synonym).

oaxacae, Kinosternon, Pritchard, 1979, Encycl. Turtles: 557 (rejected in priority but not in homonymy contexts to protect Kinosternon oaxacae Berry and Iverson, 1980, of which Pritchard's name is now a jr. synonym).

ocellata, Lacerta, Houttuyn, 1787, Anim. Mus. Houtt. Index: 24 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name is nomenclaturally non-existent and should not be assigned to any species; in addition the name is unidentifiable).

ocellatus, Draco, Oken, 1816, Lehrb. Naturg. 3:277 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of any species).

operculata, Siren, Beauvois, 1799, Trans. Am. Philos. Soc. 4:277-281, figs. 1-4 (rejected in priority but not in homonymy contexts to protect Salamandra tigrina Green, 1825, now Ambystoma t. tigrinum (Green), of which Beauvois' name is now a jr. synonym).

oryzicola, Berus, Oken, 1816, Lehrb. Naturg. 3:248 (this work by Oken has been suppressed for nomenclatural purposes (Opinion

417), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of any species).

oryzivorus, Coluber, Suckow, 1798, Anfang. Theor. Angew. Naturg. Thiere 3 Amphibien: 245 (rejected in priority but not in homonymy contexts to protect Boa reticulata Schneider, 1801, now Python reticulatus (Schneider), of which oryzivorus is now a jr. synonym). Op. 1463.

oularsawa, Coluber, Bonnaterre, 1790, Tabl. Encycl. Meth. Trois Règnès Nature: 26 (rejected in priority but not in homonymy contexts, to protect Boa reticulata Schneider, 1801, now Python reticulatus (Schneider), of which oularsawa is now a jr. synonym). Op. 1463.

papillosa, Rana, Houttuyn, 1787, Anim. Mus. Houtt. Index: 19 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name is nomenclaturally non-existent, and should not be assigned to any species; in addition the name is unidentifiable).

pelamys, Hydrophis, Oken, 1816, Lehrb. Naturg. 3: 279 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of any species).

petrefactus, Crocodylus, Bertrand, 1793, Dict. Univ. Foss. Propres Foss. Accid. 1: 183 (this work by Bertrand has been suppressed for nomenclatural purposes (Opinion 592), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of any species).

philadelphicus, Axolotus, Jarocki, 1822, Zoologia 3: 179 (rejected in priority but not in homonymy contexts to protect Salamandra tigrina Green, 1825, now Ambystoma t. tigrinum (Green), of which Jarocki's name is now a jr. synonym).

planitia, Testudo, Gmelin, 1789, in Linnaeus, Syst. Nat., ed. 13, 1: 1045 (rejected in priority but not in homonymy contexts to protect Chelonura temminckii Troost, 1835, now Macrocllemys temminckii (Troost), of which planitia is now a jr. synonym).

proboscidea, Rhinostoma, Fitzinger, 1826, Neue Classif. Rept.: 56 (confirmed as a nomen nudum, hence non-existent nomenclaturally; originally intended as a name for Heterodon rhinostoma Schlegel, 1837, now Simophis rhinostoma (Schlegel)).

proboscidea, Rhinostoma (Rhinaspis), Fitzinger, 1843, Syst. Rept.: 26 (confirmed as a nomen nudum, hence non-existent nomenclaturally; originally intended as a name for Heterodon rhinostoma Schlegel, 1837, now Simophis rhinostoma (Schlegel)).



- punctato-fasciata, Lacerta muralis, Eimer, 1881, Arch. Naturg. 47(1): 368, tab. 13, fig. 10 (rejected in priority but not in homonymy contexts to protect Lacerta muralis neapolitana fiumana Werner, 1891, now Podarcis melisellensis fiumana (Werner), of which Eimer's name is now a jr. synonym).
- punctato-striata, Lacerta muralis, Eimer, 1881, Arch. Naturg. 47(1): 340, tab. 13, figs. 4,5 (rejected in priority but not in homonymy contexts to protect Lacerta muralis neapolitana fiumana Werner, 1891, now Podarcis melisellensis fiumana (Werner), of which Eimer's name is now a jr. synonym).
- quater-radiatus, Coluber, Gmelin, 1799, Naturforscher 28: 169, tab. 3, fig. 1 (rejected in priority but not in homonymy contexts to protect Coluber scalaris Schinz, 1822, now Elaphe scalaris (Schinz), of which Gmelin's name is now a jr. synonym).
- reticulata, Amphisbaena, Thunberg, 1787, D.D. Mus. Nat. Acad. Upsaliensis: 30 (rejected in priority but not in homonymy contexts to protect Amphisbaena cinerea Vandelli, 1797, now Blanus cinereus (Vandelli), of which Thunberg's name is now a jr. synonym).
- salvaquardia, Stellio, Laurenti, 1768, Specimen Medicum: 57 (rejected in priority but not in homonymy contexts to protect Tupinambis bengalensis Daudin, 1802, now Varanus bengalensis (Daudin), of which Laurenti's name is now a jr. synonym).
- sclerotica, Elaphe, Smith, 1941, Copeia, 1941: 135, 136 (Coluber subocularis Brown, 1901, now Bogertophis subocularis (Brown) ruled not invalidated by the sr. name Bascanion suboculare Cope, 1867, a jr. synonym of Masticophis m. mentovarius (Duméril, Bibron and Duméril, 1854), hence Smith's name is a jr. objective synonym of Brown's name).
- semimembranacea, Testudo, Hermann, 1804, Observ. Zool.: 219 (rejected in priority but not in homonymy contexts to protect Trionyx (Aspidonectes) sinensis Wiegmann, 1835, now Trionyx sinensis Wiegmann, of which Hermann's name is now a jr. synonym).
- sumichrasti, Henicognathus, Bocourt, 1886, Miss. Sci. Mex. (10): 628-630, pl. 41, fig. 5. (rejected in priority but not in homonymy contexts to protect Ablabes chinensis Günther, 1889, now Sibynophis chinensis (Günther), of which Bocourt's name is now a jr. synonym).
- terrestris, Testudo, Fermin, 1765, Hist. Nat. Hollande Equinox.: 51 (this work by Fermin has been suppressed for nomenclatural

purposes (Opinion 660), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of any species).

tibiatrix, Hyla, Laurenti, 1768, Spec. Medicum: 34 (rejected in priority but not in homonymy contexts to protect Rana venulosa Laurenti, 1768, now Phrynohyas venulosa (Laurenti), of which H. tibiatrix is now a jr. synonym).

timorensis, Python, Müller, 1844, Verh. Natuurl. Gesch. Ned. Overz. Bez., Land- en Volkenk. (7): 211, 221 (rejected in priority but not in homonymy contexts to protect Liasis mackloti Duméril and Bibron, 1844, of which Müller's name is now a jr. synonym).

timoriensis, Python, Müller, 1857, Reizen en Onderzoekingen in den Indischen Archipel, gedaan op last der Nederlandsche Indische Regeering, tusschen de Jaren 1828 en 1836 2: 172 (ruled an incorrect subsequent spelling of Python timorensis Müller, 1844, hence without nomenclatural status).

trimeresurus, Coluber dipsas, Oken, 1816, Lehrb. Naturg. 3:263 (this work by Oken has been suppressed for nomenclatural purposes (Opinion 417), hence the name does not exist nomenclaturally, and should not be cited in the synonymy of any species).

unicolor, Cornufer, Tschudi, 1838, Class. Batr.: 28 (Tschudi's usage, and all others prior to the proposal of Eleutherodactylus unicolor Stejneger, 1904, are rejected in both priority and homonymy contexts, to preserve Stejneger's name, which would otherwise be a jr. homonym; Tschudi's name is actually referable to Eleutherodactylus and would, if not rejected, replace Leptodactylus inoptatus Barbour, 1914, now E. inoptatus (Barbour); Tschudi's name, as type of Cornufer, requires replacement in that role to leave the name Cornufer as long interpreted, through designation of Halophila vitiensis Girard, 1853, as type species, although that species is now generally referred to the genus Platymantis Günther, 1858, along with all other species formerly referred to Cornufer; if Platymantis is split in the future, Cornufer is available if needed; thus three names are protected by rejection of Tschudi's name).

ventricosa, Emys, Gray, 1855, Cat. Shield Rept. Coll. Brit. Mus., Pt. I, Testudinata (Tortoises): 28, pl. 14 (rejected in priority but not in homonymy contexts to protect Emys cataspila Günther, 1885, now Trachemys ornata cataspila (Günther), of which Gray's name is now a jr. synonym).

- verrucosa, Rana, Houttuyn, 1787, Anim. Mus. Houtt. Index: 19 (this work by Houttuyn has been suppressed for nomenclatural purposes (Opinion 380), hence the name is nomenclaturally non-existent and should not be assigned to any species; in addition the name is unidentifiable).
- vertebralis, Leptophis, Duméril, Bibron and Duméril, 1854, Erp. Gen. 7: 543, 544 (rejected in priority but not in homonymy contexts to protect Natrix barbouri Taylor, 1922, now Macropophis barbouri (Taylor), of which L. vertebralis is now a jr. synonym).
- viridi-squamosa, Testudo, Lacépède, 1788, Hist. Nat. Quad. Ovip. Serpens 1, Syn. Meth.: 92 (rejected in priority but not in homonymy contexts to protect Thalassochelys (Colpochelys) kempii Garman, 1880, now Lepidochelys kempii (Garman), of which Lacépède's name is now a jr. synonym).
- zonata, Hyla, Spix, 1824, Anim. Nov. Test. Ran. Brasil: 41 (ruled a jr. objective synonym of Rana venulosa Laurenti, 1768, [now Phrynohyas venulosa (Laurenti)], through action of the ICZN under its plenary powers).

## LITERATURE CITED

- Cantor, Theodore Edward. 1836. Sketch of an undescribed hooded serpent (Hamadryas hannah). Asiatick Researches, 19: 87-94.
- Cantor, Theodore Edward. 1838. Notice of the Hamadryas (H. ophiophagus), a genus of hooded serpents with poisonous fangs and maxillary teeth. Proc. Zool. Soc. London, 6: 72-75.
- International Commission on Zoological Nomenclature. 1956. Opinion 380. Suppression for nomenclatorial purposes under the plenary powers of the catalogue of the collection of Martinus Houttuyn prepared under the title "Animalium Musaei Houttuiniani Index" in connection with the sale of the collection in 1787. Op. Decl. Intern. Comm. Zool. Nomencl., 12(1):1-32.
- ICZN. 1956. Opinion 417. Rejection for nomenclatorial purposes of volume 3 (Zoologie) of the work by Lorenz Oken entitled "Oken's Lehrbuch der Naturgeschichte" published in 1815-1816. Op. Decl. Intern. Comm. Zool. Nomencl., 14(1):1-42.
- ICZN. 1961. Opinion 592. Rejection of Bertrand, 1763, Dictionnaire universel des fossiles propres et des fossiles accidentels. Bull. Zool. Nomencl., 18:114-120.
- ICZN. 1963. Opinion 660. Suppression under the plenary powers of seven specific names of turtles (Reptilia, Testudines). Bull. Zool. Nomencl., 20:187-190.

- ICZN. 1966. Opinion 789. Rejection of the pamphlet by J. Hübner, 1808, entitled *Erste Zuträge zur Sammlung exotischer Schmetterlinge*. Bull. Zool. Nomencl., 23:213-220.
- ICZN. 1974. Opinion 1024. Epirhexis Cope, 1866 (Amphibia: Salientia): suppressed under the plenary powers. Bull. Zool. Nomencl., 31(3):130-132.

Melville, Richard V. and J. D. D. Smith. 1987. Official Lists and Indexes of Names and Works in Zoology. International Trust for Zoological Nomenclature. London.

Smith, Hobart M. and Rozella B. Smith. 1973. Synopsis of the Herpetofauna of Mexico. Vol. II. Lundberg; Augusta, West Virginia.

Vanzolini, Paulo E. 1977. An Annotated Bibliography of the Land and Fresh-water Reptiles of South America (1758-1975). Vol. I (1758-1900). Mus. Zool. Univ. São Paulo; São Paulo.

### CROSS-INDEXES

The following cross-indexes will facilitate scanning the 200 names here treated for those pertinent to valid names. The list is divided into two groups: *nomina clara* (names whose allocation to known taxa is apparent) and *nomina dubia* (names whose allocation to any given taxon is uncertain).

#### NOMINA CLARA

Indented names are to be found in the preceding account. CAUTION: not all indented names are synonyms of the valid names under which they appear; most are, but in some cases the valid name is simply discussed in the account for the invalid name.

#### CAUDATA

##### A. Family-Group Names

Ambystomatidae  
  Ambystomidae

##### B. Genus-group Names

Ambystoma  
  Amblystoma  
  Axolot  
  Axolotus  
  Philhydrus  
  Phyllhydrus  
  Siredon  
  Sirenodon

Stegoporus  
 Andrias  
   Palaeotriton  
   Proteocordylus  
 Aneides  
   Anaides  
   Autodax  
 Bolitoglossa  
   Oedipus  
   Palmatotriton  
 Notophthalmus  
   Diemichylus  
   Diemyctelus  
   Diemyctylus  
   Notophthalma  
   Notophthalmia  
   Notophthalmus  
 Siren  
   Sirena  
   Sirene

#### C. Species-group Names

Ambystoma tigrinum tigrinum  
   neocaesariensis, Proteus  
   operculata, Siren  
   philadelphicus, Axolotus  
 Aneides lugubris  
   ereticauda, Triton  
 Cryptobranchus alleganiensis  
   alleghaniensis, Abranchus  
 Gyrinophilus porphyriticus duryi  
   lutescens, Triturus  
 Plethodon cinereus  
   erythronota, Salamandra  
 Pleurodeles waltl  
   funnebris, Salamandra

### SALIENTIA

#### A. Family-group Names

Leiopelmatidae  
   Leiopelmidae  
   Liopelmatina  
   Liopelmidae

#### B. Genus-group Names

Kassina  
   Eremiophilus  
 Leiopelma  
   Liopelma  
 Phrynohyas

Acrodytes  
 Syrrhophus  
 Epirhexis

C. Species-group Names

Eleutherodactylus inoptatus  
     unicolor, Cornufer  
 Eleutherodactylus unicolor  
     unicolor, Cornufer  
 Hyla boans  
     maxima, Rana  
 Phrynohyas venulosa  
     tibiatrix, Hyla  
     zonata, Hyla  
 Platymantis vitiensis  
     unicolor, Cornufer  
 Rana grayi  
     fasciata, Rana  
 Rana perezii  
     bosci, Rana

GYMNOPHIONA

A. Family-group Names

Caeciliidae  
     Caeciliidae  
     Cecilia

RHYNCHOCEPHALIA

A. Family-group Names

Sphenodontidae  
     Hatteriidae  
     Rhynchocephalidae

B. Genus-group Names

Sphenodon  
     Hatteria  
     Rhynchocephalus  
     Sphaenodon

TESTUDINES

A. Genus-group Names

Dermochelys  
     Coriudo  
     Dermatochelys  
     Dermochelis  
     Sphargis

Rhinoclemmys  
Chemelys

B. Species-group Names

Dermochelys coriacea  
    mercurialis, Sphargis  
Emydoidea blandingii  
    flava, Testudo  
    meleagris, Testudo  
Kinosternon alamosae  
    alamose, Kinosternon  
Kinosternon oaxacae  
    oaxacae, Kinosternon  
Lepidochelys kempii  
    minor, Testudo mydas  
    viridi-squamosa, Testudo  
Macrochelys temminckii  
    planitia, Testudo  
Rhinoclemmys punctularia  
    dorsata, Testudo  
Trachemys ornata cataspila  
    ventricosa, Emys  
Trionyx sinensis  
    semimembranacea, Testudo  
Trionyx steindachneri  
    californiana, Aspidonectes

SAURIA

A. Family-group Names

Agamidae  
    Agamoidea  
    Stellionidae (2)  
Teiidae  
    Tupinambidae

B. Genus-group Names

Agama  
    Stellio  
Bachia  
    Brachypus  
Chalarodon  
    Tritropis  
    Tropidogaster  
Draco  
    Dracunculus  
Ichnotropis  
    Thermophilus  
Mabuya  
    Mabouia  
    Mabouya

Mabuia  
 Mabuya  
 Phrynosoma  
   Tapaia (2)  
   Tapaja  
   Tapaya (2)  
   Tapayia  
 Uromastyx  
   Centrocerus

### C. Species-group Names

Agama agama agama  
   colonorum, Agama  
 Agama bibronii  
   bibronii, Trapelus (Psammorrhoea)  
 Anguis fragilis colchica  
   besseri, Anguis  
 Bachia bicolor  
   diglossis, Chirotis  
 Chalarodon madagascariensis  
   blainvillii, Tropidogaster  
 Lacerta trilineata  
   michahellesii, Podarcis  
 Podarcis melisellensis fiumana  
   punctato-fasciata, Lacerta muralis  
   punctato-striata, Lacerta muralis  
 Varanus bengalensis  
   dracaena, Lacerta  
   salvaquardia, Stellio  
 Varanus (Varanus) salvator  
   monitor, Lacerta

## AMPHISBAENIA

### A. Species-group Names

Amphisbaena darwini trachura  
   mildei, Amphisbaena  
 Amphisbaena dubia  
   dubia, Amphisbaena  
 Amphisbaena fuliginosa amazonica  
   dubia, Amphisbaena  
 Blanus cinereus cinereus  
   reticulata, Amphisbaena

## SERPENTES

### A. Genus-group Names

Ahaetulla  
   Dryinus  
   Dryophis  
   Passerita



Anilius  
     Tortrix  
 Bitis  
     Cobra  
 Boa  
     Constrictor  
 Boiga  
     Ibiba  
 Cemophora  
     Amplycephalus  
 Clonophis  
     Cora  
 Dasypeltis  
     Anodon  
 Dendrelaphis  
     Tachyophis  
 Dromophis  
     Philodendros  
     Philodendrus  
 Enhydris  
     Enhydrus  
 Leptophis  
     Ahaetulla  
     Dendrophis  
 Leptotyphlops  
     Typhlina  
 Lystrophis  
     Rhinostoma  
 Ophidioniscus  
     Ophidion  
 Ophiophagus  
     Dendraspis  
     Hamadryas  
 Pareas  
     Amplycephalus  
 Phimophis  
     Rhinosimus  
 Sibon  
     Asthenognathus  
 Simophis  
     Rhinaspis  
 Spalerosophis  
     Sphalerosophis

#### B. Species-group Names

Boa constrictor  
     formosissimus, Constrictor  
 Bogertophis subocularis  
     sclerotica, Elaphe  
 Cemophora coccinea  
     doliatus, Coluber  
 Crotalus atrox  
     cinereous, Crotalus

*Crotalus horridus*  
     *cincolor, Crotalus durissus*  
     *concolor, Crotalus durissus*  
*Crotalus polystictus*  
     *multimaculata, Crotalus lugubris*  
*Crotalus viridis decolor*  
     *concolor, Crotalus durissus*  
*Drymarchon corais melanurus*  
     *melanocercus, Drymarchon corais*  
*Drymobius margaritiferus*  
     *chiametla, Coluber*  
*Drymobius margaritiferus fistulosus*  
     *chiametla, Coluber*  
*Eirenis modesta*  
     *nigricollis, Coluber*  
*Elaphe scalaris*  
     *elaphis, Coluber*  
     *quater-radiatus, Coluber*  
*Hemorrhoids viridiflava carbonaria*  
     *melanepis, Coluber*  
*Heterodon platirhinos*  
     *niger, Scytale*  
*Liasis mackloti*  
     *timorensis, Python*  
     *timoriensis, Python*  
*Lystrophis dorbignyi*  
     *nasua, Vipera*  
*Macropophis barbouri*  
     *vertebralis, Leptophis*  
*Ninia atrata*  
     *atratus, Coluber*  
*Philodryas nattereri*  
     *molochina, Psammophis*  
*Philothamnus irregularis*  
     *caesius, Coluber*  
*Psammophis subtaeniata*  
     *bilineatus, Psammophis moniliger*  
*Pseudoboa nigra*  
     *niger, Scytale*  
*Python reticulatus*  
     *oryzivorus, Coluber*  
     *oularsawa, Coluber*  
*Sibynophis chinensis*  
     *sumichrasti, Henicognathus*  
*Simophis rhinostoma*  
     *proboscidea, Rhinostoma*  
     *proboscidea, Rhinostoma (Rhinaspis)*  
*Storeria occipitomaculata*  
     *leberis, Coluber*  
*Vipera ursinii renardi*  
     *foetidus, Coluber*

## CROCODYLIA

## A. Species-group Names

Alligator mississippiensis  
 alligator, Lacerta  
 lucius, Crocodilus (Alligator)  
 mississippiensis, Alligator  
 mississippiensis, Crocodilus  
 Crocodylus palustris  
 indicus, Crocodilus vulgaris

NOMINA DUBIA

A few of the invalid names listed in the preceding section are nomenclaturally non-existent, but are included there because they have commonly been included in synonymies. Thirty-seven other names, not commonly cited in synonymies, and non-existent nomenclaturally (hence not obliged to be cited), follow, with author and date.

## SALIENTIA

Bufo cruciger Oken, 1816  
Rana coerulea Houttuyn, 1787  
Rana mitrata Houttuyn, 1787  
Rana papillosa Houttuyn, 1787  
Rana verrucosa Houttuyn, 1787

## TESTUDINES

Testudo terrestris Fermin, 1765

## SAURIA

Discosomus Oken, 1816  
Petrodactylus Oken, 1816  
Pterodactylus Oken, 1816  
Scinci Oken, 1816  
Scincorum Oken, 1816  
Zygnis Oken, 1816  
Anquis granulatus Houttuyn, 1787  
Bipes lepidopus Oken, 1816  
Draco ocellatus Oken, 1816  
Lacerta areolata Houttuyn, 1787  
Lacerta fasciata Houttuyn, 1787  
Lacerta grisea Oken, 1816

Lacerta ocellata Oken, 1816  
Scincus galliwaspi Oken, 1816  
Stellio graecus Oken, 1816  
Stellio italicus Oken, 1816

## AMPHISBAENIA

Propus Oken, 1816  
Amphisbaena flavescens Houttuyn, 1787  
Amphisbaena marmorata Houttuyn, 1787

## SERPENTES

Berus Oken, 1816  
Draco Oken, 1816  
Herpeton Oken, 1816  
Berus oryzicola Oken, 1816  
Coluber cupreus Houttuyn, 1787  
Coluber dipsas trimeresurus Oken, 1816  
Coluber jacapara Houttuyn, 1787  
Crotalus durissus concolor Jan, 1859  
Hydrophis pelamys Oken, 1816  
Trigonocephalus lancifer Oken, 1816

## CROCODYLIA

Crocodylus Bertrand, 1763  
Crocodylus petrefactus Bertrand, 1763









2  
07

A CATEGORIZATION AND BIBLIOGRAPHIC SURVEY  
OF PARENTAL BEHAVIOR IN  
LEPIDOSAURIAN REPTILES



LOUIS A. SOMMA

Department of Zoology  
University of Florida



SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
NO. 81

1990



**SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE**

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with the Division of Amphibians and Reptiles.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions on preparation and submission. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington DC 20560 USA. Please include a self-addressed mailing label with requests.

## INTRODUCTION

Parental behavior is a significant aspect of the life history of a wide variety of animal taxa. Parental behavior is common to the vertebrate classes Osteichthyes (Blumer, 1979, 1982; Perrone and Zaret, 1979; Baylis, 1981; Gittleman, 1981; Gross and Sargent, 1985), Amphibia (McDiarmid, 1978; Wells, 1981; Nussbaum, 1985; Duellman and Trueb, 1986), and universal within the Aves (Kendeigh, 1952; Skutch, 1957, 1976; Welty, 1982; Silver et al., 1985) and Mammalia (Kleiman and Malcolm, 1981; Dewsbury, 1985; Kleiman, 1985; Malcolm, 1985). In contrast, relatively few reptiles exhibit parental behavior (Tinkle and Gibbons, 1977; Shine and Bull, 1979; Shine, 1985, 1988). Parental behavior is common in living crocodylians (Greer, 1970, 1971; Lang, 1987; Shine, 1988) and also may have been common in extinct archosaurs (Horner and Makela, 1979; Coombs, 1982; Horner, 1982, 1984, 1987; Horner and Gorman, 1988) and cynodonts (Graves and Duvall, 1983; Duvall, 1986). Accounts of parental behavior in four turtle species (Gopherus agassizii: Barrett and Humphrey, 1986; G. flavomarginatus: Carr, 1952; Janulaw and Appleton cited in Morafka, 1982; Appleton, 1986; Ernst and Barbour, 1989; Manouria emys: Louwman, 1982; McKeown et al., 1982; Trachemys stejnegeri malonei: Hodsdon and Pearson, 1943) are remarkable, because turtles generally lack any form of parental behavior (Shine, 1988; Ernst and Barbour, 1989). The purpose of this review is to survey the various categories of parental behavior reported for lepidosaurians (lizards, snakes, amphisbaenians and a rhychocephalian) and to provide an extensive bibliography as a guide to current and future researchers.

The phrase "parental care" refers to all nongametic and postfertilization contributions of parents to the survival of their offsprings (Wittenburger, 1981; Blumer, 1982) and is construed by some (Williams, 1966; Baylis, 1981; Keenleyside, 1981; Gross and Sargent, 1985; Congdon, 1989; Spotila and O'Connor, 1989) to include viviparity and other physiological contributions. I use the phrase "parental behavior" to limit the scope of this survey to all behavioral contributions by the parent to offspring survival after oviposition or parturition. Behaviors associated exclusively with oviposition and nest construction are not included in this survey; they are probably common to most oviparous lepidosaurians (Hahn, 1909; Hilzheimer, 1910; Blanchard, 1933; Carl, 1944; Carpenter, 1966; Platt, 1969; Rand and Rand, 1976; Duvall et al., 1979; Jones and Guillette, 1982; Green and Pauley, 1987).

The term "brooding" describes behaviors of the parent while attending the nest and progeny (sensu Somma, 1988; also see Pope, 1961; Peters, 1964; Carpenter and Ferguson, 1977). Thus, brooding does not include territorial, nest-site defense wherein the parent remains at a distance from the nest, and not in or on the nest or progeny, as seen in some iguanine and gekkonid lizards, and the tuatara.

## HISTORICAL ASPECTS

The oldest conceivable documentation of parental behavior in a lepidosaurian is in the book of Isaiah, 34:15 (McDowell et al., 1982) written some time between 745-350 BC (Asimov, 1968). In this account, the Hebrew word "lilith," at one time interpreted as an owl (Strong, 1961; Asimov, 1968), is referred to as the "arrow snake." This vernacular name may have referred to the boid Eryx jaculus (Topsell, 1608; White, 1954). Since E. jaculus is viviparous, the snake reputed to "lay eggs and hatch, and gather them under her shadow" (McDowell et al., 1982) may refer to an oviparous snake or be erroneous, if not simply a fable.

Much of the pre-Twentieth century natural history literature indicates that many authors believed parental behavior was universal in snakes (Aristotle, [d. 322 BC]; Nicander of Colophon [135-133? BC]; Gesneri, 1551-1587; Topsell, 1608; Chateaubriand, 1827; Sundowner, 1895, 1902). The Second Voyage of Sinbad the Seaman, written in the 8th century AD (Burton, 1885-1888), is an example of early fiction that mentions a giant snake (undoubtedly a python) brooding its eggs. The "cockatrice" or "basilisk" was reputed to brood her eggs (Gesneri, 1551-1587; Topsell, 1608). It is likely that this mythical beast was a fantastical description of a cobra, either Ophiophagus hannah or a species of Naja (White, 1954; also see descriptions by Pliny the Elder, [d. AD 79]; Gesneri, 1551-1587; Topsell, 1608). Similarly, brooding was attributed to the "asp" (= Naja haje?) and the "dipsas" (= Bungarus sp.?; White, 1954) by Nicander. Modern documentation confirms that these taxa brood their eggs (Table VI).

Snakes have long been credited with the ability to swallow their young to protect them from danger (Topsell, 1608; Carver, 1778; Mease, 1807; Holmes, 1823; Hunter, 1824; Chateaubriand, 1827; Gosse, 1851; Rivers, 1874; Stanley, [19??]; Burroughs, 1908; Meek, 1946). Earliest accounts of this behavior are found in hieroglyphics attributed to the ancient Egyptians circa 2300 BC (Speck, 1923). So prevalent was (and still is!) this belief, that it was incorporated into early fiction, including the pre-Elizabethian poem, The Faerie Queene (Spenser, 1590). This alleged behavior, attributed not only to snakes but also to the lizard, Lacerta vivipara (Hopley, 1882), has been reviewed and discussed by numerous authors for more than 300 years (Browne, 1646; White, 1787; Hopley, 1882; Noble, 1921; Speck, 1921, 1923; Schmidt, 1929; Ditmars and Bridges, 1937; Angel, 1950; Klauber, 1972; Russell, 1983; Shine, 1988). Despite many inquiries into the plausibility of this behavior, no scientific evidence exists for its occurrence (Klauber, 1972; Shine, 1988).

Among saurians, Scincus scincus was reputed to care for its eggs and young (Gesneri, 1551-1587; Topsell, 1608), but no modern observations support this contention (Table V). Hoy (1883) suggested that all lizards brooded their eggs. In all likelihood, his statement is based upon observations of Ophisaurus attenuatus and Eumeces septentrionalis; these are the only egg-brooding lizards that he actually observed (Hoy, 1883). The suggestion that an amphisbaenian broods its eggs (Gesneri, 1551-1587; Topsell, 1608; Aldrovandi, 1640; reviewed in Druce, 1910), has not been verified.

## CATEGORIES OF PARENTAL BEHAVIOR

The various categories of parental behavior, as reported in the literature, are listed below. The symbol in parentheses identifies the categories used in Tables I and II. The literature sources are provided in Tables V and VI.

**Coil around brood (C):** The attendant parent remains coiled around or covers the brood with its body, presumably creating a physical buffer or barrier between progeny and the external environment. This is the most common form of parental behavior.

**Nest constructed and maintained (NC):** A burrow or brood chamber is constructed by the parent and maintained while attending progeny. Although this usually involves digging a depression or burrow in the substrate, Ophiophagus hannah is known to maintain a relatively complex nest chamber constructed from surrounding vegetation (Wasey, 1892; Oliver, 1956; Leakey, 1969; Whitaker, 1977).

**Defense of brood (D):** Progeny are aggressively defended by parent in the presence of conspecifics or heterospecifics.

**Passive protection (PP):** Neonatal vipers may accrue protection from the venomous female, without her exhibiting any overt signs of aggressive behavior (i.e., Crotalus horridus: W. Martin, pers. comm.).

**Thermoregulation (T):** Attendant parent uses its body to maintain a relatively constant incubation temperature for developing eggs. Most, perhaps all, pythonines are able to become low-grade endotherms while brooding through "shivering thermogenesis," thereby raising the temperature of the female's body and developing eggs above that of ambient conditions (Vinegar et al., 1970; Harlow and Grigg, 1984; Shine, 1988). However, it has been suggested that shivering thermogenesis is not practiced by all species of pythons (Vinegar et al., 1970; Ellis and Chappell, 1987; but see [Orlov], 1986; Shine, 1988). It is also possible that an attendant parent could (1) provide a passive thermal barrier between eggs and the external environment with its body or (2) bask in the sun and transfer radiantly absorbed heat from its body to its eggs (Medsger, 1919, 1932; Noble and Mason, 1933; Cogger and Holmes, 1960). Python eggs, of at least two species, that are not brooded, and subsequently incubated at lower temperatures, take longer to hatch and exhibit a higher rate of developmental anomalies (Vinegar, 1973, 1974; Branch and Patterson, 1975).

**Hydroregulation (H):** Even though it has never been demonstrated experimentally, hydroregulation has been inferred from some squamates (Fitch, 1954; Somma, 1985b; Bels and Van den Sande, 1986; [Orlov], 1986; York and Burghardt, 1988; Somma, 1989b; Somma and Fawcett, 1989). In addition, tenuous evidence suggests that two species of snakes wet their bodies with water and then lay over the eggs, thereby reducing desiccation (Elaphe obsoleta: J. Lombard, pers. comm.; Trimeresurus wiroti: Mehrtens, 1987).

**False brooding (FB):** Many species of pythons continue to brood when their clutches are removed prior to hatching (T. Miller, pers. comm.), and a Python molurus has brooded without ovipositing (J. S. Foster, pers. comm.). The nongravid female P. m. bivittatus housed with two gravid conspecifics brooded when the gravid females oviposited and brooded their eggs. This "false brooding" consisted of the python laying in a conical coil and exhibiting shivering thermogenesis (J. S. Foster, pers. comm.; Somma, pers. observ.).

**Oophagy (OO):** The parent eats eggs or aborted ova. This is a form of parental behavior because it may prevent microbial infection of viable, healthy eggs from adjacent infected eggs (Groves, 1982; Somma, 1989a) or prevent detection of progeny by predators using olfaction (i.e., detection of chemical cues released from rotting eggs or aborted ova) to locate food (Tinkle and Gibbons, 1977; Groves, 1982; Shine, 1988). Earliest observations of this behavior may be found in Hindu scriptures dating back to 600 BC (Rao, 1957).

**Removal of nonviable eggs from nest (RE):** Nonviable eggs are removed from the nest, presumably at a safe distance from viable eggs. This behavior has been reported only in Python molurus (Griehl, 1982) and perhaps occurs in Gerrhonotus liocephalus (Greene cited in Tinkle and Gibbons, 1977).

**Parental care of neonates (CN):** A parent remains with and expresses parental behavior toward neonates after hatching or parturition.

**Neonates assisted during hatching or parturition (AN):** Attending parent assists neonates from eggs, extraembryonic membranes or nest chamber. In Eumeces obsoletus and E. septentrionalis, the mother licks amniotic fluid from neonates' bodies after hatching (Evans, 1959; Somma, 1987c).

**Manipulation or retrieval of eggs (ME):** Eggs are manipulated within the nest or moved to a new nest site. Eggs that have been removed from the nest may also be retrieved.

**Communal care of eggs (CC):** Eggs may be deposited in a communal clutch and attended by several unrelated parents. In some situations, this behavior could be interpreted as alloparental care (Banks and Schwaner, 1984; Somma, 1987a).

**Neonatal feeding facilitated (FN):** A brooding female E. obsoletus avoided taking food items in deference to its young, waiting for them to finish before she fed (Evans, 1959). This behavior has not been observed in other reptiles.

**Normally does not exhibit parental behavior (NPB):** Parental behavior has been reported (sometimes reliably) for individuals of species that normally abandon their progeny.

**Details unknown (DU):** Details of parental behavior are not known or not reported.

**Reliability uncertain (RU):** Reliability of report regarded as uncertain or possibly unreliable due to paucity of information provided or observed by the author.

**Erroneous documentation (E):** Report considered unsubstantiated due to inadequate information for a species normally lacking parental behavior.

## DISCUSSION

Over 6140 extant species of lepidosaurian reptiles are currently recognized (Bellairs, 1986); of these, parental behavior has been reported for 210 species. Only 148 species (82 lizards, 65 snakes, and Sphenodon) are represented by reliable documentation or approximately 2.4% of all lepidosaurians (Table III). These species represent 17 families, excluding the unsubstantiated documentation for the Hydrophiidae, Typhlopidae and the unidentified amphisbaenid family (Tables I, II, III). Parental behavior is common in Eumeces, Phelsuma, Uromastix, anguids, xantusiids, iguanines, oviparous boids, southeast-Asian elapids, oviparous viperids, Elaphe, Farancia, and Psammophylax (Tables I and II). In most species (96%) parental behavior is entirely maternal; however, paternal and biparental behavior have been confirmed (Table IV).

The literature in this survey (Tables V and VI) demonstrates a paucity of experimentally obtained data on parental behavior in lepidosaurians (but see Noble and Mason, 1933; Hutchison et al., 1966; Vinegar et al., 1970; Van Mierop and Barnard, 1976a, 1978; Hasegawa, 1985, Somma, 1985b; [Orlov], 1986; Ellis and Chappell, 1987; York and Burghardt, 1988; Graves, 1989; Somma and Fawcett, 1989; Vitt and Cooper, 1989; Guillette et al., in review). The majority of the literature is descriptive or anecdotal, and the adaptive functions of this life history trait remain largely conjectural. Furthermore, the only ecological/evolutionary analysis devoted solely to parental behavior in reptiles is provided by Shine (1988; see Lillywhite, 1988).

## ACKNOWLEDGEMENTS

Obtaining the literature for this survey would have been impossible without the superlative library skills of J. Mundell, L. Valentine and the late M. Covault (Interlibrary Loan Department, University Library, University of Nebraska at Omaha), D. Beaubien, B. Gilbert and K. Harmon (Interlibrary Loan Department, Marston Science Library, University of Florida), and S. Tergas (Interlibrary Loan Department, Main Library West, University of Florida). I am also grateful those who provided or informed me of important references; they are E. J. Bredin, R. L. Burke, D. B. Carter, B. Clark, P. A. Cochran, H. Ehmann, D. Ferraro, L. A. Fitzgerald, J. S. Foster, H. W. Greene, L. J. Guillette, Jr., A. Hagedorn, T. Heaton-Jones, K. Horikoshi, T. R. Johnson, G. Kattan, J. W. Lang, J. F. Lokke, S. A. Minton, Jr., L. D. Moehn, J. C. Murphy, L. Nico, J. G. Robinson, H. M. Smith, N. M. Somma, S. F. Somma, S. Stewart and S. J. Walsh. I wish to thank all of the individuals who allowed me to cite their unpublished observations as personal communications.

I am indebted to D. Haney, A. Hensley and K. Horikoshi for translating the French, German and Japanese literature, respectively. I sincerely thank

H. B. Lillywhite for his insightful comments and criticism of the manuscript. Special thanks are extended to R. Shine for his helpful comments and opinions. I owe my sincerest gratitude to J. D. Fawcett for his assistance, use of his extensive herpetological library, comments and proofreading of the manuscript, his boundless encouragement and enthusiasm for this project. Lastly, I thank J. Matter for allowing me to use his word processor and G. Kiltie for her skillful preparation of this manuscript.

The cover illustration depicts a female Eumeces s. septentrionalis from Nebraska brooding eggs. I thank John F. Lokke for creating the original artwork from which the illustration was copied.

TABLE I

**Distribution of parental behavior in lizards, amphisbaenians  
and a rhynchocephalian**

| Taxa                             | Type of<br>parental behavior | Oviparous<br>or<br>viviparous | Maternal,<br>paternal or<br>biparental |
|----------------------------------|------------------------------|-------------------------------|--|
| Agamidae                         |                              |                               |  |
| <u>Leiolepis belliana</u>        | RU, CN                       | 0                             | B?                                     |
| <u>Phrynocephalus</u> sp.        | RU, CN                       | 0                             | B                                      |
| <u>Uromastix aegyptius</u>       | D                            | 0                             | M                                      |
| <u>U. ornatus</u>                | D, CN                        | 0                             | M                                      |
| Anguidae                         |                              |                               |  |
| <u>Barisia imbricata</u>         | AN                           | V                             | M                                      |
| <u>Diploglossus bilobatus</u>    | C                            | 0                             | M                                      |
| <u>D. delasagra</u>              | C                            | 0                             | P                                      |
| <u>Elgaria coerulea</u>          | AN                           | V                             | M                                      |
| <u>E. multicaudata</u>           | C, NC, D, CC                 | 0                             | M                                      |
| <u>Gerrhonotus liocephalus</u>   | C, OO?, RE?                  | 0                             | M                                      |
| <u>Mesaspis moreleti</u>         | AN                           | V                             | M                                      |
| <u>Ophisaurus apodus</u>         | C, D                         | 0                             | M                                      |
| <u>O. attenuatus</u>             | C, OO                        | 0                             | M                                      |
| <u>O. compressus</u>             | C                            | 0                             | M                                      |
| <u>O. gracilis</u>               | C                            | 0                             | M                                      |
| <u>O. harti</u>                  | C                            | 0                             | M                                      |
| <u>O. ventralis</u>              | C, ME, OO                    | 0                             | M                                      |
| Cordylidae                       |                              |                               |  |
| <u>Cordylus cataphractus</u>     | CN                           | V                             | B                                      |
| <u>C. giganteus</u>              | CN                           | V                             | M                                      |
| Gekkonidae                       |                              |                               |  |
| <u>Ailuronyx seychellensis</u>   | C, D, OO, AN?                | 0                             | M                                      |
| <u>Chondrodactylus angulifer</u> | RU, D?                       | 0                             | M                                      |
| <u>Eublepharis macularis</u>     | RU, D, NPB                   | 0                             | M                                      |
| <u>Gekko gekko</u>               | C, D, NPB?                   | 0                             | M, P                                   |
| <u>G. petricolus</u>             | C, D, OO                     | 0                             | M, P                                   |
| <u>G. smithii</u>                | RU, DU                       | 0                             | M                                      |

|                                    |                              |       |   |
|------------------------------------|------------------------------|-------|---|
| <u>Hemidactylus turcicus</u>       | D, NPB?                      | O     | M |
| <u>Hemiphyllodactylus typus</u>    | RU, DU                       | O     | M |
| <u>Naultinus grayi</u>             | CN, D                        | V     | B |
| <u>Phelsuma borbonica</u>          | D                            | O     | M |
| <u>P. dubia</u>                    | D, OO                        | O     | M |
| <u>P. flavigularis</u>             | D, OO                        | O     | M |
| <u>P. lineata</u>                  | D, OO                        | O     | M |
| <u>P. madagascariensis</u>         | D, OO                        | O     | M |
| <u>P. standingi</u>                | ME                           | O     | M |
| <u>Phyllodactylus lanei</u>        | C, CC                        | O     | M |
| <u>Ptychozoon lionotum</u>         | C, D                         | O     | M |
| <u>Teratoscincus scincus</u>       | RU, NPB?                     | O     | M |
| Iguanidae                          |                              |       |   |
| <u>Amblyrhynchus cristatus</u>     | D                            | O     | M |
| <u>Brachylophus fasciatus</u>      | D                            | O     | M |
| <u>B. vitiensis</u>                | D                            | O     | M |
| <u>Conolophus pallidus</u>         | D                            | O     | M |
| <u>C. subcristatus</u>             | D                            | O     | M |
| <u>Crotaphytus collaris</u>        | RU, NPB                      | O     | M |
| <u>Cyclura carinata</u>            | D                            | O     | M |
| <u>C. cornuta</u>                  | D                            | O     | M |
| <u>C. cychlura</u>                 | D                            | O     | M |
| <u>C. nubila</u>                   | D                            | O     | M |
| <u>Iguana iguana</u>               | D                            | O     | M |
| <u>Phrynosoma douglassi</u>        | RU, D, NPB                   | V     | M |
| <u>Sauromalus varius</u>           | D                            | O     | M |
| <u>Sceloporus undulatus</u>        | E, CN                        | O     | B |
| Lacertidae                         |                              |       |   |
| <u>Acanthodactylus scutellatus</u> | RU, CN                       | O     | M |
| <u>Lacerta viridis</u>             | D?, ME                       | O     | M |
| Scincidae                          |                              |       |   |
| <u>Calyptotis scutirostrum</u>     | ME, NC                       | O     | M |
| <u>Corucia zebzata</u>             | CN, D, AN?                   | V     | M |
| <u>Cyclodina pseudornata</u>       | AN                           | V     | M |
| <u>Egernia cunninghami</u>         | AN, D                        | V     | M |
| <u>E. striata</u>                  | RU, CN                       | V     | M |
| <u>E. whitii</u>                   | RU, CN                       | V     | ? |
| <u>Emoia cyanura</u>               | RU, CC?                      | O     | M |
| <u>Eumeces anthracinus</u>         | C, D, NC, OO                 | O     | M |
| <u>E. callicephalus</u>            | C, OO, NC, ME                | O, V* | M |
| <u>E. chinensis</u>                | C, NC                        | O     | M |
| <u>E. copei</u>                    | CN                           | V     | M |
| <u>E. egregius</u>                 | C, D, NC, CC                 | O     | M |
| <u>E. elegans</u>                  | C                            | O     | M |
| <u>E. fasciatus</u>                | CN, C, D, ME, NC, OO, H?, CC | O     | M |
| <u>E. inexpectatus</u>             | C, OO, CC                    | O     | M |
| <u>E. laticeps</u>                 | C, D, NC, ME, OO, CN, CC     | O     | M |
| <u>E. latiscutatus</u>             | C, CN                        | O     | M |
| <u>E. lynxe</u>                    | CN                           | V     | M |
| <u>E. multivirgatus</u>            | C, CN?, CC                   | O     | M |
| <u>E. obsoletus</u>                | C, ME, NC, CN, FN, D, OO     | O     | M |



|  |                                 |   |   |
|--|---------------------------------|---|---|
| <u>E. okadae</u>                           | C, NC, OO, ME, CC               | O | M |
| <u>E. oshimensis</u>                       | DU                              | O | M |
| <u>E. parviauriculatus</u>                 | CN                              | O | M |
| <u>E. quadrilineatus</u>                   | DU                              | O | M |
| <u>E. schneiderii</u>                      | C                               | O | M |
| <u>E. septentrionalis</u>                  | C, D, NC, H, ME, CN, CC, OO, AN | O | M |
| <u>E. skiltonianus</u>                     | C, D, NC, ME, CN?               | O | M |
| <u>E. stimsoni</u>                         | RU, DU                          | O | M |
| <u>E. tetragrammus</u>                     | C                               | O | M |
| <u>E. xanthi</u>                           | C                               | O | M |
| <u>Lampropholis mustelina</u>              | ME, NC                          | O | M |
| <u>Leiolopisma otagense</u>                | AN                              | V | M |
| <u>L. smithi</u>                           | AN                              | V | M |
| <u>L. zia</u>                              | ME, NC                          | O | M |
| <u>Mabuya capensis</u>                     | AN                              | V | M |
| <u>M. macrorhyncha</u>                     | AN                              | V | M |
| <u>M. macularia</u>                        | RU, DU                          | O | M |
| <u>Neoseps reynoldsi</u>                   | C                               | O | M |
| <u>Scincus scincus</u>                     | E?, CN                          | O | M |
| <u>Sphenomorphus quoyii</u>                | AN                              | V | M |
| <u>Tiliqua rugosa</u>                      | AN                              | V | M |
| <br>Teiidae                                |                                 |   |   |
| <u>Tupinambis teguixin</u>                 | C, D, CN?, NC, AN?              | O | M |
| <u>T. rufescens</u>                        | NC, D?                          | O | M |
| <br>Varanidae                              |                                 |   |   |
| <u>Varanus gouldii</u>                     | E, AN                           | O | ? |
| <u>V. griseus</u>                          | RU, DU                          | O | M |
| <u>V. komodoensis</u>                      | RU, AN, NPB?                    | O | M |
| <u>V. mitchelli</u>                        | RU, C                           | O | M |
| <u>V. salvator</u>                         | RU, D?, NPB                     | O | M |
| <u>V. varius</u>                           | AN                              | O | M |
| <br>Xantusiidae                            |                                 |   |   |
| <u>Xantusia henshawi</u>                   | AN                              | V | M |
| <u>X. vigilis</u>                          | AN                              | V | M |
| <br>Trogonophidae or<br>Amphisbaenidae (?) |                                 |   |   |
| 'Amphisbaina'<br>(= unidentified species)  | RU, C                           | O | M |
| <br>Sphenodontidae                         |                                 |   |   |
| <u>Sphenodon punctatus</u>                 | D                               | O | M |

\*One instance of viviparity in E. callicephalus has been reported by Taylor (1985).

TABLE II

## Distribution of parental behavior in snakes

| Taxa   | Type of parental behavior    | Oviparous or viviparous | Maternal, paternal or biparental |
|--|------------------------------|-------------------------|----------------------------------|
| Boidae   |                              |                         |                                  |
| <u>Aspidites melanocephalus</u>                        | C, T                         | O                       | M                                |
| <u>Boa constrictor</u>                                 | RU, C, D, NPB                | V                       | M                                |
| <u>Casarea dussemieri</u>                              | RU, C                        | O                       | M                                |
| <u>Chondropython viridis</u>                           | C, T, D, H                   | O                       | M                                |
| <u>Epicrates cenchria</u>                              | AN, OO, D, CN                | V                       | M                                |
| <u>E. striatus</u>                                     | OO                           | V                       | M                                |
| <u>E. subflavus</u>                                    | E, C                         | V*                      | M                                |
| <u>Eunectes murinus</u>                                | AN, OO                       | V                       | M                                |
| <u>E. notaeus</u>                                      | AN, OO                       | V                       | M                                |
| <u>Liasis albertisii</u>                               | C, D, T                      | O                       | M                                |
| <u>L. boa</u>  | C                            | O                       | M                                |
| <u>L. childreni</u>                                    | C                            | O                       | M                                |
| <u>L. fuscus</u>                                       | C, T, D, H                   | O                       | M                                |
| <u>L. olivaceus</u>                                    | C                            | O                       | M                                |
| <u>L. papuanus</u>                                     | C, T                         | O                       | M                                |
| <u>L. perthensis</u>                                   | C                            | O                       | M                                |
| <u>Morelia amethystina</u>                             | C, T, CC                     | O                       | M                                |
| <u>M. bredli</u>                                       | C                            | O                       | M                                |
| <u>M. spilota</u>                                      | C, T, H, D, NC               | O                       | M                                |
| <u>Python anchietae</u>                                | C, T                         | O                       | M                                |
| <u>P. curtus</u>                                       | C, T                         | O                       | M                                |
| <u>P. molurus</u>                                      | C, T, D, FB, RE, CN?, H?, ME | O                       | M                                |
| <u>P. regius</u>                                       | C, T, D, H, ME               | O <sup>+</sup>          | M                                |
| <u>P. reticulatus</u>                                  | C, T?, D                     | O                       | M                                |
| <u>P. sebae</u>  | C, T?, D, ME, RE             | O                       | M                                |
| <u>P. timoriensis</u>                                  | C, T                         | O                       | M                                |
| 'Lilith' or 'arrow snake'<br>(= <u>Eryx jaculus?</u> ) | E?, C                        | V                       | M                                |
| Colubridae   |                              |                         |                                  |
| <u>Ahaetulla nasuta</u>                                | OO                           | V                       | M                                |
| <u>Amphiesma stolata</u>                               | C                            | O                       | M                                |
| <u>Atretium schistosum</u>                             | RU, C                        | O                       | M                                |
| <u>Cemophora coccinea</u>                              | E, C, OO                     | O                       | M                                |
| <u>Cerberus rynchops</u>                               | RU, CN                       | V                       | M                                |
| <u>Clelia clelia</u>                                   | C, H?                        | O                       | M                                |
| <u>Coronella austriaca</u>                             | RU, CN                       | V                       | M                                |
| <u>Diadophis punctatus</u>                             | C, NPB, CC?                  | O                       | M                                |
| <u>Elaphe climacophora</u>                             | C                            | O                       | M                                |
| <u>E. flavolineata</u>                                 | C                            | O                       | M                                |
| <u>E. guttata</u>                                      | RU, C, NPB                   | O                       | M                                |
| <u>E. obsoleta</u>                                     | C, D, H?, NPB, CC?           | O                       | M, B?                            |
| <u>E. quadrivirgata</u>                                | C, D                         | O                       | M                                |
| <u>E. quatuorlineata</u>                               | RU, C, NPB                   | O                       | M                                |
| <u>E. schrenki</u>                                     | RU, C                        | O                       | M                                |

|  |                 |       |       |
|--|-----------------|-------|-------|
| <u>Farancia abacura</u>  | C, NC, CN?      | 0     | M     |
| <u>F. erythrogramma</u>  | RU, C           | 0     | M     |
| <u>Heterodon platirhinos</u>   | E, C, D         | 0     | M     |
| <u>Hydrodynastes gigas</u>   | RU, C           | 0     | M     |
| <u>Lampropeltis triangulum</u>   | RU**, C, NPB    | 0     | M     |
| <u>Lycodon aulicus</u>   | RU, C           | 0     | M     |
| <u>L. striatus</u>   | RU, C           | 0     | M     |
| <u>Masticophis flagellum</u>   | E, CN           | 0     | B     |
| <u>Natrix natrix</u>   | C, D, NPB       | 0     | M     |
| <u>Oligodon taeniolatus</u>  | RU, C           | 0     | M     |
| <u>Opisthotropis latouchii</u>   | C, H?           | 0     | M     |
| <u>Pituophis catenifer</u>   | E, C, D         | 0     | M     |
| <u>Psammophylax rhombeatus</u>   | C, CC           | 0     | M     |
| <u>P. tritaeniatus</u>   | C               | 0     | M     |
| <u>P. variabilis</u>   | C               | 0 (V) | M     |
| <u>Ptyas korros</u>  | RU, C           | 0     | M     |
| <u>P. mucosus</u>  | C               | 0     | M     |
| <u>Rhabdophis subminiata</u>   | C               | 0     | M     |
| <u>Sinonatrix percarinata</u>  | C               | 0     | M     |
| <u>Tropidonophis mairii</u>  | C               | 0     | M     |
| <u>Xenochrophis piscator</u>   | C, NC           | 0     | M     |
| <u>Elapidae</u>  |                 |       |       |
| <u>Bungarus caeruleus</u>  | C               | 0     | M     |
| <u>B. candidus</u>   | C               | 0     | M     |
| <u>B. ceylonicus</u>   | C, CC?, NC, CN? | 0     | B?    |
| <u>B. fasciatus</u>  | C, CN?          | 0     | M, B? |
| 'Dipsas' (= <u>Bungarus</u> sp.?)  | RU, CN?         | 0     | M     |
| <u>Calliophis maculiceps</u>   | RU, D?          | 0     | M     |
| <u>Demansia papuensis</u>  | RU, DU          | 0     | M     |
| <u>Micrurus fulvius</u>  | RU, C, NPB      | 0     | M     |
| <u>M. psyches</u>  | RU, DU          | 0     | M     |
| <u>Naja melanoleuca</u>  | C               | 0     | M, B? |
| <u>N. naja</u>   | C, D, NC, OO    | 0     | M, B  |
| <u>N. nigricollis</u>  | RU, DU          | 0     | M     |
| 'Asp' (= <u>Naja</u> <u>haje</u> ?)  | RU, D           | 0     | M     |
| <u>Ophiophagus hannah</u>  | C, NC           | 0     | M     |
| 'Cockatrice' or 'basilisk'<br>(= <u>Ophiophagus hannah</u> or<br><u>Naja</u> sp.?) | RU, C           | 0     | M     |
| <u>Pseudechis butleri</u>  | RU, C, D        | 0     | M     |
| <u>Pseudonaja textilis</u>   | RU, C, NPB      | 0     | M     |
| <u>Hydrophiidae</u>  |                 |       |       |
| <u>Pelamis platurus</u>  | E, CN, C        | V     | M     |
| <u>Laticaudidae</u>  |                 |       |       |
| <u>Laticauda colubrina</u>   | C, D, CN?       | 0     | M     |
| <u>L. semifasciata</u>   | RU, DU          | 0     | ?     |
| <u>Leptotyphlopidae</u>  |                 |       |       |
| <u>Leptotyphlops dulcis</u>  | C               | 0     | M     |
| <u>L. humilis</u>  | E?, C           | 0     | M     |

|                                       |             |       |   |
|---------------------------------------|-------------|-------|---|
| Typhlopidae                           |             |       |   |
| <u>Ramphotyphlops</u> <u>braminus</u> | RU, C, NPB  | O     | M |
| <u>Rhinotyphlops</u> <u>caecus</u>    | RU, DU      | O     | M |
| Viperidae                             |             |       |   |
| <u>Agkistrodon</u> <u>contortrix</u>  | RU, CN, D   | V     | M |
| <u>A. piscivorus</u>                  | RU, CN, D?  | V     | M |
| <u>Calloselasma</u> <u>rhodostoma</u> | C, H, D     | O     | M |
| <u>Causus</u> <u>rhombeatus</u>       | C           | O     | M |
| <u>Crotalus</u> <u>sp.</u>            | RU, CN      | V     | M |
| <u>C. adamanteus</u>                  | E?, CN, D   | V     | M |
| <u>C. atrox</u>                       | CN, PP      | V     | M |
| <u>C. horridus</u>                    | CN, PP      | V     | M |
| <u>C. viridis</u>                     | CN, D, PP   | V     | M |
| <u>Deinagkistrodon</u> <u>acutus</u>  | C           | O     | M |
| <u>Lachesis</u> <u>muta</u>           | C, D        | O     | M |
| <u>Porthidium</u> <u>nummifer</u>     | RU, CN      | V     | M |
| <u>Sistrurus</u> <u>catenatus</u>     | RU, CN      | V     | M |
| <u>Trimeresurus</u> <u>kaulbacki</u>  | DU, D?      | O     | M |
| <u>T. monticola</u>                   | C, D        | O     | M |
| <u>T. okinavensis</u>                 | C, D        | O (V) | M |
| <u>T. wiroti</u>                      | C, H?       | O     | M |
| <u>Vipera</u> <u>aspis</u>            | RU, CN, PP? | V     | M |
| <u>V. berus</u>                       | CN, PP      | V     | M |

\*This species is viviparous (Lynn and Grant, 1940), not oviparous as reported by Gosse (1851). Perhaps Gosse confused an oviparous colubrid for E. subflavus.

\*\*Female L. triangulum remain coiled around their eggs for a brief period after oviposition in order to compress them into an adherent mass before abandoning them (McCauley, 1945; Green and Pauley, 1987). This may account for the considerable number of references (Table VI) suggesting that this species broods its eggs.

+There is an unverified report of viviparity in P. regius (Anonymous, 1941).

TABLE III

Summary of the number of species and genera exhibiting parental behavior within lepidosaurian taxa (based on Tables I and II)

| Taxa                                | Number* of genera | Number* of species |
|-------------------------------------|-------------------|--------------------|
| Squamata                            | 107 (69)          | 209 (147)          |
| Amphisbaenia                        | 1 (0)             | 1 (0)              |
| Trogonophidae or Amphisbaenidae (?) | 1 (0)             | 1 (0)              |
| Sauria                              | 48 (36)           | 104 (82)           |
| Agamidae                            | 3 (1)             | 4 (2)              |
| Anguidae                            | 6 (6)             | 13 (13)            |

|                  |         |          |
|------------------|---------|----------|
| Cordylidae       | 1 (1)   | 2 (2)    |
| Gekkonidae       | 11 (7)  | 18 (13)  |
| Iguanidae        | 9 (6)   | 14 (11)  |
| Lacertidae       | 2 (1)   | 2 (1)    |
| Scincidae        | 13 (11) | 41 (35)  |
| Teiidae          | 1 (1)   | 2 (2)    |
| Varanidae        | 1 (1)   | 6 (1)    |
| Xantusiidae      | 1 (1)   | 2 (2)    |
| Serpentes        | 58 (33) | 104 (65) |
| Boidae           | 10 (7)  | 28 (24)  |
| Colubridae       | 25 (14) | 36 (20)  |
| Elapidae         | 8 (3)   | 14 (7)   |
| Hydrophiidae     | 1 (0)   | 1 (0)    |
| Laticaudidae     | 1 (1)   | 2 (1)    |
| Leptotyphlopidae | 1 (1)   | 2 (1)    |
| Typhlopidae      | 2 (0)   | 2 (0)    |
| Viperidae        | 10 (7)  | 19 (12)  |
| Rhynchocephalia  | 1 (1)   | 1 (1)    |
| Sphenodontidae   | 1 (1)   | 1 (1)    |

\*Number in parentheses represents actual total when erroneous and uncertain accounts are deleted from analysis (see Tables I and II).

TABLE IV

Prevalence of maternal, paternal and biparental behavior in lepidosaurians

| Parent exhibiting behavior | Number* of species | Percent* of total |
|----------------------------|--------------------|-------------------|
| Maternal only              | 193 (142)          | 92.0 (96)         |
| Paternal only              | 1 (1)              | 0.5 (0.7)         |
| Biparental only            | 7 (2)              | 3.0 (1.3)         |
| Maternal and biparental    | 4 (1)              | 2.0 (0.7)         |
| Paternal or maternal       | 2 (2)              | 1.0 (1.3)         |
| Sex of parent unknown      | 3 (0)              | 1.5 (0)           |

\*Number in parentheses represents actual value when erroneous and uncertain sources are deleted from analysis.

TABLE V

Bibliographic sources for reports of lizard, amphisbaenians and  
rhynchocephalian parental behavior

| Taxa                             | Sources  |
|----------------------------------|--|
| Agamidae                         |  |
| <u>Leiolepis belliana</u>        | Boulenger, 1903  |
| <u>Phrynocephalus</u> sp.        | Bertin and Burton, 1967  |
| <u>Uromastix aegyptius</u>       | Mendelssohn and Bouskila, 1989; H. Mendelssohn, pers. comm.  |
| <u>U. ornatus</u>                | Mendelssohn and Bouskila, 1989; G. Perry, pers. comm.; H. Mendelssohn, pers. comm.                           |
| Anguidae                         |  |
| <u>Barisia imbricata</u>         | Guillette and Hotton, 1986   |
| <u>Diploglossus bilobatus</u>    | Taylor, 1956   |
| <u>D. delasagra</u>              | Barbour and Ramsden, 1919  |
| <u>Elgaria coerulea</u>          | Stewart in Guillette and Hotton, 1986, and pers. comm.   |
| <u>E. multicarinata</u>          | Langerwerf, 1981; Jes, 1987  |
| <u>Gerrhonotus liocephalus</u>   | Greene and Dial, 1966; Greene in Tinkle and Gibbons, 1977  |
| <u>Mesaspis moreleti</u>         | Greene in Guillette and Hotton, 1986   |
| <u>Ophisaurus apodus</u>         | Petzold, 1971; Langerwerf, 1981, 1984; Claffey and Johnson, 1982a, b; Huff, 1985                             |
| <u>O. attenuatus</u>             | ?Hoy, 1883; Collins, 1959; Blair, 1961; Fitch, 1970, 1986, 1989; Vogt, 1981                                  |
| <u>O. compressus</u>             | Bartlett, 1985   |
| <u>O. gracilis</u>               | Wall, 1908; Smith, 1935; Jayaram, 1974; Daniel, 1983   |
| <u>O. harti</u>                  | Pope, 1929, 1955   |
| <u>O. ventralis</u>              | Noble and Mason, 1932, 1933; Telford, 1952; Vinegar, 1968; Villiard, 1969; Mount, 1975; Somma, pers. observ. |
| Cordylidae                       |  |
| <u>Cordylus cataphractus</u>     | Branch, 1988; S. Jacobs, pers. comm.   |
| <u>C. giganteus</u>              | Patterson and Bannister, 1987; S. Jacobs, pers. comm.  |
| Gekkonidae                       |  |
| <u>Ailuronyx sechellensis</u>    | High, [1976]; Miller, 1980; McKeown and Miller, 1985; Slavens, 1987  |
| <u>Chondrodactylus angulifer</u> | Miller, 1983a  |
| <u>Eublepharis macularis</u>     | Miller, 1980   |
| <u>Gekko gecko</u>               | Honegger, 1969; Koch, 1972; Zaworski, 1987a, c, 1988; T. Miller, pers. comm.                                 |
| <u>G. petricolus</u>             | Zaworski, 1987a, b   |
| <u>G. smithii</u>                | Tho and Ho, 1979   |
| <u>Hemidactylus turcicus</u>     | Somma, pers. obs.  |
| <u>Hemiphyllodactylus typus</u>  | Eijsden, 1978  |
| <u>Naultinus grayi</u>           | Robb, 1980; J. Fawcett, pers. comm.  |

- Phelsuma borbonica Miller, 1982  
P. dubia Osadnik, 1984  
P. flavigularis Osadnik, 1984  
P. lineata Osadnik, 1984  
P. madagascariensis Osadnik, 1984; Rösler, 1988  
P. standlingi Digney and Tytle, 1983  
Phyllodactylus lanei Z. Uribe, pers. comm.; A. Ramírez, pers. comm.  
Ptychozoon lionotum Waitkus, 1983; Tytle et al., 1987  
Teratoscincus scincus Miller, 1983b
- Iguanidae
- Amblyrhynchus cristatus Heller, 1903; Eibl-Eibsfeldt, 1966; Trillmich, 1979; Fitch, 1982; Dellinger, 1989  
Brachylophus fasciatus Cogger, 1974; Gibbons and Watkins, 1982  
B. vitiensis Gibbons and Watkins, 1982; Gibbons, 1984/85  
Conolophus pallidus Christian and Tracy, 1982  
C. subcristatus Werner, 1982  
Crotaphytus collaris Burt and Hoyle, 1934  
Cyclura carinata Iverson, 1977, 1979  
C. cornuta Shaw, 1969; Wiewandt, 1977, 1979; Boylan, 1984  
C. cychlura Carey, 1975  
C. nubila Shaw, 1954; Crutchfield, 1982, 1986; Thompson in Blair, 1983a, b  
Iguana iguana Alvarez del Toro, 1960; Mertens, 1960; Wiewandt, 1982; Ellison, 1985  
Phrynosoma douglassi Lockwood, 1883  
Sauromalus varius Lawler and Jarchow, 1986; Lawler in Gilbert, 1987; Castillo S., 1989  
Sceloporus undulatus Hay, 1892
- Lacertidae
- Acanthodactylus scutellatus G. Perry, pers. comm.  
Lacerta viridis Mertens, 1960; Burton and Burton, 1984
- Scincidae
- Calyptotis scutirostrum Ehmann, 1988  
Corucia zebrata Hediger, 1937, 1986; Slavens, 1983; Honegger, 1985; Mehaffey, 1986; Peterson, 1986; A. Anderson, pers. comm.  
Cyclodina pseudornata J. Fawcett, pers. comm.  
Egernia cunninghami Niekisch, 1975, 1980; Zimmermann, 1986  
E. striata Pianka and Giles, 1982  
E. whitii McPhee, 1979  
Emoia cyanura J. Fitch in Fitch, 1970  
Eumeces anthracinus Clausen, 1938; Dowling, 1950; Hamilton, 1958; Anderson, 1965; Cooper et al., 1973; Collins, 1975  
E. callicephalus Campbell and Simmons, 1961; Zweifel, 1962; Williamson, 1986; Tanner, 1987, and pers. comm.  
E. chinensis [Wang, 1966]  
E. copei L. Guillette, pers. comm.  
E. egregius Hamilton and Pollack, 1958; Mount, 1961, 1963; Somma, pers. observ.

- E. elegans  
E. fasciatus\*
- Mell, 1929; Hikida, 1981  
 Ditmars, 1904, 1907; Allard, 1909; Ruthven, 1911; Dunn, 1920; Blanchard, 1922; Bishop, 1926; Burt, 1928, 1937; Corrington, 1929; Klots, 1930; Noble and Mason, 1932, 1933; Burt and Burt, 1935, Taylor, 1935; Conant, 1938, 1951; McCauley, 1939, 1945; Cagle, 1940; Anderson, 1942, 1965; McClellan et al., 1943; Minton, 1944, 1972; H. Smith, 1946; Neill, 1948; Evans and Roeker, 1951; Fitch, 1954, 1967; Parmalee, 1955; Kennedy, 1956; Reynolds, 1959; Tinkle, 1959, P. Smith, 1961; Leviton, [1972]; Snyder, 1972; Burghardt, 1973; Mount, 1975; Fitch and von Achen, 1977; Vogt, 1981; Groves, 1982; Lang, 1982, 1983; Cooper et al., 1983; Cooper and Vitt, 1985; Stewart and Duvall, 1985; Vitt and Cooper, 1986, 1989; Green and Pauley, 1987; Johnson, 1987; Cochran, 1989; Somma, pers. observ.
- E. inexpectatus
- Smith, 1946; Duellman and Schwartz, 1958; Hamilton, 1958; Molchos, 1971; ?Loop and Scoville, 1972; Mount, 1975; Vitt and Cooper, 1986; Dundee and Rossman, 1989; Somma, pers. observ.
- E. laticeps
- Hurter, 1911; Noble and Mason, 1932, 1933; Taylor, 1935; Cook, 1943; McClellan et al., 1943; Mansueti, 1948; Martof, 1956; Smith, 1961; Mount, 1975; Johnson, 1979; Moehn, 1980; Schuette, 1980; Ashton and Ashton, 1985; Cooper and Vitt, 1985; Hammond, 1985; Vitt and Cooper, 1985a, b, 1989; Green and Pauley, 1987; Johnson, 1987; Meshaka et al., 1988; Somma, pers. observ.
- E. laticutatus  
E. lynxe  
E. multivirgatus
- Sengoku, 1979; Hikida, 1981; [Mathui, 1985]  
 L. Guillette, pers. comm.  
 Gehlbach, 1965; Van Devender and Van Devender, 1975; A. Aquino, pers. comm.; J. Lynch, pers. comm.
- E. obsoletus
- Taylor, 1935; Smith, 1946; Fitch, 1955, 1956, 1964, 1967, 1970; Fouquette and Lindsay, 1955; Evans, 1959; Hall, 1972; Hall and Fitch, 1972; J. Lynch, pers. comm.; Somma, pers. observ.
- E. okadae
- [Hikida, 1975]; Sengoku, 1979; Hasegawa, 1984, 1985, and pers. comm.; [Mathui, 1985]
- E. oshimensis  
E. parviauriculatus  
E. quadrilineatus  
E. schneiderii  
E. septentrionalis
- Toyama, 1975  
 Tanner, 1987, and pers. comm.  
 Mell, 1929  
 Zimmermann, 1986  
 ?Hoy, 1883; Breckenridge, 1941, 1943, 1944; Smith and Slater, 1949; Nelson, 1963; Bredin, 1981, 1989 [pers. comm.]; Lang, 1982, 1983; [Gerholdt], 1984b; Somma, 1985a, b, c, 1987a, b, c, 1989a; Somma and Fawcett, 1985, 1989; McAllister, 1987; (reviewed in Somma and Cochran, 1989)



- E. skiltonianus Heller in McClain, 1899\*\*; Van Denburgh, 1922; Woodbury, 1931; Tanner, 1943, 1957
- E. stimsoni Taylor, 1935
- E. tetragrammus Strecker, 1908; Werler, 1951; Behler and King, 1979
- E. xanthi Pope in Schmidt, 1927; Mell, 1929
- Lampropholis mustelina Ehmann, 1988
- Leiolopisma otagense Smithells in Sharrell, 1966
- L. smithii J. Fawcett, pers. comm.
- L. zia Ehmann, 1988
- Mabuya capensis Rose, 1929, 1950; FitzSimons, 1943
- M. macrorhyncha Rebouças-Spieker and Vanzolini, 1978
- M. macularia Daniel, 1983
- Neoseps reynoldsi Telford, 1959
- Sphenomorphus quoyii Shine, 1988
- Scincus scincus Gesneri, 1551-1587; Topsell, 1608
- Tiliqua rugosa Mertens, 1960; Hitz, 1983
- Teiidae
- Tupinambis teguixin Reese, 1922; Krieg, 1925; Fitzgerald et al., 1989, In press
- T. rufescens Fitzgerald et al., 1989, In press
- Varanidae
- Varanus gouldii Berney, 1936
- V. griseus Auffenberg, 1981
- V. komodoensis Lallemand, 1929; Pfeffer, 1959; Auffenberg, 1981
- V. mitchelli Gow in Shine, 1988
- V. salvator Biswas and Kar, 1981
- V. varius Cogger, 1967, and in Shine, 1988; Mertens, 1987; Carter, 1989, and pers. comm.
- Xantusiidae
- Xantusia henshawi Shaw, 1949
- X. vigilis Cowles, 1944; Miller, 1954
- Trogonophidae or Amphisbaenidae (?)
- 'Amphisbaina' (= species?) Gesneri, 1551-1587; Topsell, 1608; Aldrovandi, 1640
- Sphenodontidae
- Sphenodon punctatus Thompson, 1988, and in Shine, 1988 and Heaton-Jones, 1989; Guillette et al., in review

\*Prior to 1932, Eumeces inexpectatus and E. laticeps were included within the species E. fasciatus (Cope, 1900; Davis, 1968; Steiner, 1986). The fasciatus group was divided into three species by Taylor (1932a, b, 1935); E. fasciatus, E. inexpectatus and E. laticeps. Consequently, some references pertaining to E. fasciatus prior to 1932 may actually refer to either, E. inexpectatus or E. laticeps.

\*\*Identity of species as described in this account is uncertain. This could refer to Eumeces gilberti; if so, it is the only record of brooding in this species.

TABLE VI

## Bibliographic sources for reports of snake parental behavior

| Taxa                            | Sources   |
|---------------------------------|---|
| Boidae                          |   |
| <u>Aspidites melanocephalus</u> | Ross, 1978; Boos, 1979; Murphy et al., 1981; Barker, 1982, 1985; Charles et al., 1985   |
| <u>Boa constrictor</u>          | Lanworn, 1972; Wells, 1981  |
| <u>Casarea dussumieri</u>       | Bloxam, 1984  |
| <u>Chondropython viridis</u>    | Kibler in Lederer, 1944; Pope, 1961; Kratzer, 1962; MacKay, 1973; Switak, 1975; Walsh, 1977, [1980]; Ross 1978; Olexa, 1979; Christian, 1981; Engelmann and Obst, 1981; Slavens, 1982, 1983, 1984, 1985, 1987; Hudson, 1983; Van Mierop et al., 1983; Zulich, 1983, 1985; [Orlov], 1986; Wexo, 1987 |
| <u>Epicrates cenchria</u>       | Boos, 1976; Brunner, 1979; Groves, 1981; Walsh and Davis, 1984  |
| <u>E. striatus</u>              | Huff, 1980; Slavens, 1987   |
| <u>E. subflavus</u>             | Gosse, 1851   |
| <u>Eunectes murinus</u>         | Neill and Allen, 1962; Holmstrom and Behler, 1981   |
| <u>E. notaeus</u>               | Holmstrom, 1981; Slavens, 1985, 1988; Townson, 1985   |
| <u>Liasis albertisii</u>        | Kinghorn, 1956; Johnson, 1975; Ross and Larman, 1977; Ross, 1978; Tarbet, 1984; Slavens, 1986; B. Clark, pers. comm.  |
| <u>L. boa</u>                   | Barker, 1982; Mehrtens, 1987  |
| <u>L. childreni</u>             | Ross, 1973, 1980a, 1983; Dunn, 1979a; Sheargold, 1979; Barker, 1982; Slavens, 1988  |
| <u>L. fuscus</u>                | Kinghorn, 1956; Gow, 1976; Ross and Larman, 1977; Ross, 1978, 1980b; Boos, 1979, 1983; McPhee, 1979; Barker, 1982; Weidner in Funk, 1982; Orlov, 1982; Bulian and Broer, 1984; Charles et al., 1985; [Orlov], 1986; Mehrtens, 1987; Shine, 1988   |
| <u>L. olivaceus</u>             | Kinghorn, 1956  |
| <u>L. papuanus</u>              | Tryon, 1985; Tryon and Whitehead, 1988  |
| <u>L. perthensis</u>            | Stafford, 1986  |
| <u>Morelia amethystina</u>      | Pope, 1961; Ross, 1978; Boos, 1979; McPhee, 1979; Parker, [1982]; Banks and Swaner, 1984; Charles et al., 1985; Grow, 1987; B. Clark, pers. comm.   |

M. bredli  
M. spilota

Gow, 1981, 1983  
Allan in G. Krefft, 1869; P. Krefft, 1926;  
Fleay, 1956; Cogger and Holmes, 1960;  
Gow, 1976, 1983; Ross, 1978; Worrel in  
Boos, 1979; McPhee, 1979; Harlow in  
Shine, 1980; Weidner in Funk, 1982;  
Broer, 1983; Burton and Burton, 1984;  
Harlow and Grigg, 1984; Charles et al.,  
1985; Bels and Van den Sande, 1986; Slip,  
1986; Lombard, 1987; Slip and Shine,  
1988a, b; Anonymous, [19??]

Python anchietae  
P. curtus

McLain, 1983; Laszlo, 1984; Branch, 1988  
Noble, 1935; Stemmler, 1969; Vinegar et  
al., 1970; Reitingner and Lee, 1978;  
Katuska, 1983; Stafford, 1986; Trutnau,  
1986; T. Weidner, pers. comm.

P. molurus

Bennett, 1824; Lamarrepiquot [=Lamarre-  
Piquot], 1835a; Valenciennes, 1841a, b;  
Duméril, 1842; Lamarre-Piquot, 1835b,  
1842, 1858a, b; Forbes, 1881; Holland in  
Hopley, 1882; Marshall, 1893; Pinkert,  
1893; Kern, 1907; Wall, 1912, 1921;  
Abercromby, 1913; Doflein, 1914; Krogh,  
1916; Leigh, 1926, 1936; Lederer, 1928,  
1944, 1956; Schlott, 1935; Kopstein,  
1938; Patsch, 1943; Smith, 1943; Walker  
and Stoddart in Angel, 1950; Deraniyagala  
1955; Stemmler-Morath, 1956; Vogel,  
[1958?]; Anonymous, 1960a, Dowling, 1960;  
Lutz, 1962; Deoras, 1965; Wendt, 1965;  
Hutchison et al., 1966; Yadav, 1967;  
Vinegar et al., 1970; Wagner, 1973, 1976;  
Coborn, 1975, 1985; Foekema, 1975;  
Acharjyo and Misra, 1976; Van Mierop and  
Barnard, 1976a, b, 1978; Frank, 1977;  
Acharjyo, 1978; Getreuer, 1979; Townson,  
1980, [1989]; Frye, 1981; Griehl, 1982;  
Clark and Tytle, 1983; Gurung, 1983; Van  
Mierop et al., 1983; [Gerholdt], 1984a,  
Conners, 1985; Slavens, 1985, 1987, 1988;  
Alderton, 1986; Michaels, 1986; Trutnau,  
1986; Whitaker and Whitaker, 1986; Clark,  
1988; Obst et al., 1988; Schleich and  
Kästle, 1988; Cox, 1989

P. regius

Pitman, 1938; Schivre, 1972; Logan, 1973;  
Peters, 1976; Boos, 1979; Van Mierop and  
Bessette, 1981; Malone, 1982; Orlov,  
1982; Lehmann and Lehmann, 1983; Laszlo,  
1984; Barten, 1986; [Orlov], 1986;  
Trutnau, 1986; Ellis and Chappell, 1987;  
Slavens, 1987; Kirschner and Ochsenbein,  
1988

P. reticulatus

Abbott in Wray, 1862; Köhler, 1907;  
Ditmars, 1910; Hilzheimer, 1910;

- P. sebae  
 Benedict, 1932; Kopstein, 1938; Lederer, 1944, 1956; Taylor, 1965; Honegger, 1970, 1970/71, 1975; Müller, 1970; Vinegar et al., 1970; Foekema, 1971; La Panouse and Pellier, 1973; Johnson, 1977; Trutnau, 1980, 1986; Slavens, 1984  
 Günther, 1862; Sclater, 1862; [Günther], 1886; F. FitzSimons, [1912], 1930; Werner, 1930 (in Angel, 1950); Benedict and Mann in Ditmars, 1931; Benedict, 1932; Benedict et al., 1932; Pitman, 1938; Lederer, 1942, 1944, 1956; Broadley, 1959; Anonymous, 1960<sub>a,b</sub>; Dowling, 1960, 1961; Sweeney, 1961; V. FitzSimons, 1962, 1970; Meyer-Holzappel, 1969; Schütte, 1970; Vinegar et al., 1970; Munnig Schmidt, 1971, 1973; Patterson, 1974; Branch and Patterson, 1975; Broadley and Cock, 1975<sub>a, b</sub>; Pienaur et al., 1978; Dunn, 1979<sub>b</sub>; Slavens, 1985; Trutnau, 1986; Patterson and Bannister, 1987; Branch, 1988; Schleich and Kästle, 1988; Shine, 1988  
 Murphy et al., 1978; Barker, 1982
- P. timoriensis  
 'Lilith' or 'arrow snake'  
 (= Eryx jaculus?)  
 Isaiah 34:15, [c. 745-350 BC], (McDowell et al., 1982)

## Colubridae

- Ahaetulla nasuta  
Amphiesma stolata  
Atretium schistosum  
Cemophora coccinea  
Cerberus rynchops  
Clelia clelia  
Coronella austriaca  
Diadophis punctatus
- Rieppel, 1970  
 Wall, 1911, 1921; Mell, 1929; Daniel, 1983  
 Murthy, 1986  
 Ditmars, 1907  
 Whitaker, 1978; Trutnau, 1986  
 Brazil, 1914, and in Roosevelt, 1914  
 Appleby, 1971  
 ?McCauley, 1945; Cook, 1954; Fowlie, 1965;  
 ?Brodie et al., 1969; Somma, pers. observ.
- Elaphe climacophora  
E. flavolineata  
E. guttata
- Fukada, 1965  
 Kopstein, 1938  
 Kelly et al., 1936; Haast and Anderson, 1981; Kent in Shine, 1988; T. Miller, pers. comm.
- E. obsoleta
- Ditmars, 1907; Medsger, 1919, 1932;  
 ?Netting, 1927; ?McCauley, 1945; Pope, 1946; M. Fisher, pers. comm.; J. Lombard, pers. comm.
- E. quadrivirgata  
E. quatuorlineata  
E. schrenki  
Farancia abacura
- Fukada, 1965; Orlow, 1982  
 Vogel, [1958?]  
 Kudryavtsev and Frolov, 1984  
 Ridgeway, 1883 (in Hay, 1892; Wright and Wright, 1957; Minton, 1972); Meade, 1937, 1940, 1945, 1946; Conant and Downs, 1940;

- Goldstein, 1941; Cagle, 1942; Reynolds and Solberg, 1942; Cook, 1954; Riemer, 1957; Tinkle, 1959; Hahn and Wilson, 1966; Crawford, 1984; Mehrtens, 1987; Dundee and Rossman, 1989
- Farancia erythrogramma Fry in Wright and Wright, 1957; Neill, 1964a; Ashton and Ashton, 1981
- Heterodon platirhinos Hay, 1892, 1893; Hahn 1909
- Hydrodynastes gigas Vogel 1964
- Lampropeltis triangulum Ditmars, 1907; Noble, 1920; Babcock, 1929; Anonymous, 1940; Minton, 1972; Minton and Minton, 1973; Marsec in Shine, 1988
- Lycodon aulicus Herklots, 1935
- L. striatus Wall, 1921
- Masticophis flagellum Meek, 1946
- Natrix natrix Stradling in Hopley, 1882; Gallwey, 1932; Berridge, 1935; Smith, 1951; Parker, 1963; Appleby, 1971
- Oligodon taeniolatus Daniel, 1983
- Opisthotropis latouchii Pope, 1929
- Pituophis catenifer Carl, 1944
- Psammophylax rhombeatus F. FitzSimons, [1912]; V. FitzSimons, 1962, 1970; Le Roux, 1964; Bourquin, 1970; Visser, 1971; De Waal, 1978; Branch, 1981, 1988; Broadley, 1983; Jacobsen, 1985; Trutnau, 1986; Patterson and Bannister, 1987
- P. tritaeniatus Sweeney, 1961; Isemonger, 1968; Branch, 1981; Hedges, 1983; Patterson and Bannister, 1987
- P. variabilis Spawls in Broadley, 1977
- Ptyas korros Mell, 1929
- P. mucosus Wall, 1907, 1921; Mell, 1929; Kopstein, 1938; Daniel, 1983
- Rhabdophis subminiata Mell, 1929
- Sinonatrix percarinata Pope, 1929, 1935
- Tropidonophis mairii ?Sundowner, 1895\*, 1902; Bredl in Shine, 1988
- Xenochrophis piscator Abercromby, 1913; Mell, 1929; Whitaker, 1978; Daniel, 1983; Whitaker and Whitaker, 1986
- Elapidae
- Bungarus caeruleus Wall, 1921; Daniel, 1983; Whitaker and Whitaker, 1986
- B. candidus Mell, 1929; Shaw and Shebbeare, 1931; Soderberg, 1973
- B. ceylonicus Green, 1905
- B. fasciatus Evans, 1905; Wall, 1921; Mell, 1929; Soderberg, 1973; Yahya, 1985
- 'Dipsas' (= Bungarus sp.?) Nicander of Colophon [135-133? BC]
- Calliophis maculiceps Frith, 1977 (also illustrated in Phelps, 1981)
- Demansia papuensis Parker, [1982]

- Micrurus fulvius  
M. psyches  
Naja melanoleuca  
N. naja
- 'Asp' (= Naja haje?)  
N. nigricollis  
Ophiophagus hannah
- 'Cockatrice' or 'Basilisk'  
 (= Ophiophagus hannah or  
Naja sp.?)  
Pseudechis butleri  
Pseudonaja textilis
- Hydrophiidae  
Pelamis platurus
- Laticaudidae  
Laticauda colubrina  
L. semifasciata
- Leptotyphlopidae  
Leptotyphlops dulcis  
L. humilis
- Typhlopidae  
Ramphotyphlops braminus  
Rhinotyphlops caecus
- Viperidae  
Agkistrodon contortrix  
A. piscivorus  
Calloselasma rhodostoma
- Campbell, 1973  
 Mole, 1924  
 Tryon, 1979; Dowling, 1986  
 [Appuhamy, 1810] (see Deraniyagala, 1955);  
 Fayrer, 1870; Kipling, 1894+; Wall, 1921;  
 Mell, 1929; Jennison, 1931; Kopstein,  
 1938; Smith, 1943; Simmon, 1944; Tweedie,  
 1954; Deraniyagala, 1955; Rao, 1957;  
 Duckett, 1964; Deoras, 1965; Petzold,  
 1968; Miller, 1970; Campbell and Quinn,  
 1975, Daniel, 1983; Whitaker and  
 Whitaker, 1986  
 Nicander of Colophon [135-133? BC]  
 Håkansson, 1981  
 Fayrer, 1870; Nicholson, 1870; Wasey, 1892;  
 Evans, 1903; Joynson, 1917; Wall, 1924;  
 Berridge, 1935; Mustill, 1936; Smith,  
 1936; Oliver, 1956; Leakey, 1969; Ionides  
 and Leakey in Soderberg, 1973;  
 Burchfield, 1977; Whitaker, 1977, 1978;  
 Reitingger and Lee, 1978; Daniel, 1983;  
 Gurung, 1983; Whitaker and Whitaker,  
 1986; Dattatri, 1987; Mehrtens, 1987;  
 Shine, 1988  
 Gesneri, 1551-1587; Topsell, 1608  
 Fitzgerald and Mengden, 1987  
 Fleay, 1943; Edwards and Wells in Shine,  
 1988; Shine, 1989
- Bertin and Burton, 1967
- Semper, 1881; ?Sundowner, 1895, 1902\*;  
 Smedley, 1931; Neill, 1964<sub>b</sub>, Taylor, 1965  
 Herre and Rabor, 1949
- Hibbard, 1964  
 Whitfield, 1983++
- Mell, 1929  
 Bogert, 1940
- Anderson, 1942; Fitch, 1960; Kennedy, 1964  
 Wharton, 1960, 1966  
 Smith, 1915, 1943; Tweedie, 1954; Leakey,  
 1969; Campden-Main, 1970; Reitingger and  
 Lee, 1978; Liat, 1982; York and  
 Burghardt, 1988; Gloyd and Conant, 1989

|                               |  |
|-------------------------------|--|
| <u>Causus rhombeatus</u>      | F. FitzSimons, [1912]; Woodward, 1933;<br>Sweeney, 1961; Broadley, 1983  |
| <u>Crotalus</u> sp.           | Audubon, 1909  |
| <u>C. adamanteus</u>          | Meek, 1946   |
| <u>C. atrox</u>               | Price, 1988  |
| <u>C. horridus</u>            | Anderson, 1942, 1965; Lokke, 1985; Martin,<br>1986a,b, 1989, and pers. comm.;<br>?Bartlett, 1987; ?Brown, 1987; ?Reinert<br>and Zappalorti, 1988   |
| <u>C. viridis</u>             | Gloyd, 1937; Jackley and Shelton in<br>Klauber, 1972; Duvall et al., 1985;<br>Graves, 1988, 1989   |
| <u>Deinagkistrodon acutus</u> | Fleck, 1987  |
| <u>Lachesis muta</u>          | Mole, 1924, and in Ditmars, 1910;<br>Donisthorpe, 1947; Ramsey and Travis,<br>1960; Wehekind, 1960; Emsley, 1977;<br>Caycedo, 1978; Frieberg, 1982 |
| <u>Porthidium nummifer</u>    | Picado T., 1931  |
| <u>Sistrurus catenatus</u>    | Greene and Oliver, 1965; ?Vogt, 1981;<br>Reinert and Kodrich, 1982   |
| <u>Trimeresurus kaulbacki</u> | Obst et al., 1988  |
| <u>T. monticola</u>           | Leigh, 1910; Pope, 1929, 1935  |
| <u>T. okinavensis</u>         | ?Fukada, 1964; Koba et al., 1970   |
| <u>T. wiroti</u>              | Mehrtens, 1987   |
| <u>Vipera aspis</u>           | Lanworn, 1972; Naulleau, 1987; Dowling,<br>1986  |
| <u>V. berus</u>               | Brittain, 1866 (in Hopley, 1882); Service,<br>1902; Smith, 1951; Appleby, 1971; Street,<br>1979; Naulleau, 1987                                    |

\*It is not known what species Sundowner actually observed but T. mairii and L. colubrina seem likely candidates. The credibility of Sundowner's [= Tichborne, H. (sic?)] (1895, 1902) observations are at best questionable, and mostly fabricated (Johnson and Smith, 1985). Nevertheless, it is likely that he also observed Australian pythons (species unspecified) brooding their eggs (Sundowner, 1895, 1902).

+Although a work of fiction, Kipling (1894) was one of the earliest published accounts of brooding in Naja naja (mistakenly referred to as Ophiophagus hannah) in English. He based his story, "Rikki-tikki-tavi," on a personal communication from an anonymous herpetologist (Kipling, 1894).

++Whitfield's (1983) mention of brooding in L. humilis is likely a mistaken reference to Hibbard's (1964) observations on L. dulcis.

## REFERENCES

- Abercromby, A. F. 1913. Some notes on the breeding habits of some Ceylon snakes and reptiles. *Spolia Zeylanica* 9: 144-147.
- Acharjyo, L. N. 1978. An incubating python. *Hornbill* 1978 (October-December): 36-37.
- Acharjyo, L. N., and R. Misra. 1976. Aspects of reproduction and growth of the Indian python, *Python molurus molurus*, in captivity. *Brit. J. Herp.* 5:562-565.
- Alderton, D. 1986. *A Petkeeper's Guide to Reptiles & Amphibians*. Salamander Books; London.
- Aldrovandi, U. 1640. *Serpentum et Draconū Historiae*. Nicolai Tebaldini; Bononiae.
- Allard, H. A. 1909. Notes on some salamanders and lizards of north Georgia. *Science* 30: 122-124.
- Alvarez del Toro, M. 1960. *Los Reptiles de Chiapas*. Instituto Zoológico del Estado; Tuxtla Gutiérrez, México.
- Anderson, P. 1942. Amphibians and reptiles of Jackson County, Missouri. *Bull. Chicago Acad. Sci.* 6: 203-220.
- Anderson, P. 1965. *The Reptiles of Missouri*. University of Missouri; Columbia.
- Angel, F. 1950. *Vie et Moeurs des Serpents*. Payot; Paris.
- Anonymous. 1940. Snakes. Pp. 338-389. In R. De Sola (ed), *American Wild Life, Illustrated*. Wise & Company; New York.
- Anonymous. 1941. Ball python upsets zoological ideas. *Sci. News Lett.* 39: 265.
- Anonymous. 1960a. ["This snake is not wired for sound"]. *Anim. King.* 63: [113] [photo].
- Anonymous. 1960b. [Excellent success has been achieved...]. *Anim. King.* 63: 146 [photo].
- Anonymous. [19??]. *Bush Dwellers of Australia*. Australian News and Information Bureau; [Canberra].
- Appleby, L.G. 1971. *British Snakes*. John Baker (Publishers); London.
- Appleton, A.B. (1986). Captive management and breeding of the Bolson tortoise (*Gopherus flavomarginatus*). [9]: 101-119. In S. McKeown, F. Caporaso, and K. H. Peterson (eds), *9th International Herpetological Symposium on Captive Propagation and Husbandry*. Zoological Consortium; Thurmont, Maryland.
- [Appuhamy, A.V.L.L. 1810. *The Sarapaduteya Nototh Sarapothpaththiya, a Palmadulla Snake Bite Treatise*.] [unpublished Sinhala serpent lore extracted from ancient palm leaf manuscripts (see Deraniyagala, 1955)].
- Aristotle. [d. 322 BC]. [History of Animals]. Translated in Cresswell, R. (1907). *Aristotle's History of Animals*. George Bell & Sons; London.
- Ashton, R.E., Jr., and P. S. Ashton. 1981. *Handbook of Reptiles and Amphibians of Florida. Part One. The Snakes*. Windward Publishing; Miami.
- Ashton, R.E., Jr., and P. S. Ashton. 1985. *Handbook of Reptiles and Amphibians of Florida. Part Two. Lizards, Turtles & crocodilians*. Windward Publishing; Miami.
- Asimov, I. 1968. *Asimov's Guide to the Bible. Vol. One. The Old Testament*. Doubleday & Company; Garden City.
- Audubon, J. W. 1909. *Audubon's Western Journal: 1849-1850*. Arthur H. Clark Company; Cleveland.



- Auffenberg, W. 1981. The Behavioral Ecology of the Komodo Monitor. University Presses of Florida; Gainesville.
- Babcock, H.L. 1929. The Snakes of New England. Boston Society of Natural History; Boston.
- Banks, C., and T. D. Schwaner. 1984. Two cases of interspecific hybridization among captive Australian boid snakes. *Zoo Biol.* 3: 221-227.
- Barbour, T., and C. T. Ramsden. 1919. The herpetology of Cuba. *Mem. Mus. Comp. Zool. Harvard* 47: 69-213.
- Barker, D.G. 1982. Maintenance and reproduction of pythons at the Dallas Zoo. [5]: 86-92. In R. A. Hahn (ed), 5th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Barker, D. G. 1985. Maintenance and reproduction of black-headed pythons at the Dallas Zoo. [8]:106-108. In S. McKeown and F. Caporaso (eds), 8th Annual International Herpetological Symposium on Captive Propagation & Husbandry. Zoological Consortium; Thurmont, Maryland.
- Barrett, S. L., and J. A. Humphrey. 1986. Agonistic interactions between Gopherus agassizii (Testudinidae) and Heloderma suspectum (Helodermatidae). *Southwest. Nat.* 31: 261-263.
- Barten, S.L. 1986. Egg retention in captive snakes. *Bull. Chicago Herpetol. Soc.* 21: 65-71.
- Bartlett, R.D. 1985. Notes on the natural history and reproductive strategy of the island glass lizard, Ophisaurus compressus. *Brit. Herpetol. Soc. Bull.* (11): 19-21.
- Bartlett, R.D. 1987. *In Search of Reptiles and Amphibians*. E.J. Brill; Leiden/New York/K benhavn/Köln.
- Baylis, J.R. 1981. The evolution of parental care in fishes, with reference to Darwin's rule of male sexual selection. *Envir. Biol. Fishes* 6: 223-251.
- Behler, J.L., and F.W. King 1979. The Audubon Society Field Guide to North American Reptiles and Amphibians. Alfred A. Knopf; New York.
- Bellairs, A.d'A. 1986. Reptiles. Pp. 60-67. In T.R. Halliday and K. Adler (eds), The Encyclopedia of Reptiles and Amphibians. Facts on File Publications; New York.
- Bels, V.L., and P.A. Van den Sande. 1986. Breeding the Australian carpet python Morelia spilotes variegata at Antwerp Zoo. *Int. Zoo Yearb.* 24/25: 231-238.
- Benedict, F.G. 1932. The physiology of large reptiles with special reference to the heat production of snakes, tortoises, lizards and alligators. *Carnegie Inst. Washington Publ.* (425): 1-539.
- Benedict, F.G., E.L. Fox, and V. Coropatchinsky. 1932. The incubating python: a temperature study. *Proc. Natl. Acad. Sci.* 18: 209-212.
- Bennet, E.T. 1829. *The Tower Menagerie*. Robert Jennings; London.
- Berney, F.L. 1936. Gould's monitor (Varanus gouldiae). *Queensland Nat.* 1936: 12-14.
- Berridge, W.S. 1935. *All about Reptiles and Batrachians*. George G. Harrap & Company; London.
- Bertin, L., and M. Burton. 1967. Reptiles (Class Reptilia). Pp. 283-330. In C.R. Murphy (ed), *The Larousse Encyclopedia of Animal Life*. Paul Hamlyn; London.
- Bishop, S.C. 1926. Records of some amphibians and reptiles from Kentucky. *Copeia* (152): 118-120.

- Biswas, S., and S. Kar. 1981. Some observations on nesting habits and biology of Varanus salvator (Laurenti) of Bhitarkanika Sanctuary, Orissa. *J. Bombay Nat. Hist. Soc.* 78: 303-308.
- Blair, A.P. 1961. Notes on Ophisaurus attenuatus attenuatus (Anguidae). *Southwest. Nat.* 6: 201.
- Blair, D.[W.] 1983a. Dragons of Cayman. *Oceans* 16(2):31-33.
- Blair, D.W. 1983b. The Cayman Island rock iguana Cyclura nubila caymanensis. *San Diego Herpetol. Soc. Newsl.* 5(9): 1-2.
- Blanchard, F.N. 1922. The amphibians and reptiles of western Tennessee. *Occ. Pap. Mus. Zool. Univ. Michigan* (117): 1-18.
- Blanchard, F.N. 1933. Eggs and young of the smooth green snake, Liopeltis vernalis (Harlan). *Pap. Mich. Acad. Sci. Arts Lett.* 17: 493-508.
- Bloxam, Q. 1984. A preliminary report on the captive management and reproduction of the Round Island boa (Casarea dussumieri). [7]:115-117. In P.J. Tolson (ed), 7th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Blumer, L.S. 1979. Male parental care in the bony fishes. *Quart. Rev. Biol.* 54: 149-161.
- Blumer, L.S. 1982. A bibliography and categorization of bony fishes exhibiting parental care. *Zool. J. Linn. Soc.* 75: 1-22.
- Bogert, C.M. 1940. Herpetological results of the Vernay Angola expedition with notes on African reptiles in other collections. Part I. Snakes, including an arrangement of the African Colubridae. *Bull. Amer. Mus. Nat. Hist.* 77: 1-107.
- Boos, H.[E.A.] 1976. Behavioral information needed. *AAZPA Newsl.* 17: 13.
- Boos, H.E.A. 1979. Some breeding records of Australian pythons Pythonidae. *Int. Zoo Yearb.* 19: 87-89.
- Boos, H.E.A. 1983. Australo-Asian pythons with some breeding records in captivity. *ASRA J.* 2(1): 22-32.
- Boulenger, G.A. 1903. Report on the batrachians and reptiles. Fasc. Malayensis, *Zool., Anthro. Zool. Res. Exp. Perak Siam Malay States*, 1901-1902 (1): 131-176.
- Bourquin, O. 1970. A note on the growth of spotted skaapesteker embryos (Psammophylax rhombeatus). *J. Herpetol. Assoc. Africa* (6): 8-9.
- Boylan, T. 1984. Breeding the rhinoceros iguana Cyclura c. cornuta at Sydney Zoo. *Int. Zoo Yearb.* 23: 144-148.
- Branch, B. [= W.R.]. 1988. Field Guide to the Snakes and Other Reptiles of Southern Africa. Ralph Curtis-Books-Publishing; Sanibel Island, Florida.
- Branch, W.R. 1981. Venomous snakes of southern Africa. 3. Concluding part: Colubridae. *Bull. Maryland Herpetol. Soc.* 17: 125-150.
- Branch, W.R., and R.W. Patterson. 1975. Notes on the development of embryos of the African rock python, Python sebae (Serpentes: Boidae). *J. Herp.* 9: 243-248.
- Brazil, V. 1914. La Défense Contre l'Ophidisme. Pocaï-Weiss & C.; Saint-Paul.
- Breckenridge, W.J. 1941. Amphibians and reptiles of Minnesota with special reference to the black-banded skink, Eumeces septentrionalis (Baird). Unpublished Ph.D. Dissertation; University of Minnesota, [Minneapolis].
- Breckenridge, W.J. 1943. The life history of the black-banded skink Eumeces septentrionalis septentrionalis (Baird). *Amer. Midl. Nat.* 29: 591-606.
- Breckenridge, W.J. 1944. Reptiles and Amphibians of Minnesota. University of Minnesota Press; Minneapolis.
- Bredin, E.J. 1981 (pers. comm.). Distribution of the northern prairie skink in Manitoba. Unpublished MS Report No. 81-7, Manitoba Department of

- Natural Resources, Wildlife Branch, Biological Services. [cited with permission of author.]
- Bredin, E.J. 1989 (pers. comm.). Status report on the northern prairie skink, Eumeces septentrionalis, in Canada. [Unpublished MS Report], The Amphibian and Reptile Subcommittee, Committee on the Status of Endangered Wildlife in Canada. [cited with permission of author.]
- Broadley, D.G. 1959. The herpetology of southern Rhodesia. Part 1. Snakes. Bull. Mus. Comp. Zool. Harvard 120: 1-100.
- Broadley, D.G. 1977. A revision of the African snakes of the genus Psammophylax Fitzinger (Colubridae). Occ. Pap. Natl. Mus. Monum. Rhodesia 1976B 6: 1-44.
- Broadley, D.G. 1983. FitzSimons' Snakes of Southern Africa. Delta Books; Johannesburg.
- Broadley, D. G., and E. V. Cock. 1975a. Snakes of Rhodesia. Longman Rhodesia (Bundu Series); Salisbury.
- Broadley, D. G., and E. V. Cock. 1975b. Snakes of Zimbabwe. [1982 reprinted, retitled edition]. Longman Zimbabwe (Bundu Series); Salisbury.
- Brodie, E.D., Jr., R.A. Nussbaum, and R.M. Storm. 1969. An egg-laying aggregation of five species of Oregon reptiles. Herpetologica 25: 223-227.
- Broer, W. 1983. Erfolgreiche Haltung und Nachzucht des Teppich- oder Rautenpythons, Python spilotus variegatus (Gray, 1844) (Serpentes: Boidae). Salamandra 19: 84-93.
- Brown, W.S. 1987. Hidden life of the timber rattler. Natl. Geogr. 172: 128-138.
- Browne, T. 1646. Pseudodoxia Epidemica: or, Enquiries into the Very Many Received Tenets, and Commonly Presumed Truths. Edward Dod; London.
- Brunner, J.C. 1979. Captive breeding of Columbian rainbow boas Epicrates cenchris crassus. [2]:39-47. In R.A. Hahn (ed), 2nd Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Bulian, J., and W. Broer. 1984. Ein seltener Python und seine Nachzucht: Liasis mackloti (Serpentes: Boidae). Salamandra 20: 205-211.
- Burchfield, P.M. 1977. Breeding the king cobra Ophiophagus hannah at Brownsville Zoo. Int. Zoo Yearb. 17: 136-140.
- Burghardt, G.M. 1973. Chemical release of prey attack: extension to naive newly hatched lizards, Eumeces fasciatus. Copeia 1973: 178-181.
- Burroughs, J. 1908. Leaf and Tendril. Houghton Mifflin Company; Boston and New York.
- Burt, C.E. 1928. The lizards of Kansas. Trans. Acad. Sci. St. Louis 26: 1-84.
- Burt, C.E. 1937. The lizards of the southeastern United States. Trans. Kansas Acad. Sci. 40: 349-366.
- Burt, C.E., and M.D. Burt. 1935. Further records of the ecology and distribution of amphibians and reptiles in the Middle West. Amer. Midl. Nat. 16: 311-336.
- Burt, C.E., and W.L. Hoyle. 1934. Additional records of the reptiles of the central prairie region of the United States. Trans. Kansas Acad. Sci. 37: 193-216.
- Burton, M., and R. Burton. 1984. Encyclopedia of Reptiles, Amphibians & Other Cold-blooded Animals. BPC Publishers; [London].
- Burton, R.F. 1885-1888. The Book of a Thousand Nights and a Night. The Burton Club; [London].

- Cagle, F.R. 1940. Eggs and natural nests of Eumeces fasciatus. Amer. Midl. Nat. 23: 227-233.
- Cagle, F.R. 1942. Herpetological fauna of Jackson and Union Counties, Illinois. Amer. Midl. Nat. 28: 164-200.
- Campbell, H., and R.S. Simmons. 1961. Notes on the eggs and young of Eumeces callicephalus Bocourt. Herpetologica 17: 212-213.
- Campbell, J.A. 1973. A captive hatching of Micrurus fulvius tenere (Serpentes, Elapidae). J. Herp. 7: 312-315.
- Campbell, J.A., and H.R. Quinn. 1975. Reproduction in a pair of Asiatic cobras, Naja naja (Serpentes, Elapidae). J. Herp. 9: 229-233.
- Campden-Main, S.M. 1970. A Field Guide to the Snakes of South Vietnam. United States National Museum; Washington, D.C.
- Carey, W.M. 1975. The rock iguana, Cyclura pinguis, on Anegada, British Virgin Islands, with notes on Cyclura ricordi and Cyclura cornuta on Hispaniola. Bull. Florida State Mus. Biol. Sci. 19: 189-233.
- Carl, G.C. 1944. The Reptiles of British Columbia. British Columbia Provincial Museum; Victoria.
- Carpenter, C.C. 1966. The marine iguana of the Galápagos Islands, its behavior and ecology. Proc. California Acad. Sci. Fourth Ser. 34: 329-376.
- Carpenter, C.C., and G.W. Ferguson. 1977. Variation and evolution of stereotyped behavior in reptiles. 7: 335-508. In C. Gans and D.W. Tinkle (eds), Biology of the Reptilia. Ecology and Behavior A. Academic Press; New York.
- Carr, A. 1952. Handbook of Turtles. The Turtles of the United States, Canada, and Baja California. Comstock Publishing Associates; Ithaca and London.
- Carter, D.B. 1989. Evidence of parental involvement in the release of hatchling Varanus varius from termite mounds. First World Congr. Herp. [Unpubl.] Abst. (Canterbury, United Kingdom) 1: [58]
- Carver, J. 1778. Travels Through the Interior Parts of North-America, in the Years 1766, 1767, and 1768. J. Walter and S. Crowder; London.
- Castillo S., C. 1989. Behavior and social organization in Sauromalus varius, Dickerson (Reptilia, Sauria, Iguanidae) in captivity. First World Congr. Herp. [Unpubl.] Abst. (Canterbury, United Kingdom) 1: [261].
- Caycedo, G.C. 1978. Perdido en el Amazonas. Carlos Valencia Editores; Bogotá.
- Charles, N., R. Field, and R. Shine. 1985. Notes on the reproductive biology of Australian pythons, genera Aspidites, Liasis and Morelia. Herpetol. Rev. 16: 45-48.
- Chateaubriand, F.R. de 1827. Voyage en Amérique. Lacroix; Paris.
- Christian, K.A., and C.R. Tracy. 1982. Reproductive behavior of Galápagos land iguanas, Conolophus pallidus, on Isla Santa Fe, Galápagos. Pp. 366-379. In G.M. Burghardt and A.S. Rand (eds), Iguanas of the World. Their Behavior, Ecology, and Conservation. Noyes Publications; Park Ridge, New Jersey.
- Christian, T. 1981. A field trip to the McIlwraith Range - Cape York Peninsula. Monitor-Bull. Victorian Herp. Soc. 1(2): 40-56.
- Claffey, O., and B. Johnson. 1982a. Captive reproduction of the sheltopusik (Ophisaurus apodus). AAZPA Reg. Conf. Proc. 1982: 103-106.
- Claffey, O., and B. Johnson. 1982b. Captive reproduction of the sheltopusik (Ophisaurus apodus). Animal Keepers Forum 9: 329-332.
- Clark, B. 1988. Breeding the albino Burmese python. Vivarium 1: 18-21.

- Clark, B., and T. Tytle. 1983. Python molurus bivittatus (Burmese python). Morphology. Herpetol. Rev. 14: 121.
- Clausen, R.T. 1938. Notes on Eumeces anthracinus in central New York. Copeia 1938: 3-7.
- Coborn, J. 1975. Report on the reproduction of captive Indian pythons (Python molurus bivittatus). Brit. J. Herp. 5: 471-472.
- Coborn, J. 1985. Howell Beginner's Guide to Snakes. Howell Book House; New York.
- Cochran, P.A. 1989. Confirmation of the five-lined skink (Eumeces fasciatus) in northeastern Wisconsin. Bull. Chicago Herpetol. Soc. 24: 30-31.
- Cogger, H.G. 1967. Australian Reptiles in Colour. East-West Center Press; Honolulu.
- Cogger, H.G. 1974. Voyage of the banded iguana. Australian Nat. Hist. 18: 144-149.
- Cogger, H.G., and A. Holmes. 1960. Thermoregulatory behaviour in a specimen of Morelia spilotes variegata Gray (Serpentes: Boidae). Proc. Linn. Soc. New South Wales 85: 328-333.
- Collins, H.H., Jr. 1959. Complete Field Guide to American Wildlife. East, Central and North. Harper and Row, Publishers; New York.
- Collins, J.T. 1975. Observations on reproduction in the southern coal skink (Eumeces anthracinus pluvialis Cope). Trans. Kansas Acad. Sci. 77: 126-127.
- Conant, R. 1938. The reptiles of Ohio. Amer. Midl. Nat. 20: 1-200.
- Conant, R. 1951. The Reptiles of Ohio. University of Notre Dame Press; Notre Dame, Indiana.
- Conant, R., and A. Downs, Jr. 1940. Miscellaneous notes on the eggs and young of reptiles. Zoologica 25: 33-48.
- Congdon, J.D. 1989. Proximate and evolutionary constraints on energy relations of reptiles. Physiol. Zool. 62: 356-373.
- Connors, S. 1985. Snake breeding projects at the Cheyenne Mountain Zoo. Colorado Herpetol. Soc. [Newsl.] 13(5): 1-2.
- Cook, F.A. 1943. Alligators and lizards of Mississippi. Surv. Bull. Mississippi Game Fish Comm. Jackson, Mississippi (1): 1-20.
- Cook, F.A. 1954. Snakes of Mississippi. Surv. Bull. State Game Fish Comm. Jackson, Mississippi [2]: 1-40.
- Coombs, W.P., Jr. 1982. Juvenile specimens of the ornithischian dinosaur Psittacosaurus. Paleontology (London) 25: 89-108.
- Cooper, J.E., R.L. Franz, F. Groves, J.D. Hardy, Jr., H.S. Harris, Jr., D.S. Lee, P. Wemple, and R.G. Tuck. 1973. Endangered amphibians and reptiles of Maryland. Bull. Maryland Herpetol. Soc. 9: 42-99.
- Cooper, W.E., Jr., and L.J. Vitt. 1985. Blue tails and autotomy: enhancement of predation avoidance in juvenile skinks. Z. Tierpsychol. 70: 265-276.
- Cooper, W.E., Jr., L.J. Vitt, L.D. Vangilder, and J.W. Gibbons. 1983. Natural nests and brooding behavior of Eumeces fasciatus. Herpetol. Rev. 14: 65-66.
- Cope, E.D. 1900. The crocodylians, lizards, and snakes of North America. Rept. U.S. Natl. Mus. 1898: 153-1294.
- Corrington, J.D. 1929. Herpetology of the Columbia, South Carolina region. Copeia (172): 58-83.
- Cowles, R.B. 1944. Parturition in the yucca night lizard. Copeia 1944: 98-100.
- Cox, M.J. 1989. Notes on the natural history (breeding) of Naja kaouthia and Python molurus bivittatus in Thailand. Bull. Chicago Herpetol. Soc. 24: 127-128.

- Crawford, T. 1984. Herping notes from the deep south. *Iowa Herpetol. Soc. Newsl.* 8(1): 2-3.
- Crutchfield, T.E. 1982. Courtship and nesting behavior in Cyclura nubila lewisi. [5]: 54-56. In R.A. Hahn (ed), 5th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Crutchfield, T.E. 1986. The first captive propagation of the Grand Cayman Island iguana Cyclura n. lewisi. [9]:165-170. In S. McKeown, F. Caporaso and K.H. Peterson (eds), 9th International Herpetological Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Daniel, J.C. 1983. *The Book of Indian Reptiles*. Bombay Natural History Society; Bombay.
- Dattatri, S. 1987. Breeding the king cobra (Ophiophagus hannah) in captivity. *J. Bombay Nat. Hist. Soc.* 84: 222-227.
- Davis, D.M. 1968. A study of variation in the North American lizards of the fasciatus group of the genus Eumeces (Scincidae). Unpublished Ph.D. Dissertation; Duke University, [Durham].
- Dellinger, T. 1989. Energetic costs of reproduction in a polygynous reptile: the marine iguana. *First World Congr. Herp.* [Unpubl.] Abst. (Canterbury, United Kingdom) 1: [83]. [Information not provided in abstract.]
- Deoras, P.J. 1965. *Snakes of India*. Secretary, National Book Trust; New Delhi.
- Deraniyagala, P.E.P. 1955. A Colored Atlas of Some Vertebrates from Ceylon. Volume Three. Serpentine Reptilia. Ceylon National Museums; Colombo.
- De Waal, S.W.P. 1978. The Squamata (Reptilia) of the Orange Free State, South Africa. *Mem. Nas. Mus. (Bloemfontein)* 11: 1-160.
- Dewsbury, D.A. 1985. Paternal behavior in rodents. *Amer. Zool.* 25: 841-852.
- Digney, T., and T. Tytle. 1983. Captive maintenance and propagation of the lizard genus Phelsuma. [6]:141-156. In D. L. Marcellini (ed), 6th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Ditmars, R.L. 1904. Observations on lacertilians. *Eighth Ann. Rept. New York Zool. Soc.* [8]: 146-160.
- Ditmars, R.L. 1907. *The Reptile Book*. Doubleday, Page & Company; Garden City.
- Ditmars, R.L. 1910. *Reptiles of the World*. Macmillan Company; New York.
- Ditmars, R.L. 1931. *Snakes of the World*. Macmillan Company; New York.
- Ditmars, R.L., and W. Bridges. 1937. *Wild Animal World. Behind the Scenes at the Zoo*. D. Appleton-Century Company; New York/London.
- Doflein, F. 1914. Das Tier als Glied des Naturganzen. II Band: 1-960. In R. Helle and F. Doflein (eds.), *Tierbau und Tierleben*. B. G. Teubner; Leipzig und Berlin.
- Donisthorpe, H. 1947. Lizards and snakes. Pp. 418-439. In Anonymous (ed), *Wild Life the World Over, Illustrated*. Wise & Company; New York.
- Dowling, H.G. 1950. A new southeastern record for the coal skink. *Copeia* 1950: 235.
- Dowling, H.G. 1960. Thermistors, air conditioners & insecticides. *Anim. King.* 63: 202-207.
- Dowling, H.G. 1961. How old are they and how big do they grow? *Anim. King.* 64: 171-175.

- Dowling, H.G. 1986. Snakes. Pp. 112-127. In T.R. Halliday and K. Adler (eds), The Encyclopedia of Reptiles and Amphibians. Facts on File Publications; New York.
- Druce, G.C. 1910. The amphisbaena and its connexions in ecclesiastical art and architecture. Archeol. J. 67(268); Ser. 2 17(4): 285-317.
- Duckett, J.E. 1964. The cobras of Batu Kawan, Province Wellesley. Malayan Nat. J. 18: 210-211.
- Duellman, W.E., and A. Schwartz. 1958. Amphibians and reptiles of southern Florida. Bull. Florida State Mus. Biol. Sci. 3: 181-324.
- Duellman, W.E., and L. Trueb. 1986. Biology of Amphibians. McGraw-Hill Book Company; New York.
- Duméril, [A.-M.-C.] 1842. Sur le développement de la chaleur dans les oeufs des serpents, et sur l'influence attribuée à l'incubation de la mère. Compt. Rend. Hebd. Séan. Acad. Sci. Paris 14: 193-203.
- Dundee, H.A., and D.A. Rossman. 1989. The Amphibians and Reptiles of Louisiana. Louisiana State University Press; Baton Rouge.
- Dunn, E. R. 1920. Some reptiles and amphibians from Virginia, North Carolina, Tennessee and Alabama. Proc. Biol. Soc. Washington 33: 129-138.
- Dunn, R. W. 1979a. Breeding Children's pythons Liasis childreni at Melbourne Zoo. Int. Zoo Yearb. 19: 89-90.
- Dunn, R. W. 1979b. Breeding African pythons Python sebae at Melbourne Zoo. Int. Zoo Yearb. 19: 91-92.
- Duvall, D. 1986. A new question of pheromones: aspects of possible chemical signaling and reception in the mammal-like reptiles. Pp. 219-238. In N. Hotton, III, P.D. MacLean, J.J. Roth, and E.C. Roth (eds.), The Ecology and Biology of Mammal-like Reptiles. Smithsonian Institution Press; Washington and London.
- Duvall, D., M.B. King, and K.J. Gutzwiller. 1985. Behavioral ecology and ethology of the prairie rattlesnake. Natl. Geogr. Res. 1: 80-111.
- Duvall, D., J. Trupiano, and H.M. Smith. 1979. An observation of maternal behavior in the Mexican desert spiny lizard, Sceloporus rufidorsum. Trans. Kansas Acad. Sci. 82: 60-62.
- Ehmann, H. 1988. Post-oviposition egg care in three Australian skink lizards. Herpetofauna (Sydney) 18: 26-29.
- Eibl-Eibesfeldt, I. 1966. Das Verteidigen der Eiablageplätze bei der Hood-Meerechse (Amblyrhynchus cristatus venustissimus). Z. Tierpsychol. 23: 627-631.
- Eijsden, E. H. T. van. 1978. Over de gecko's Gehyra mutilata en Hemiphyllodactylus t. typus, immigranten op Mauritius. Lacerta 36: 72-79.
- Ellis, T. L., and M.A. Chappell. 1987. Metabolism, temperature relations, maternal behavior, and reproductive energetics in the ball python (Python regius). J. Comp. Physiol. B 157: 393-402.
- Ellison, T.L. 1985. Iguana iguana in captivity. [8]: 93-95. In S. McKeown and F. Caporaso (eds), 8th Annual International Herpetological Symposium on Captive Propagation & Husbandry. Zoological Consortium; Thurmont, Maryland.
- Emsley, M. 1977. Snakes of Trinidad and Tobago. Bull. Maryland Herpetol. Soc. 13: 201-304.
- Engelmann, W.-E., and F.J. Obst. 1981. Mit gespaltener Zunge. Aus der Biologie und Kulturgeschichte der Schlangen. Herder Feiburg; Basel, Wien.
- Ernst, C.H., and R.W. Barbour. 1989. Turtles of the World. Smithsonian Institution Press; Washington, D.C. and London.



- Evans, G.H. 1903. The king-cobra, or hamadryad-*Naia bungarus* (Boulenger), *Ophiophagus elaps* (Günther). J. Bombay Nat. Hist. Soc. 14: 409-418.
- Evans, G.H. 1905. Breeding of the banded krait (*Bungarus fasciatus*) in Burma. J. Bombay Nat. Hist. Soc. 16: 519-520.
- Evans, H.E., and R. M. Roecker. 1951. Notes on the herpetology of Ontario, Canada. Herpetologica 7: 69-71.
- Evans, L.T. 1959. A motion picture study of maternal behavior of the lizard, *Eumeces obsoletus* Baird and Girard. Copeia 1959: 103-110.
- Fayrer, T. 1870. The Thanatophidia of India, Being a Description of the Venomous Snakes of the Indian Peninsula with an Account of the Influence of Their Poison on Life and a Series of Experiments. J. and A. Churchill; London.
- Fitch, A.V. 1964. Temperature tolerances of embryonic *Eumeces*. Herpetologica 20: 184-187.
- Fitch, H.S. 1954. Life history and ecology of the five-lined skink, *Eumeces fasciatus*. Univ. Kansas Publ. Mus. Nat. Hist. 8: 1-156.
- Fitch, H.S. 1955. Habits and adaptations of the great plains skink (*Eumeces obsoletus*). Ecol. Monogr. 25: 59-83.
- Fitch, H.S. 1956. Temperature responses of free-living amphibians and reptiles of northeastern Kansas. Univ. Kansas Publ. Mus. Nat. Hist. 8: 419-475.
- Fitch, H.S. 1960. Autecology of the copperhead. Univ. Kansas Publ. Mus. Nat. Hist. 13: 85-288.
- Fitch, H.S. 1967. Ecological studies of lizards on the University of Kansas Natural History Reservation. Pp. 30-34. In W.W. Milstead (ed), Lizard Ecology: a Symposium. University of Missouri Press; Columbia.
- Fitch, H.S. 1970. Reproductive cycles in lizards and snakes. Univ. Kansas Misc. Publ. Mus. Nat. Hist. (52): 1-247.
- Fitch, H.S. 1982. Reproductive cycles in tropical reptiles. Occ. Pap. Mus. Nat. Hist. Univ. Kansas (96): 1-53.
- Fitch, H.S. 1986. Ecology of the slender glass lizard. Soc. Stud. Amphib. Rept.-Herp. Leag. Meet. [Unpubl. Abst.] (Springfield, Missouri) 1986: 79.
- Fitch, H.S. 1989. A field study of the slender glass lizard, *Ophisaurus attenuatus*, in northeastern Kansas. Occ. Pap. Mus. Nat. Hist. Univ. Kansas (125): 1-50.
- Fitch, H.S., and P.L. von Achen. 1977. Spatial relationships and seasonality in the skinks *Eumeces fasciatus* and *Scincella laterale* in northeastern Kansas. Herpetologica 33: 303-313.
- Fitzgerald, L.A., J.M. Chani, and O.E. Donadio. 1989. *Tupinambis* lizards in Argentina: implementing management of a traditionally exploited resource. First World Congr. Herp. [Unpubl.] Abst. (Canterbury, United Kingdom) 1: [101].
- Fitzgerald, L.A., J.M. Chani, and O.E. Donadio. In press. *Tupinambis* lizards in Argentina: implementing management of a traditional exploitation system. In J.G. Robinson and K.H. Redford (eds), Neotropical Wildlife Use and Conservation. University of Chicago Press; Chicago.
- Fitzgerald, M., and G.A. Mengden. 1987. Captive breeding and oviparity in *Pseudechis butleri* (Serpentes: Elapidae). Amphibia-Reptilia 8: 165-170.
- FitzSimons, F.W. [1912]. The Snakes of South Africa. Their Venom and the Treatment of Snake Bite. T. Maskew Miller; Cape Town.
- FitzSimons, F.W. 1930. Pythons and Their Ways. George G. Harrap & Company; London.



- FitzSimons, V.F.[M.] 1943. The lizards of South Africa. Transvaal Mus. Mem. (1): 1-528.
- FitzSimons, V.F.M. 1962. Snakes of Southern Africa. Purnell & Sons; Cape Town.
- FitzSimons, V.F.M. 1970. A Field Guide to the Snakes of Southern Africa. Collins; London.
- Fleay, D. 1943. The brown snake - Dangerous fellow. Victorian Nat. 59: 147-152.
- Fleay, D. 1956. Talking of Animals. Jacaranda Press; Brisbane.
- Fleck, J. 1987. Erst-Nachzucht von Agkistrodon acutus (Günther, 1888) (Serpentes: Crotalidae). Salamandra 23: 193-203.
- Foekema, G.M.M. 1971. Broeden pythons moeilijk in gevangenschap? Lacerta 29: 107-110.
- Foekema, G.M.M. 1975. Ontwikkeling en voortplanting van Python molurus bivittatus in een huiskamerterrarium. Lacerta 33: 123-139.
- Forbes, W.A. 1881. Observations on the incubation of the Indian python (Python molurus), with special regard to the alleged increase of temperature during the process. Proc. Zool. Soc. London 1881: 960-967.
- Fouquette, M.J., Jr., and H.L. Lindsay, Jr. 1955. An ecological survey of reptiles in parts of northwestern Texas. Texas J. Sci. 7: 402-421.
- Fowlie, J.A. 1965. The Snakes of Arizona. Their Derivation, Speciation, Distribution, Description, and Habits - A Study in Evolutionary Herpeto-zoogeographic Phylogenetic Ecology. McGraw Hill; [New York].
- Frank, W. 1979. Boas and Other Non-venomous Snakes. [English Language Edition]. T.F.H. Publications; Neptune, New Jersey.
- Freiberg, M. 1982. Snakes of South America. T.F.H. Publications; Neptune, New Jersey.
- Frith, C.B. 1977. A survey of the snakes of Phuket Island and the adjacent mainland areas of peninsular Thailand. Nat. Hist. Bull. Siam Soc. 26: 263-316.
- Frye, F.L. 1981. Biomedical and Surgical Aspects of Captive Reptile Husbandry. Veterinary Medicine Publishing Company; Edwardsville, Kansas.
- Fukada, H. 1964. Biological studies on snakes XI. Eggs and young of Trimeresurus okinavensis Boulenger. Bull. Kyoto Gakugei Univ. Ser. B (24): 7-11.
- Fukada, H. 1965. Breeding habits of some Japanese reptiles (Critical review). Bull. Kyoto Gakugei Univ. Ser. B (27): 65-82.
- Funk, R.S. 1982. Intra-brood variation in boas and pythons. [5]:17-29. In R.A. Hahn (ed), 5th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Gallwey, E. 1932. Eggs of grass snake. Naturalist (London) 910: 320.
- Gehlbach, F.R. 1965. Herpetology of the Zuni Mountains region, northwestern New Mexico. Proc. U.S. Natl. Mus. 116: 243-332.
- [Gerholdt, J.] 1984<sub>a</sub>. Herp hints. Minnesota Herpetol. Soc. Newsl. 4(6): 6.
- [Gerholdt, J.] 1984<sub>b</sub>. Phantastic phenological phinds. Minnesota Herpetol. Soc. Newsl. 4(7): 10-12.
- Gesneri [= Gessner], C. 1551-1587. Historiae Animalium. Zürich.
- Getreuer, W.K. 1979. Voortplanting en ontwikkeling in het terrarium, van Python molurus bivittatus. Lacerta 37: 71-82.
- Gibbons, J.R.H. 1984/85. On the trail of the crested iguana. Anim. King. 87(6): 40-45.
- Gibbons, J.R.H., and I.F. Watkins. 1982. Behavior, ecology, and conservation of South Pacific banded iguanas, Brachylophus, including a newly

- discovered species. Pp. 418-441. In G.M. Burghardt and A.S. Rand (eds), *Iguanas of the World. Their Behavior, Ecology, and Conservation*. Noyes Publications; Park Ridge, New Jersey.
- Gilbert, B. 1987. Lizards that take to the desert like ducks to water. *Smithsonian* 18(5): 78-87.
- Gittleman, J.L. 1981. The phylogeny of parental care in fishes. *Anim. Behav.* 29: 936-941.
- Gloyd, H.K. 1937. A herpetological consideration of faunal areas in southern Arizona. *Bull. Chicago Acad. Sci.* 5: 79-136.
- Gloyd, H.K., and R. Conant. 1989. Snakes of the *Agkistrodon* Complex: A Monographic Review. Society for the Study of Amphibians and Reptiles; Athens, Ohio.
- Goldstein, R.C. 1941. Notes on the mud snake in Florida. *Copeia* 1941: 49-50.
- Gosse, P.H. 1851. *A Naturalist's Sojourn in Jamaica*. Longman, Brown, Green and Longmans; London.
- Gow, G.F. 1976. *Snakes of Australia*. Angus & Robertson Publishers; Sydney.
- Gow, G.F. 1981. A new species of *Python* from central Australia. *Australian J. Herp.* 1: 29-34.
- Gow, G.F. 1983. *Snakes of Australia*. Revised Edition. Angus & Robertson Publishers; Sydney.
- Graves, B.M. 1988. Functions and causes of selected aspects of birthing rookery use by prairie rattlesnakes (*Crotalus viridis viridis*) in Wyoming. Unpublished Ph.D. dissertation, University of Wyoming, [Laramie].
- Graves, B.M. 1989. Defensive behavior of female prairie rattlesnakes (*Crotalus viridis*) changes after parturition. *Copeia* 1989: 791-794.
- Graves, B.M., and D. Duvall. 1983. A role for aggregation pheromones in the evolution of mammallike reptile lactation. *Amer. Nat.* 122: 835-839.
- Green, E.E. 1905. On the nesting of the snake, *Bungarus ceylonicus*. *Spolia Zeylanica* 3: 158.
- Green, N.B., and T.K. Pauley. 1987. *Amphibians and Reptiles in West Virginia*. University of Pittsburgh Press; Pittsburgh.
- Greene, H.W., and B.E. Dial. 1966. Brooding behavior by female alligator lizards. *Herpetologica* 22: 303.
- Greene, H.W., and G.V. Oliver, Jr. 1965. Notes on the natural history of the western massasauga. *Herpetologica* 21: 225-228.
- Greer, A.E. 1970. Evolutionary and systematic significance of crocodilian nesting habits. *Nature (London)* 227: 523-524.
- Greer, A.E. 1971. Crocodilian nesting habits and evolution. *Fauna* 2: 20-28.
- Griehl, K. 1984. *Snakes. Giant Snakes and Non-venomous Snakes in the Terrarium*. English Language Edition. Barron's Educational Series; Woodbury, New York.
- Gross, M.R., and R.C. Sargent. 1985. The evolution of male and female parental care in fishes. *Amer. Zool.* 25: 807-822.
- Groves, J.D. 1981. Observations and comments on the post-parturient behaviour of some tropical boas of the genus *Epicrates*. *Brit. J. Herp.* 6: 89-91.
- Groves, J.D. 1982. Egg-eating behavior of brooding five-lined skinks, *Eumeces fasciatus*. *Copeia* 1982: 969-971.
- Grow, D. 1987. Amethystine pythons hatch at the Oklahoma City Zoo. *AAZPA Newsl.* 28(6): 16.
- Guillette, L.J., Jr., and N. Hotton, III. 1986. The evolution of mammalian reproductive characteristics in therapsid reptiles. Pp. 239-250. In N.

- Hotton, III, P.D. MacLean, J.J. Roth, and E.C. Roth (eds), *The Ecology and Biology of Mammal-like Reptiles*. Smithsonian Institution Press; Washington and London.
- Guillette, L.J., Jr., A. Cree, and T.S. Gross. In review. Plasma prostaglandin and steroid concentrations during natural nest digging, oviposition and nest guarding in the tuatara, Sphenodon punctatus. *Biol. Reprod.*
- Günther, A.[C.L.G.] 1862. [Remarks on the incubation of the female python in the Society's gardens.] *Proc. Zool. Soc. London* 1862: 1-2.
- [Günther, A.C.L.G.] 1886. Python. 20:144. In T.S. Baynes and W.R. Smith (eds), *Encyclopaedia Britannica*. Ninth Edition. Adam and Charles Black; Edinburgh.
- Gurung, K.K. 1983. *Heart of the Jungle*. André Deutsch with Tiger Tops PVT; London and Kathmandu, Nepal.
- Haast, W.E., and R. Anderson. 1981. *Complete Guide to Snakes of Florida*. Phoenix Publishing Company; Miami.
- Hahn, D.E., and L.D. Wilson. 1966. Variation in two broods of Farancia abacura reinwardti Schlegel (Serpentes: Colubridae) from Louisiana. *J. Ohio Herpetol. Soc.* 5: 159-160.
- Hahn, W.L. 1909. Notes on the mammals and cold-blooded vertebrates of the Indiana University Farm, Mitchell, Indiana. *Proc. U.S. Nat. Mus.* 35: 545-581.
- Håkansson, N.T. 1981. An annotated checklist of reptiles known to occur in the Gambia. *J. Herp.* 15: 155-161.
- Hall, R.J. 1972. Ecology of a population of the great plains skink (Eumeces obsoletus). *Univ. Kansas Sci. Bull.* 49: 357-388.
- Hall, R.J., and H.S. Fitch. 1972. Further observations on the demography of the great plains skink (Eumeces obsoletus). *Trans. Kansas Acad. Sci.* 74: 93-98.
- Hamilton, W.J., Jr. 1958. Technophagy in the Florida five-lined skink. *Herpetologica* 14: 28.
- Hamilton, W.J., Jr., and J.A. Pollack. 1958. Notes on the life history of the red-tailed skink. *Herpetologica* 14: 25-28.
- Hammond, S. 1985. Reproductive behavior in the broadhead skink, Eumeces laticeps. *Notes NOAH* 12(4): 7-8.
- Harlow, P., and G. Grigg. 1984. Shivering thermogenesis in a brooding diamond python, Python spilotes spilotes. *Copeia* 1984: 959-965.
- Hasegawa, M. 1984. Biennial reproduction in the lizard Eumeces okadae on Miyake-Jima, Japan. *Herpetologica* 40: 194-199.
- Hasegawa, M. 1985. Effect of brooding on egg mortality in the lizard Eumeces okadae on Miyake-Jima, Izu Islands, Japan. *Copeia* 1985: 497-500.
- Hay, O. P. 1892. The batrachians and reptiles of the state of Indiana. *Indiana Dept. Geol. Nat. Res. Seventeenth Ann. Rept.* 1891 17: 409-609.
- Hay, O.P. 1893. On the breeding habits, eggs, and young of certain snakes. *Proc. U.S. Natl. Mus.* 1892 15: 385-397.
- Heaton-Jones, T.G. 1989. March meeting proceedings. *Gainesville Herpetol. Soc. Newsl.* 5(8): 32-34.
- Hedges, N.G. 1983. *Reptiles and Amphibians of East Africa*. Kenya Literature Bureau; Nairobi.
- Hediger, H. 1937. *Selttsame Reptilien und Amphibien der Solomon-Inseln*. *Natur Volk* 67: 590-595.
- Hediger, H. 1968. Der Akademiestreit um die brütende Python. *Neue Zürcher Zeitung* 1968(351): 5.

- Hediger, H. 1986. Strange reptiles and amphibians of the Solomon Islands. Bull. Chicago Acad. Sci. 21: 49-50. [English translation of Hediger, 1937.]
- Heller, E. 1903. Papers from the Hopkins Stanford Galapagos Expedition, 1898-1899. XIV. Reptiles. Proc. Washington Acad. Sci. 5: 39-98.
- Herklots, G.A.C. 1935. Land snakes of Hong Kong. Part III. Hong Kong Nat. 6: 192-207.
- Herre, A.W.C.T., and D.S. Rabor. 1949. Notes on Philippine sea snakes of the genus Laticauda. Copeia 1949: 282-284.
- Hibbard, C.W. 1964. A brooding colony of the blind snake, Leptotyphlops dulcis dissecta Cope. Copeia 1964: 222.
- High, J. [1976]. Natural History of the Seychelles. Government Printer; Port Victoria.
- [Hikida, T. 1975. An observation of the brooding habit in Eumeces okadae. Niigata Herpetol. J.] [in Japanese] 1(3): 6-7.
- Hikida, T. 1981. Reproduction in the Japanese skink (Eumeces laticutatus) in Kyoto. Zool. Mag. 90: 85-95.
- Hilzheimer, M. 1910. Die Brutflege der Reptilien. Aus Natur. (Leipzig) 6: 361-367.
- Hitz, R. 1983. Pflege und Nachzucht von Trachydosaurus rugosus Gray, 1827 im Terrarium (Sauria: Scincidae). Salamandra 19: 198-210.
- Hodsdon, L.A., and J.F.W. Pearson. 1943. Notes on the discovery and biology of two Bahamen fresh-water turtles of the genus Pseudemys. Proc. Florida Acad. Sci. 6(2): 17-23.
- Holmes, I. 1823. An Account of the United States of America. Caxton Press, Henry Fisher; London.
- Holmstrom, W.F., Jr. 1981. Observations on the reproduction of the yellow anaconda Eunectes notaeus at the New York Zoological Park. Int. Zoo Yearb. 21: 92-94.
- Holmstrom, W.F., Jr., and J.L. Behler. 1981. Post-parturient behavior of the common anaconda, Eunectes murinus. Zool. Gart. N.F. (Jena) 51: 353-356.
- Honegger, R.E. 1969. Notes on some amphibians and reptiles at Zurich Zoo. Int. Zoo Yearb. 9: 24-28.
- Honegger, R.E. 1970. Beitrag zur Fortpflanzungsbiologie von Boa constrictor und Python reticulatus (Reptilia, Boidae). Salamandra 6: 73-79.
- Honegger, R.E. 1970/71. Beitrag zur Fortpflanzungsbiologie einiger tropischer Reptilien. Freunde Kölner Zoo 12: 175-179.
- Honegger, R.E. 1975. Breeding and maintaining reptiles in captivity. Pp. 1-12. In R.D. Martin (ed), Breeding Endangered Species in Captivity. Academic Press; London.
- Honegger, R.E. 1985. Additional notes on the breeding and captive management of the prehensile-tailed skink (Corucia zebrata). Herpetol. Rev. 16: 21-23.
- Hopley, C.C. 1882. Snakes: Curiosities and Wonders of Serpent Life. Griffith and Farran; London.
- Horner, J.R. 1982. Evidence of colonial nesting and 'site fidelity' among ornithischian dinosaurs. Nature (London) 297: 675-676.
- Horner, J.R. 1984. The nesting behavior of dinosaurs. Sci. Amer. 250(4): 130-137.
- Horner, J.R. 1987. Ecologic and behavioral implications derived from a dinosaur nesting site. 2:50-63. In S.J. Czerkas and E.C. Olson (eds), Dinosaurs Past and Present. Natural History Museum of Los Angeles County/University of Washington Press; Seattle and London.

- Horner, J.R., and J. Gorman. 1989. *Digging Dinosaurs*. Workman Publishing; New York.
- Horner, J.R., and R. Makela. 1979. Nest of juveniles provides evidence of family structure among dinosaurs. *Nature* (London) 282: 296-298.
- Hoy, P.R. 1883. Catalogue of the cold-blooded vertebrates of Wisconsin. *Geol. Wisconsin Surv.* 1873-1878 1: 422-435.
- Hudson, R. 1983. The reptile reproduction program at the Fort Worth Zoo. [6]:328-349. In D.L. Marcellini (ed), 6th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Huff, T.A. 1980. Captive propagation of the subfamily Boinae with emphasis on the genus Epicrates. (1):125-134. In J.B. Murphy and J.T. Collins (eds), *Reproductive Biology and Diseases of Captive Reptiles*. SSAR Contr. Herp. Society for the Study of Amphibians and Reptiles; [Athens, Ohio].
- Huff, T.A. 1985. An overview of lizard husbandry and propagation with emphasis on the work at the Reptile Breeding Foundation. Pp. 51-76. In R.L. Gray (ed), *Proceedings of the Northern California Herpetological Society and Bay Area Amphibian and Reptile Society 1985 Conference on Captive Propagation and Husbandry of Reptiles and Amphibians*. Northern California Herpetological Society; Davis.
- Hunter, J.D. 1824. *Memoirs of a Captivity Among the Indians of North America, From Childhood to the Age of Nineteen: with Anecdotes Descriptive of Their Manners and Customs. To Which is Added, Some Account of the Soil, Climate, and Vegetable Productions of the Territory Westward of the Mississippi*. Longman, Hurst, Rees, Orme, Brown and Green; London.
- Hurter, J. 1911. Herpetology of Missouri. *Trans. Acad. Sci. St. Louis* 20:59-274.
- Hutchison, V.H., H.G. Dowling, and A. Vinegar. 1966. Thermoregulation in a brooding female Indian python, Python molurus bivittatus. *Science* 151: 694-696.
- Isemonger, R.M. 1968. *Snakes of Africa*. Books of Africa; Cape Town.
- Iverson, J.B. 1977. Behavior and ecology of the rock iguana Cyclura carinata. Unpublished Ph.D. Dissertation; University of Florida, [Gainesville].
- Iverson, J.B. 1979. Behavior and ecology of the rock iguana Cyclura carinata. *Bull. Florida State Mus. Biol. Sci.* 24: 175-358.
- Jacobsen, N.H.G. 1985. *Uit die Natuur ons Reptiele*. CUM-Boeke; Roodepoort, South Africa.
- Jayaram, K.C. 1974. Ecology and distribution of fresh-water fishes, Amphibia and reptiles. 23:517-584. In M.S. Mani (ed), *Ecology and Biogeography of India*. Monogr. Biol. Dr. W. Junk b.v., Publishers; The Hague.
- Jennison, G. 1931. Cobras bred at Belle Vue Zoological Gardens, Manchester. *Proc. Zool. Soc. London* 1931: 1413.
- Jes, H. 1987. *Lizards in the Terrarium*. English Edition. Barron's Educational Series; New York.
- Johnson, B.J., and H.M. Smith. 1985. Sundowner, the forgotten ophidiophile. *Bull. Maryland Herpetol. Soc.* 21: 119-134.
- Johnson, C.R. 1975. Defensive display behaviour in some Australian and Papuan-New Guinean pygopodid lizards, boid, colubrid and elapid snakes. *Zool. J. Linn. Soc.* 56: 265-282.
- Johnson, T.R. 1977. Adult female reticulated python with eggs at St. Louis Zoo Reptile House on 12 February 1977. *Herpetol. Rev.* 8: 147 [photo].

- Johnson, T.R. 1979. Notes on an egg clutch and a new county record for the broadhead skink (Eumeces laticeps). St. Louis Herpetol. Soc. Bull. 7(1): 12.
- Johnson, T.R. 1987. The Amphibians and Reptiles of Missouri. Conservation Commission of the State of Missouri; Jefferson City.
- Jones, R.E., and L.J. Guillette, Jr. 1982. Hormonal control of oviposition and parturition in lizards. *Herpetologica* 38: 80-93.
- Joynson, H.W. 1917. A hamadryad's nest. *J. Nat. Hist. Soc. Siam* 2: 255.
- Katuska, C.J. 1983. Husbandry and management of the genus Python. St. Louis Herpetol. Soc. Newsl. 4(4): 8-18.
- Keenleyside, M.H.A. 1981. Parental care patterns of fishes. *Amer. Nat.* 117: 1019-1022.
- Kelly, H.A., A.W. Davis, and H.C. Robertson. 1936. Snakes of Maryland. Natural History Society of Maryland; Baltimore.
- Kendeigh, S.C. 1952. Parental care and its evolution in birds. *Illinois Biol. Monogr.* 22: 1-356.
- Kennedy, J.P. 1956. An arboreal nest of the five-lined skink, Eumeces fasciatus, in eastern Texas. *Southwest. Nat.* 1: 158-159.
- Kennedy, J.P. 1964. Natural history notes on some snakes of eastern Texas. *Texas J. Sci.* 16: 210-215.
- Kern, E. 1907. Riesenschlangen in der Gegangenschaft ausgebrütet. *D. Umschau, Wissensch. Mag.* 11: 1050-1052.
- Kinghorn, J.R. 1956. The Snakes of Australia. Angus & Robertson; Sydney.
- Kipling, R. 1894. The Second Jungle Book. Harper & Brothers; [London].
- Kirschner, A., and A. Ochsenbein. 1988. Beobachtungen zu Haltung und Nachzucht von Python regius (Shaw, 1802) (Serpentes: Boidae). *Salamandra* 24: 193-202.
- Klauber, L.M. 1972. Rattlesnakes. Their Habits, Life Histories, and Influence on Mankind. Second Edition. University of California Press; Berkeley and Los Angeles.
- Kleiman, D.G. 1985. Paternal care in New World primates. *Amer. Zool.* 25: 857-859.
- Kleiman, D.G., and J.R. Malcolm. 1981. The evolution of male parental investment in mammals. Pp. 347-388. In D.G. Gubernick and P.H. Klopfer (eds), *Parental Care in Mammals*. Plenum Press; New York.
- Klots, A.B. 1930. Notes on Amphibia and Lacertilia collected at Weymouth, N. J. *Copeia* (173): 109-111.
- Koba, K., K. Tanaka, K. Yoshizaki, and E. Nakamoto. 1970. Eggs and hatching of the hime-habu, Trimeresurus okinavensis Boulenger. *Snake* 2: 111-121.
- Koch, H. 1972. Vie en captivité et reproduction de Gekko gekko L. *Aquarama* 6: 44-45.
- Köhler, W. 1907. Riesenschlangen in Gefangenschaft ausgebrütet. *Blätt. Aquar.-Terrarienk.* (Stuttgart) 18: 498.
- Kopstein, F. 1938. Eien Beitrag zur Eierkunde und zur Fortpflanzung der Malaiischen Reptilien. *Bull. Raffles Mus.* (Singapore) (14): 81-167.
- Kratzer, H. 1962. Überraschende Nachzucht von Chondropython viridis. *Aquar.-Terrar.- Z.* (Stuttgart) 15: 117-119.
- Kreffft, G. 1869. The Snakes of Australia. An Illustrated and Descriptive Catalogue of All the Known Species. Thomas Richards, Government Printer; Sydney.
- Kreffft, P. 1926. Das Terrarium. 2 Auflage. Pfenningstorff; Berlin.
- Krieg, H. 1925. Biologische Reisestudien in Südamerika. IV. Beobachtungen über die 'Iguana' (Tupinambis teguixin L.). *Z. Wissensch. Biol. Abt. A.; Z. Morph. Ökol. Tiere* A3: 441-451.

- Krogh, A. 1916. *The Respiratory Exchange of Animals and Man*. Longmans, Green and Company; London.
- Kudryavtsev, S.V., and V.E. Frolov. 1984. On the reproductive biology of the Russian rat snake Elaphe s. schrencki (Squamata: Colubridae). [7]:134-138. In P.J. Tolson (ed), 7th Annual Reptile Symposium on Captive Propagation and husbandry. Zoological Consortium; Thurmont, Maryland.
- Lallemant, G.F.M.A. 1929. De Reuzen leguaan (Varanus komodoensis) op de eilanden Komodo en Rintja. *Trop. Natur* 18(8): 125-128.
- Lamarre-Picquot [= Lamarre-Picquot, P.]. 1835a. Ophiologie. L'Institut. Sect. I (Paris, Bureau 19[de l'Institut.]) 3: 70.
- Lamarre-Picquot, P. 1835b. Réponse pour servir de réfutation aux opinions et à la critique du rapport de M. Constant Duméril, etc. 64 pp. [Unpublished brochure].
- Lamarre-Picquot, [P.] 1842. Troisième mémoire sur l'incubation et autres phénomènes observés chez les ophidiens. *Compt. Rend. Hebd. Séan. Acad. Sci.* 14: 164.
- Lamarre-Picquot, [P.] 1858a. [l'Incubation des ophidiens]. *Compt. Rend. Hebd. Sean. Acad. Sci.* 47:458.
- Lamare-Picquot [=Lamarre-Picquot, P.]. 1858b. [Adresse une note supplémentaire...]. *Compt. Rend. Hebd. Séan. Acad. Sci.* 47: 525-526.
- Lang, J.W. [1982.] Distribution and abundance of the five-lined skink (Eumeces fasciatus) in Minnesota. Unpublished Final Report to the Nongame Wildlife Program, Minnesota Department of Natural Resources; St. Paul.
- Lang, J.W. 1983. Blue devils in Rock Valley: our search for the five-lined skink. *Minnesota Volunteer* 1983(July-August): 40-46.
- Lang, J.W. 1987. Crocodylian behaviour: implications for management. Pp. 273-293. In G.J.W. Webb and S.C. Manolis (eds), *Wildlife Management: Crocodiles and Alligators*. Surrey Beatty & Sons; Chipping Norton, Australia.
- Langerwerf, B. 1981. The southern alligator lizard, Gerrhonotus multicarinatus Blainville 1935: its care and breeding in captivity. *Brit. Herpetol. Soc. Bull.* (4): 21-25.
- Langerwerf, B. 1984. Techniques for large-scale breeding of lizards from temperate climates in greenhouse enclosures (breeding many species of lizards in captivity, aiming the maintenance of populations of each species outside their natural habitat). *Acta Zool. Pathol. Antverp.* (78): 163-176.
- Lanworn, R.A. 1972. *The Book of Reptiles*. Hamlyn Publishing Group; Feltham, England.
- La Panouse, R. de, and C. Pellier. 1973. Ponte d'un python réticulé (Python reticulatus) élevé en terrarium, et incubation des oeufs. *Bull. Mus. Natl. Hist. Nat. Ser. 3, Zool.* (Paris) 79(105): 37-48.
- Laszlo, J. 1984. Further notes on reproductive patterns of amphibians and reptiles in relation to captive breeding. *Int. Zoo Yearb.* 23: 166-174.
- Lawler, H.E., and J.L. Jarchow. 1986. A captive management plan for large iguanine lizards using the Isla San Esteban chuckwalla Sauromalus varius Dickerson as a model. [9]:137-164. In S. McKeown, F. Caporaso, and K.H. Peterson (eds), 9th International Herpetological Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Leakey, J.H.E. 1968. Observations made on king cobras in Thailand during May 1966. *J. Natl. Res. Counc. Thailand* 5: 1-10.
- Lederer, G. 1928. Zucht ereignis im Frankfurter Aquarium. *D. Umschau, Wissensch. Mag.* 32: 425-426.



- Lederer, G. 1944. Nahrungserwerb, Entwicklung, Paarung und Brutfürsorge von Python reticulatus (Schneider). Zool. Jahrb. Abt. f; Anat. Ont. Tiere (Jena) 68: 363-398.
- Lederer, G. 1956. Fortpflanzungsbiologie und Entwicklung von Python molurus molurus (Linné) und Python molurus bivittatus (Kühl). Aquar.- Terrar.- Z. (Stuttgart) 9: 243-248.
- Lehmann, C., and K.P. Lehmann. 1983. Haltung und Vermehrung des Königspythons, Python regius, in Terrarium. Aquar. Terr. (Berlin) 30:100-103.
- Leigh, C. 1910. An oviparous Indian viper. Field (London) 115(2975): 3.
- Leigh, C. 1926. Notes on snakes. J. Bombay Nat. Hist. Soc. 31: 227-228.
- Leigh, C. 1936. Breeding of pythons. Field (London) 1936: 1556.
- Le Roux, S.F. 1964. A note on the spotted skaapstekker (Psammophylax rhombeatus Linnaeus). African Wild Life 18: 113-115.
- Leviton, A.E. [1972]. Reptiles and Amphibians of North America. Doubleday and Company; New York.
- Liat, L.B. 1982. Poisonous Snakes of Peninsular Malaysia. Second Edition. Malayan Nature Society; Kuala Lumpur.
- Lillywhite, H.B. 1988. Biology of the Reptilia. Vol. 16. Ecology B. Defense and Life History. [Book review]. Copeia 1988: 1102-1104.
- Lockwood, S. 1883. Maternal anxiety in a horned toad. Amer. Nat. 17: 682-683.
- Logan, T. 1973. Observations on the ball python (Python regius) in captivity at Houston Zoological Gardens. J. Herpetol. Assoc. Africa (10): 5-8.
- Lokke, J. 1985. A question of parental care in the timber rattlesnake Crotalus horridus. Nebraska Herpetol. Newsl. 6(4): 4-5.
- Lombard, J. 1987. News. 2 May, 1987. Nebraska Herpetol. Newsl. 8(3): 1-2.
- Loop, M.S., and S.A. Scoville. 1972. Response of newborn Eumeces inexpectatus to prey-object extracts. Herpetologica 28: 254-256.
- Louwman, J.W.W. 1982. Breeding the six-footed tortoise Geochelone emys at Wassenaar Zoo. Int. Zoo Yearb. 22: 153-156.
- Lutz, R. 1962. Fortpflanzung des Tigerpython. Aquar.- Terrar.- Z. (Stuttgart) 15: 277-279.
- Lynn, W.G., and C. Grant. 1940. The herpetology of Jamaica. Bull. Inst. Jamaica Sci. Ser. (1): 1-148.
- Mackay, R.D. 1973. The green python. Wildlife Australia 10(4): 108.
- Malcolm, J.R. 1985. Paternal care in canids. Amer. Zool. 25: 853-856.
- Malone, D.R. 1982. Captive care and breeding of the ball python. Bull. Chicago Herpetol. Soc. 17: 96-98.
- Mansueti, R. 1948. "Scorpion" of the treetops. Nat. Hist. 57: 213-215+240.
- Marshall, W. 1893. Brütende Schlangen. Blätt. Aquar.- Terrarienk. (Stuttgart) 4: 158-161.
- Martin, W.H. 1986a. Reproduction of the timber rattlesnake in northwestern Virginia. Herp. Leag. - Soc. Stud. Amphib. Rept. Meet. [Unpubl. Abst.] (Springfield, Missouri) 1986: 94.
- Martin, W.H. 1986b. Life history of the timber rattlesnake, Crotalus horridus. Investigator's Annual Report, U.S. Fish and Wildlife Service, National Park Service, U.S. Department of the Interior. Pp. 1.
- Martin, W.H. 1989. The timber rattlesnake, Crotalus horridus in the Appalachian Mountains of eastern North America. First World Congr. Herp. [Unpubl.] Abst. (Canterbury, United Kingdom) 1: [193].
- Martof, B. 1956. A contribution to the biology of the skink, Eumeces laticeps. Herpetologica 12: 111-114.



- [Mathui, K. 1985. Amphibians and Reptiles of Japan. Shogakukan; Tokyo.] [in Japanese].
- McAllister, C.T. 1987. Eumeces septentrionalis obtusirostris (southern prairie skink). Reproduction. Herpetol. Rev. 18: 75.
- McCauley, R.H., Jr. 1939. Differences in the young of Eumeces fasciatus and Eumeces laticeps. Copeia 1939: 93-95.
- McCauley, R.H., Jr. 1945. The Reptiles of Maryland and the District of Columbia. Robert H. McCauley, Jr.; Hagerstown, Maryland.
- McClellan, W.H., R. Mansueti, and F. Groves. 1943. The lizards of central and southern Maryland. Proc. Nat. Hist. Soc. Maryland (8): 1-42.
- McDiarmid, R.W. 1978. Evolution of parental care in frogs. Pp. 127-147. In G.M. Burghardt and M. Bekoff (eds), The Development of Behavior: Comparative and Evolutionary Aspects. Garland Press; New York.
- McDowell, W. A., A. Farstad, W. White, and J.D. Price. (Eds.) 1982. The Holy Bible. The New King James Version. Thomas Nelson; Nashville.
- McKeown, S., J.O. Juvik, and D.E. Meier. 1982. Observations on the reproductive biology of the land tortoises Geochelone emys and Geochelone yniphora in the Honolulu Zoo. Zoo Biol. 1: 223-235.
- McKeown, S., and M.J. Miller. 1985. A brief note on the natural history, captive maintenance and propagation of the Seychelles giant skin-sloughing gecko Ailuronyx sechellensis. [8]:96-101. In S. McKeown and F. Caporaso (eds), 8th Annual International Herpetological Symposium on Captive Propagation & Husbandry. Zoological Consortium; Thurmont, Maryland.
- McLain, J.M. 1983. Notes on boid reproduction at the Houston Zoological Park. [6]:141-156. In D.L. Marcellini (ed), 6th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- McLain, R.B. 1899. Contributions to North American herpetology. Critical notes on a collection of reptiles, from the western coast of the United States. [Robert Baird McLain]; Wheeling, West Virginia. [also reprinted. 1971. Miscellaneous Publications. Facsimile Reprints in Herpetology. Society for the Study of Amphibians and Reptiles; Athens, Ohio.]
- McPhee, D.R. 1979. The Observer's Book of Snakes and Lizards of Australia. Frederick Warne (Publishers); London.
- Meade, G.P. 1937. Breeding habits of Farancia abacura in captivity. Copeia 1937: 12-15.
- Meade, G.P. 1940. Maternal care of eggs by Farancia. Herpetologica 2: 15-20.
- Meade, G.P. 1945. Further observations on Louisiana captive snakes. Copeia 1945: 73-75.
- Meade, G.P. 1946. The natural history of the mud snake. Sci. Month. 63: 21-29.
- Mease, J. 1807. A Geological Account of the United States Comprehending a Short Description of Their Animal, Vegetable and Mineral Productions, Antiquities and Curiosities. B. & T. Kite; Philadelphia.
- Medsger, O.P. 1919. Egg-laying habits of the pilot snake, Callopeltis obsoletus. Copeia (69): 28-29.
- Medsger, O.P. 1932. Nature Rambles. Summer. An Introduction to Country-lore. Frederick Warne & Company; New York and London.
- Meek, G. 1946. Creatures of Mystery. J. W. Burke Company [for] Gray Meek; Macon, Georgia.
- Mehaffey, D.T. 1986. Notes on captive propagation of the Solomon Island lizard Corucia zebrata at Fort Worth Zoo. Gainesville Herpetol. Soc. Newsl. 2(4): 12-15.

- Mehrtens, J.M. 1987. Living Snakes of the World in Color. Sterling Publishing Company; New York.
- Mell, R. 1929. Beiträge zur Fauna Sinica. IV. Grundzüge einer Ökologie der Chinesischen Reptilien und einer Herpetologischen Tiergeographie Chinas. Walter De Gruyter & Company; Berlin und Leipzig.
- Mendelssohn, H., and A. Bouskila. 1989. Comparative ecology of Uromastix aegyptius and Uromastix ornatus in southern Israel and in southern Sinai. First World Congr. Herp. [Unpubl.] Abst. (Canterbury, United Kingdom) 1: [198].
- Mertens, R. 1960. The World of Amphibians and Reptiles. English Translation. McGraw-Hill Book Company; New York.
- Mertens, T.J. 1987. Some preliminary notes and observations on the tree goanna (Varanus varius) of Bowman Park, South Australia. Unpublished Report to Bowman Park Trust. 16 pp.
- Meshaka, W.E., S.E. Trauth, and B.P. Butterfield. 1988. Natural nests, eggs, and hatchlings of the broad-headed skink, Eumeces laticeps, from northwestern Arkansas. Bull. Chicago Herpetol. Soc. 23: 104-105.
- Meyer-Holzappel, M. 1969. Notes on the breeding and egg-laying of some reptiles at Berne Zoo. Int. Zoo Yearb. 9: 20-23.
- Michaels, S.J. 1986. Husbandry and captive propagation of the light phase Indian python, Python molurus molurus, with a report of a predominantly male clutch of offspring. Bull. Chicago Herpetol. Soc. 21: 53-64.
- Miller, H. 1970. The cobra, India's "good snake." Natl. Geogr. 138: 392-409.
- Miller, M.J. 1980. The leopard gecko, Eublepharis macularis (Gray). Bull. Chicago Herpetol. Soc. 15: 10-15.
- Miller, M.J. 1982. Phelsumas. A case of monotypic care of a polytypic genus. [5]:103-118. In R.A. Hahn (ed), 5th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Miller, M.J. 1983a. Breeding the ground gecko Chondrodactylus angulifer angulifer (Peters, 1870) in captivity. [6]:171-175. In D.L. Marcellini (ed), 6th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Miller, M.J. 1983b. Breeding the Turkistan plate-tailed gecko Teratoscincus scincus (Schlegel, 1858) in captivity. [6]:176-180. In D.L. Marcellini (ed), 6th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Miller, M.[J.] 1984. Captive husbandry and propagation of geckos. Bull. Chicago Herpetol. Soc. 19: 41-54.
- Miller, M.R. 1954. Further observations on reproduction in the lizard Xantusia vigilis. Copeia 1954: 38-40.
- Minton, S.A., Jr. 1972. Amphibians and reptiles of Indiana. Indiana Acad. Sci. Monogr. (3): 1-346.
- Minton, S.A., Jr., and M.R. Minton. 1973. Giant Reptiles. Charles Scribner's Sons; New York.
- Moehn, L.D. 1980. Microhabitat preference in the broadhead skink. Bull. Chicago Herpetol. Soc. 15: 49-53.
- Molchos, D. 1971. Photograph of Eumeces inexpectatus with her nest in situ, in a "hammock" near Miami, Florida. Herpetol. Rev. 3: 64 [photo].
- Mole, R.R. 1924. The Trinidad Snakes. Proc. Zool. Soc. London 1924: 235-278.
- Morafka, D.J. 1981. The status and distribution of the Bolson tortoise (Gopherus flavomarginatus). (12): 71-94. In R.B. Bury (ed), North

- American Tortoises: Conservation and Ecology. Wildlife Research Report. United States Department of the Interior, Fish and Wildlife Service; Washington, D.C.
- Mount, R.H. 1961. The natural history of the red-tailed skink, Eumeces egregius Baird. Unpublished Ph.D. Dissertation; University of Florida, [Gainesville].
- Mount, R.H. 1963. The natural history of the red-tailed skink, Eumeces egregius Baird. Amer. Midl. Nat. 70: 356-385.
- Mount, R.H. 1975. The Reptiles and Amphibians of Alabama. Auburn University Agricultural Experiment Station; Auburn.
- Müller, M. 1970. Einige beobachtungen zur Fortpflanzungsbiologie bei Riesenschlange im Zoologischen Garten Leipzig. Aquar. Terrar. (Leipzig) 17: 162-164.
- Munnig Schmidt, C.H. 1971. Waarnemingen bij het broeden van Python sebae. Lacerta 29: 105-106.
- Munnig Schmidt, C.H. 1973. Verslag van een geslaagde kweek met de Africaanse python (Python sebae). Lacerta 31: 91-101.
- Murphy, J.B., D.G. Barker, and B.W. Tryon. 1978. Miscellaneous notes on the reproductive biology of reptiles. 2. Eleven species of the family Boidae, genera Candoia, Corallus, Epicrates and Python. J. Herp. 12: 385-390.
- Murphy, J.B., W.L. Lamoreaux, and D.G. Barker. 1981. Miscellaneous notes on the reproductive biology of reptiles. 4. Eight species of the family Boidae, genera Acrantophis, Aspidites, Candoia, Liasis and Python. Trans. Kansas Acad. Sci. 84: 39-49.
- Murthy, T.S.N. 1986. The Snake Book of India. R. P. Singh Gahlot for International Book Distributors; Dehra Dun.
- Mustill, F.J. 1936. A hamadryad's nest and eggs. J. Bombay Nat. Hist. Soc. 39: 186-187.
- Nauelleau, G. 1987. Les serpentes de France. 2<sup>e</sup> édition. Rev. Française Aquariol. 11(3-4): 1-56.
- Neill, W.T. 1948. The lizards of Georgia. Herpetologica 4: 153-158.
- Neill, W.T. 1964a. Taxonomy, natural history, and zoogeography of the rainbow snake, Farancia erythrogramma (Palisot de Beauvois). Amer. Midl. Nat. 71: 257-295.
- Neill, W.T. 1964b. Viviparity in snakes: some ecological and zoogeographical considerations. Amer. Midl. Nat. 98: 35-55.
- Neill, W.T., and R. Allen. 1962. Parturient anaconda, Eunectes gigas Latreille, eating own abortive eggs and foetal membranes. Quart. J. Florida Acad. Sci. 25: 73-75.
- Nelson, W.F. 1963. Natural history of the northern prairie skink, Eumeces septentrionalis septentrionalis (Baird). Unpublished Ph.D. Dissertation; University of Minnesota, [Minneapolis].
- Netting, M.G. 1927. A note on the egg-laying of Elaphe obsoleta (Say). Copeia (162): 29-30.
- Nicander of Colophon. [135-133? BC]. [Theriaca]. Translated in A.S.F. Gow and A.F. Scholfield. 1953. Nicander. The Poems and Poetical Fragments. The Cambridge University Press; London.
- Nicholson, E. 1870. Indian Snakes. Higginbotham & Co.; Madras.
- Niekisch, M. 1975. Pflege und Nachzucht von Egernia cunninghami (Sauria, Scincidae). Salamandra 11: 130-135.
- Niekisch, M. 1980. Terraristische Beobachtungen zur Biologie von Egernia cunninghami (Reptilia: Sauria: Scincidae). Salamandra 16: 162-176.

- Noble, G.K. 1920. An early record for the oviposition and hatching of the milk snake. *Copeia* (88): 98-100.
- Noble, G.K. 1921. Do snakes swallow their young for protection? *Copeia* (98): 54-57.
- Noble, G.K. 1935. The brooding habit of the blood python and of other snakes. *Copeia* 1935: 1-3.
- Noble, G.K., and E.R. Mason. 1932. The relation of water regulation to habitat selection of reptiles. *Science* 76: 545-546.
- Noble, G.K., and E.R. Mason. 1933. Experiments on the brooding habits of the lizards Eumeces and Ophisaurus. *Amer. Mus. Novit.* (619): 1-29.
- Nussbaum, R.A. 1985. The evolution of parental care in salamanders. *Misc. Publ. Mus. Zool. Michigan* (169): 1-50.
- Obst, F.J., K. Richter, and U. Jacob. 1988. *The Completely Illustrated Atlas of Reptiles and Amphibians for the Terrarium. English-language Edition.* T. F. H. Publications; Neptune City, New Jersey.
- Olexa, A. 1979. The green python, Chondropython viridis (Schlegel, 1872) and its reproduction in captivity. *Gazella* 2: 83-87.
- Oliver, J.A. 1956. Reproduction in the king cobra, Ophiophagus hannah Cantor. *Zoologica* 41: 145-152.
- [Orlov, N.L.] 1986. [Facultative endogenous thermoregulation in python snakes (Boidae, Pythoninae) and correlation between endothermic reactions and behavioural heat regulation in these snakes.] [in Russian Cyrillic] *Zool. Zhur.* 65: 551-559.
- Orlov [=Orlov], N.L. 1982. Die Fortpflanzung der Pythons P. regius und Liasis mackloti und der Kletternattern Elaphe climacophora und E. quadrivirgata im Terrarium. *Herpetofauna (Weinstadt)* (19): 25-30.
- Osadnik, G. 1984. An investigation of egg laying in Phelsuma (Reptilia: Sauria: Gekkonidae). *Amphibia-Reptilia* 5: 125-134.
- Parker, F. [1982]. The snakes of Western Province. *Wildlife in Papua New Guinea (Div. Wildlife, Konedobu)* (82/1): 1-78.
- Parker, H.W. 1963. *Snakes.* Robert Hale; London.
- Parmalee, P.W. 1955. Reptiles of Illinois. *Illinois State Mus. Pop. Sci. Ser.* 5: 1-88.
- Patsch, H.-J. 1943. Fortpflanzungsbiologische Beobachtungen an Python bivittatus. *Zool. Gart. N.F. (Leipzig)* 15: 132-133.
- Patterson, R.W. 1975. Hatching the African python Python sebae in captivity. *Int. Zoo Yearb.* 14: 81-82.
- Patterson, R.[W.], and A. Bannister. 1987. *Reptiles of Southern Africa.* C. Struik Publishers; Cape Town.
- Perrone, M., and T.M. Zaret. 1979. Parental care patterns of fishes. *Amer. Nat.* 113: 351-361.
- Peters, J.A. 1964. *A Dictionary of Herpetology.* Hafner Publishing Company; New York.
- Peters, U. 1976. Gelungene Riesenschlangenzucht im Tarongo Zoo. *Aquar.-Mag. (Stuttgart)* 12: 525-527.
- Peterson, K.H. 1989. Further comments on anecdotal data. *Bull. Chicago Herpetol. Soc.* 24: 190-191.
- Petzold, H.-G. 1968. Zur Fortpflanzungsbiologie asiatischer Kobras (Naja naja). *Zool. Gart. N.F. (Leipzig)* 36: 133-146.
- Petzold, H.-G. 1971. Blindschleiche und Scheltopusik. *D. Neue Brehm - Bucherei (Wittenberg)* 448: 85-86.

- Pfeffer, P. 1959. Observations sur le varan de Komodo Varanus komodoensis.  
Ouwens 1912. Terre Vie 106: 195-243.
- Phelps, T. 1981. Poisonous Snakes. Blandford Press; Poole, Dorset, U.K.
- Pianka, E.R., and W.F. Giles. 1982. Notes on the biology of two species of nocturnal skinks, Egernia inornata and Egernia striata, in the Great Victoria Desert. West. Australian Nat. 15(2): 8-13.
- Picado T., C. 1931. Serpientes Venenosas de Costa Rica. Sus Venenos. Seroterapia anti-ofídica. Sauter, Arias & Co.; San José.
- Pienaar, U. DeV., W.D. Haacke, and N.H.G. Jacobsen. 1978. The Reptiles of the Kruger National Park. National Parks Board of South Africa; Pretoria.
- Pinkert, --. 1893. Brütende Riesenschlangen. Zool. Gart. (Frankfurt Main) 34: 319.
- Pitman, C.R.S. 1938. A Guide to the Snakes of Uganda. Uganda Society; Kampala.
- Platt, D.R. 1969. Natural history of the hognose snakes Heterodon platyrhinos and Heterodon nasicus. Univ. Kansas Publ. Mus. Nat. Hist. 18: 253-420.
- Pliny the Elder [=G. Plinius Secundus]. [d. AD 79]. Naturalis Historia. Translated in J. Bostock and H.T. Riley (eds). 1855. The Natural History of Pliny. Vol. II. Henry G. Bohn; London.
- Pope, C.H. 1929. Notes on reptiles from Fukien and other Chinese provinces. Bull. Amer. Mus. Nat. Hist. 58: 335-487.
- Pope, C.H. 1935. The Reptiles of China. Turtles, Crocodilians, Snakes, Lizards. Natural History of Central Asia. Vol. X. American Museum of Natural History; New York.
- Pope, C.H. 1946. Snakes of the Northeastern United States. New York Zoological Society; New York.
- Pope, C.H. 1955. The Reptile World. Alfred A. Knopf; New York.
- Pope, C.H. 1961. The Giant Snakes. Alfred A. Knopf; New York.
- Price, A.H. 1988. Observations on maternal behavior and neonate aggregation in the western diamondback rattlesnake, Crotalus atrox (Crotalidae). Southwest. Nat. 33: 370-373.
- Ramsey, R.W., and A. Travis. 1960. The Common Wildlife of Panama. Russell W. Ramsey and Amos Travis, Printing Plant Mount Hope; Canal Zone.
- Rand, W.M., and A.S. Rand. 1976. Agonistic behaviour in nesting iguanas: a stochastic analysis of dispute settlement dominated by the minimization of energy cost. Z. Tierpsychol. 40: 279-299.
- Rao, [H.]S. 1957. History of our knowledge of the Indian fauna through the ages. J. Bombay Nat. Hist. Soc. 54: 251-280.
- Rebouças-Spieker, R., and P.E. Vanzolini. 1978. Parturition in Mabuya macrorhyncha Hoge, 1946 (Sauria, Scincidae), with a note on the distribution of maternal behavior in lizards. Pap. Avulsos Zool. São Paulo 32: 95-99.
- Reese, A.M. 1922. A note on the breeding habits of tegu. Copeia (110): 69-72.
- Reinert, H.K., and W.R. Kodrich. 1982. Movements and habitat utilization by the massasauga, Sistrurus catenatus catenatus. J. Herp. 16: 162-171.
- Reinert, H.K., and R.T. Zappalorti. 1988. Field observation of the association of adult and neonatal timber rattlesnakes, Crotalus horridus, with possible evidence for conspecific trailing. Copeia 1988: 1057-1059.
- Reitinger, F.F., and J.K.S. Lee. 1978. Common Snakes of South East Asia and Hong Kong. Heinemann Educational Books; Kowloon, Hong Kong.

- Reynolds, A.E. 1959. Eggs and young of the lizard, Eumeces fasciatus. Proc. Indiana Acad. Sci. 68: 367-378.
- Reynolds, F.A., and A.N. Solberg. 1942. Notes on the life history of the mud snake. Copeia 1942: 25-26.
- Riemer, W.J. 1957. The snake Farancia abacura: an attended nest. Herpetologica 13: 31-32.
- Rieppel, O. 1970. Nachwuchs bei Dryophis nasutus (Lacépède) 1789. Aqua Terra 7: 85-88.
- Rivers, G.M. 1874. The rattlesnake--Its poison and antidote. South. Med. Rec. 4: 505-517.
- Robb, J. 1980. New Zealand Amphibians and Reptiles in Colour. William Collins Publishers; Auckland.
- Roosevelt, T. 1926. Through the Brazilian Wilderness. Charles Scribner's Sons; New York.
- Rose, W. 1929. Veld and Vlei. An Account of South African Frogs, Toads, Lizards, Snakes, & Tortoises. Specialty Press of South Africa; Cape Town.
- Rose, W. 1950. The Reptiles and Amphibians of Southern Africa. Maskew Miller; Cape Town.
- Rösler, H. 1988. Über das "Eirfressen" in Terrarium bei Arten der Gattung Phelsuma Gray, 1825 (Sauria: Gekkonidae). Salamandra 24: 20-26.
- Ross, R.[A.] 1973. Successful mating and hatching of Children's python, Liasis childreni. HISS News-J. 1: 181-182.
- Ross, R.A. 1978. The Python Breeding Manual. Institute for Herpetological Research; [Stanford].
- Ross, R.[A.] 1980a. Erfolgreiche Zucht des Gefleckten Python (Liasis childreni). Herpetofauna (Weinstadt) (9): 28-29.
- Ross, R.[A.] 1980b. The breeding of pythons (subfamily Pythoninae) in captivity. (1):135-139. In J.B. Murphy and J.T. Collins (eds), Reproductive Biology and Diseases of Captive Reptiles. SSAR Contr. Herp. Society for the Study of Amphibians and Reptiles; [Athens, Ohio].
- Ross, R.[A.] 1983. Reproduction of the Children's python (Liasis childreni) in a terrarium. Litt. Serpentiolum (Engl. Ed.) 3: 18-21.
- Ross, R.[A.], and R. Larman. 1977. Captive breeding in two species of python Liasis albertisii and L. mackloti. Int. Zoo Yearb. 17: 133-136.
- Russell, F.E. 1983. Snake Venom Poisoning. Reprint Edition. Scholium International; Great Neck, New York.
- Ruthven, A.G. 1911. A biological survey of the sand dune region on the south shore of Saginaw Bay, Michigan. Michigan Geol. Biol. Surv. Ser 2 (4): 1-347.
- Schivre, M. 1972. Observations sur la reproduction de Python regius (Shaw). Aquarama 6: 67.
- Schleich, H.H., and W. Kästle. 1988. Reptile Egg-shells. SEM Atlas. Gustav Fischer Verlag; Stuttgart/New York.
- Schlott, M. 1935. [Nachrichten aus Zoologischen Gärten] Breslau. Zool. Gart. N.F. (Leipzig) 8: 74-77.
- Schmidt, K.P. 1927. Notes on Chinese reptiles. Bull. Amer. Mus. Nat. Hist. 54: 467-551.
- Schmidt, K.P. 1929. The truth about snake stories. Sci. Amer. 141: 134-136.
- Schuette, B. 1980. Two broad-headed skink nests. J. St. Louis Herpetol. Soc. 7(3-4): 13-14.
- Schütte, G.W. 1970. Reptile incubation. Lammergeyer (11): 85.
- Sclatter, P.L. 1862. Notes on the incubation of Python sebae, as observed in the Society's Gardens. Proc. Zool. Soc. London 1862: 365-368.



- Semper, K. 1881. *Animal Life as Affected by the Natural Conditions of Existence*. D. Appleton and Company; New York.
- Sengoku, S. 1979. [Amphibians and Reptiles.] [in Japanese]. Ienohikari Corporation; Tokyo.
- Service, R. 1902. The adder in Solway. *Ann. Scottish Nat. Hist.* 44: 153-162.
- Sharrell, R. 1966. *The Tuatara, Lizards and Frogs of New Zealand*. Collins; London.
- Shaw, C.E. 1949. Notes on two broods of xantusiids. *Herpetologica* 5: 23-26.
- Shaw, C.E. 1954. Captive-bred Cuban iguanas Cyclura macleayi macleayi. *Herpetologica* 10: 73-78.
- Shaw, C.E. 1969. Breeding the rhinoceros iguana Cyclura cornuta cornuta at San Diego Zoo. *Int. Zoo Yearb.* 9: 45-48.
- Shaw, G.E., and E.O. Shebbeare. 1931. The snakes of northern Bengal and Sikkim. Poisonous snakes. *J. Darjeeling Nat. Hist. Soc.* 5: 3-8.
- Sheargold, T. 1979. Notes on the reproduction of Children's pythons (Liasis childreni) Gray 1842. *Herpetofauna (Sydney)* 10(2): 2-4.
- Shine, R. 1980. "Costs" of reproduction in reptiles. *Oecologia (Berlin)* 46: 92-100.
- Shine, R. 1985. The evolution of viviparity in reptiles: an ecological analysis. 15: 605-694. In G. Gans and F. Billet (eds), *Biology of the Reptilia. Development B.* John Wiley & Sons; New York.
- Shine, R. 1988. Parental care in reptiles. 16: 275-329. In G. Gans and R.B. Huey (eds), *Biology of the Reptilia. Ecology B. Defense and Life History.* Alan R. Liss; New York.
- Shine, R. 1989. Constraints, allometry, and adaptation: food habits and reproductive biology of Australian brownsnakes (Pseudonaja: Elapidae). *Herpetologica* 45: 195-207.
- Shine, R., and J.J. Bull. 1979. The evolution of live-bearing in lizards and snakes. *Amer. Nat.* 113: 905-923.
- Silver, R., H. Andrews, and G. F. Ball. 1985. Parental care in an ecological perspective: a quantitative analysis of avian subfamilies. *Amer. Zool.* 25: 823-840.
- Simmon, E.S. 1944. The breeding habits of the cobra (Naja tripudians Merrem) and the green whipsnake (Dryophis mycterizans). *J. Bombay Nat. Hist. Soc.* 44: 480-481.
- Skutch, A.F. 1957. The incubation patterns of birds. *Ibis* 99: 69-93.
- Skutch, A.F. 1976. *Parent Birds and Their Young*. University of Texas Press; Austin, Texas.
- Slavens, F.L. 1982. *Inventory of Live Reptiles and Amphibians in Captivity Current January 1, 1982*. Frank L. Slavens; Seattle.
- Slavens, F.L. 1983. *Inventory of Live Reptiles and Amphibians in Captivity Current January 1, 1983*. Frank L. Slavens; Seattle.
- Slavens, F.L. 1984. *Inventory of Live Reptiles and Amphibians in Captivity Current January 1, 1984*. Frank L. Slavens; Seattle.
- Slavens, F.L. 1985. *Inventory of Live Reptiles and Amphibians in Captivity Current January 1, 1985*. Frank L. Slavens; Seattle.
- Slavens, F.L. 1986. *Inventory of Live Reptiles and Amphibians in Captivity Current January 1, 1986*. Frank L. Slavens; Seattle.
- Slavens, F.L. 1987. *Inventory of Live Reptiles and Amphibians in Captivity Current January 1, 1987*. Frank L. Slavens; Seattle.
- Slavens, F.L. 1988. *Inventory, Longevity, & Breeding Notes--Reptiles and Amphibians in Captivity Current January 1, 1988*. Frank L. Slavens; Seattle.

- Slip, D.J. 1986. Ecology and thermal biology of the diamond python, Morelia spilota spilota. Unpublished Master's Thesis; University of Sydney.
- Slip, D.J., and R. Shine. 1988a. The reproductive biology and mating system of diamond pythons, Morelia spilota (Serpentes: Boidae). *Herpetologica* 44: 396-404.
- Slip, D.J., and R. Shine. 1988b. Reptilian endothermy: a field study of thermoregulation by brooding pythons. *J. Zool. (London)* 216: 367-378.
- Smedley, N. 1931. Oviparity in a sea-snake, Laticauda colubrina (Schneid). *Bull. Raffles Mus. (Singapore)* (5): 54-59.
- Smith, H.C. 1936. A hamadryad's (Naia bungarus) nest and eggs. *J. Bombay Nat. Hist. Soc.* 39: 186.
- Smith, H.M. 1946. *Handbook of Lizards. Lizards of the United States and Canada.* Comstock Publishing Company; Ithaca and London.
- Smith, H.M., and J.A. Slater. 1949. The southern races of Eumeces septentrionalis (Baird). *Trans. Kansas Acad. Sci.* 52: 438-448.
- Smith, M.[A.] 1915. Notes on some snakes from Siam. *J. Bombay Nat. Hist. Soc.* 23: 784-789.
- Smith, M.A. 1935. *The Fauna of British India, Including Ceylon and Burma. Reptilia and Amphibia. Vol. II. -Sauria.* Taylor and Francis; London.
- Smith, M.[A.] 1937. Breeding habits of the Indian cobra. *J. Siam Soc. Nat. Hist. Suppl.* 11: 62-63.
- Smith, M.A. 1943. *The Fauna of British India Ceylon and Burma. Reptilia and Amphibia. Vol. III.-Serpentes.* Taylor and Francis; London.
- Smith, M.[A.] 1951. *The British Amphibians & Reptiles.* Collins; London.
- Smith, P.W. 1961. The amphibians and reptiles of Illinois. *Illinois Nat. Hist. Surv. Bull.* 28: 1-298.
- Snyder, D.H. 1972. *Amphibians and Reptiles of Land Between the Lakes.* Tennessee Valley Authority; [Knoxville].
- Soderberg, P. 1973. On eleven Asian elapid snakes with specific reference to their occurrence in Thailand. *Nat. Hist. Bull. Siam Soc.* 24: 203-317.
- Somma, L.A. 1985a. Notes on maternal behavior and post-brooding aggression in the prairie skink Eumeces septentrionalis. *Nebraska Herpetol. Newsl.* 6(4): 9-12.
- Somma, L.A. 1985b. Brooding behavior in the northern prairie skink, Eumeces septentrionalis septentrionalis (Baird), and its relationship to the hydric environment of the nest substrate. Unpublished M.A. Thesis; University of Nebraska at Omaha, [Omaha].
- Somma, L.A. 1985c. Egg brooding in the prairie skink, Eumeces septentrionalis, and its relationship to the hydric environment of the nest substrate. *Soc. Stud. Amphib. Rept. -Herp. Leag. Meet.* [Unpubl. Abst.] (Tampa, Florida) 1985: 78.
- Somma, L.A. 1987a. Alloparental care in the prairie skink, Eumeces septentrionalis: a case of mistaken identity? *Nebraska Herpetol. Newsl.* 8(2): 5-8.
- Somma, L.A. 1987b. Reproduction of the prairie skink, Eumeces septentrionalis, in Nebraska. *Great Basin Nat.* 47: 373-374.
- Somma, L.A. 1987c. Maternal care of neonates in the prairie skink, Eumeces septentrionalis. *Great Basin Nat.* 47: 536-537.
- Somma, L.A. 1988. Comments on the use of the term "brooding" to describe parental behaviour in squamate reptiles. *Amphibia-Reptilia* 9: 89-91.
- Somma, L.A. 1989a. Oophagous behavior in brooding prairie skinks, Eumeces septentrionalis. *Herpetol. Rev.* 20: 3-4.
- Somma, L.A. 1989b. The water vapor conductance of squamate reptilian eggs: the influence of scaling on nesting ecology. *First World Congr. Herp.* [Unpubl.] Abst. (Canterbury, United Kingdom) 1: [278].



- Somma, L.A., and P.A. Cochran. 1989. Bibliography and subject index of the prairie skink, Eumeces septentrionalis (Baird) (Sauria: Scincidae). Great Basin Nat. 49: in press.
- Somma, L.A., and J.D. Fawcett. 1985. An adaptive function for egg brooding in the prairie skink, Eumeces septentrionalis. Anim. Behav. Soc. [Unpubl. Meet.] Abst. (Raleigh, North Carolina) 1985: 71.
- Somma, L.A., and J.D. Fawcett. 1989. Brooding behaviour of the prairie skink, Eumeces septentrionalis, and its relationship to the hydric environment of the nest. Zool. J. Linn. Soc. 95: 245-256.
- Speck, F.G. 1921. The origin of the belief that snakes swallow their young for protection. Copeia (98): 51-54.
- Speck, F.G. 1923. Snake folk-lore: the snake who swallows her young. J. Amer. Folk-lore 36: 298-300.
- Spenser, E. 1590. The Faerie Queene. William Ponfonbie; London.
- Spotila, J.R., and M.P. O'Conner. 1989. Eggs as organisms--the difficulties of staying in one place. First World Congr. Herp. [Unpubl.] Abst. (Canterbury, United Kingdom) 1: [279].
- Stafford, P.J. 1986. Pythons and Boas. T.F.H. Publications; Neptune City, New Jersey.
- Stanley, C. [19??]. The Life and Adventures of the American Cow-boy. Clark Stanley Snake Oil Liniment Company; Providence, Rhode Island.
- Steiner, T.M. 1986. Eumeces inexpectatus. Cat. Amer. Amphib. Rept. (385): 1-2.
- Stemmler, O. 1969. Eine erfolgreiche Zucht von Kurzpythons (Python curtus brongersmai Stull 1938) im Terrarium. Aquar. Terrar. (Leipzig) 15: 156-161.
- Stemmler-Morath, C. 1956. Beitrag zur Gefangenschafts- und Fortpflanzungsbiologie von Python molurus L. Zool. Gart. N.F. (Leipzig) 21: 347-364.
- Stewart, J.R., and D. Duvall. 1985. Ovariectomy fails to block egg-brooding behavior in nesting, female five-lined skinks (Eumeces fasciatus). Copeia 1985: 777-779.
- Strecker, J.K. 1908. Notes on the breeding habits of Phrynosoma cornutum and other Texas lizards. Proc. Biol. Soc. Washington 21: 165-170.
- Street, D. 1979. The Reptiles of Northern and Central Europe. B. T. Batsford; London.
- Strong, J. 1961. The Exhaustive Concordance of the Bible. Abingdon Press; New York/Nashville.
- Sundowner. [=Tichborne, H. (sic?)] 1895. Snakes. European Mail; London.
- Sundowner. [=Tichborne, H. (sic?)] 1902. The Tale of the Serpent. Chatto & Windus; London.
- Sweeney, R.C.H. 1961. Snakes of Nyasaland. The Nyasaland Society and the Nyasaland Government; Zomba.
- Switak, K.H. 1975. Der Grüne Baumpython aus dem Land der Menschenfresser. Aquar. Mag. 9: 366-372.
- Tanner, W.W. 1943. Notes on the life history of Eumeces skiltonianus skiltonianus. Great Basin Nat. 4: 81-88.
- Tanner, W.W. 1957. A taxonomic and ecological study of the western skink (Eumeces skiltonianus). Great Basin Nat. 17: 59-94.
- Tanner, W.W. 1987. Lizards and turtles of western Chihuahua. Great Basin Nat. 47: 383-421.
- Tarbet, S.A. 1984. Reproduction of D'albert's python, Liasis albertisi at the Oklahoma City Zoo. [7]:132-133. In P.J. Tolson (ed), 7th Annual

- Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Taylor, E.H. 1932a. Eumeces inexpectatus: a new American lizard of the family Scincidae. Univ. Kansas Sci. Bull. 20: 251-261.
- Taylor, E.H. 1932b. Eumeces laticeps: a neglected species of skink. Univ. Kansas Sci. Bull. 20: 263-271.
- Taylor, E.H. 1935. A taxonomic study of the cosmopolitan scincoid lizards of the genus Eumeces with an account of the distribution and relationships of its species. Univ. Kansas Sci. Bull. 23: 1-643.
- Taylor, E.H. 1956. A review of the lizards of Costa Rica. Univ. Kansas Sci. Bull. 38: 1-332.
- Taylor, E.H. 1965. The serpents of Thailand and adjacent waters. Univ. Kansas Sci. Bull. 45: 609-1096.
- Taylor, T. 1985. Eumeces callicephalus (mountain skink). Reproduction. Herpetol. Rev. 16: 27.
- Telford, S.R., Jr. 1952. A herpetological survey in the vicinity of Lake Shipp, Polk County, Florida. Quart. J. Florida Acad. Sci. 15: 175-185.
- Telford, S.R., Jr. 1959. A study of the sand skink, Neoseps reynoldsi Stejneger. Copeia 1959: 110-119.
- Tho, Y.P., and S.Y. Ho. 1979. Observations on a batch of Gekko stentor eggs. Malayan Nat. J. 32: 265-269.
- Thompson, M.B. 1988. Nesting behavior of tuatara, Sphenodon punctatus. Comb. Meet. Herp. Leag., Amer. Elasm. Soc., Early Life Hist. Sect. A.F.S., Soc. Stud. Amphib. Rept., Amer. Soc. Ichth. Herp. [Unpubl.] Abst. Prog. (Ann Arbor, Michigan) 1988: 182.
- Tinkle, D.W. 1959. Observations of reptiles and amphibians in a Louisiana swamp. Amer. Midl. Nat. 62: 189-205.
- Tinkle, D.W., and J.W. Gibbons. 1977. The distribution and evolution of viviparity in reptiles. Misc. Publ. Mus. Zool. Univ. Michigan (154): 1-55.
- Topsell, E. 1608. The Historie of Serpents. Or, the Second Booke of Liuing Creatures. William Taggard; London.
- Townson, S. 1980. Observations on the reproduction of the Indian python in captivity, with special reference to the interbreeding of the two subspecies, Python molurus molurus and Python molurus bivittatus. Pp. 69-80. In S. Townson, N.J. Millichamp, D.G.D. Lucas and A.J. Millwood (eds), The Care and Breeding of Captive Reptiles. British Herpetological Society; [London].
- Townson, S. [1989]. A captive Burmese python incubating eggs. British Herpetological Society. [Unpubl. pamphlet]. [p. 3, photo].
- Townson, S. 1985. The captive reproduction and growth of the yellow anaconda (Eunectes notaeus). Pp. 33-43. In S. Townson and K. Lawrence (eds), Reptiles: Breeding, Behaviour and Veterinary Aspects. British Herpetological Society; London.
- Townson, S., and S. Todd. 1976. Captive breeding of the Indian python Python molurus bivittatus. Brit. Herpetol. Soc. Newsl. 14: 25-26.
- Toyama, M. 1975. On the embryos obtained from eggs of Eumeces oshimensis. Japan. J. Herp. 6: 39-42.
- Trutnau, L. 1980. Über die Lebensweise und Pflege des Netzpythons Python reticulatus (Schneider). Herpetofauna (Weinstadt) (4): 26-31.
- Trutnau, L. 1986. Nonvenomous Snakes. English Language Edition. Barron's Educational Series; Woodbury, New York.
- Trillmich, K. 1979. Feeding behaviour and social behaviour of the marine iguana. Noticias Galápagos 29: 17-20.

- Tryon, B.W. 1979. Reproduction in captive forest cobras, Naja melanoleuca (Serpentes: Elapidae). J. Herp. 13: 499-504.
- Tryon, B.[W.] 1985. First Papuan python hatching reported. AAZPA Newsl. 26(3): 17.
- Tryon, B.W., and J. Whitehead. 1988. Reproduction in a little-known New Guinea python, Liasis papuanus (Peters and Doria). Zoo Biol. 7:371-379.
- Tweedie, M.W.F. 1954. The Snakes of Malaya. Government Printing Office; Singapore.
- Tytle, T., S. Wheeler, and D. Grow. 1987. Ptychozoon lionatum (gliding gecko). Egg guarding. Herpetol. Rev. 18: 36.
- Valenciennes, [A.] 1841a. Observations faites pendant l'incubation d'une femelle du python à deux raies (Python bivittatus, Kuhl.) pendant les mois de mai et de juin 1841. Compt. Rend. Hebd. Séan. Acad. Sci. Paris 13: 126-133.
- Valenciennes, [A.] 1841b. Observations faites pendant l'incubation d'une femelle du python à deux raies (Python bivittatus, Kuhl.) pendant les mois de mai et de juin 1841. Ann. Sci. Nat., 2nd Ser. (Paris) 16: 65-72.
- Van Denburgh, J. 1922. The Reptiles of Western North America. Volume I. Lizards. Occ. Pap. California Acad. Sci. 10: 1-611.
- Van Devender, T.R., and W. Van Devender. 1975. Ecological notes on two Mexican skinks (genus Eumeces). Southwest Nat. 20: 279-282.
- Van Mierop, L.H.S., and S.M. Barnard. 1976a. Thermoregulation in a brooding female Python molurus bivittatus (Serpentes: Boidae). Copeia 1976: 398-401.
- Van Mierop, L.H.S., and S.M. Barnard. 1976b. Observations on the reproduction of Python molurus bivittatus. J. Herp. 10: 333-340.
- Van Mierop, L.H.S., and S.M. Barnard. 1978. Further observations on thermoregulation in brooding female Python molurus bivittatus (Serpentes: Boidae). Copeia 1978: 615-621.
- Van Mierop, L.H.S., and E.L. Bessette. 1981. Reproduction of the ball python, Python regius, in captivity. Herpetol. Rev. 12: 20-22.
- Van Mierop, L.H.S., T. Walsh, and D.L. Marcellini. 1983. Reproduction of Chondropython viridis (Reptilia, Serpentes, Boidae). [6]:265-274. In D.L. Marcellini (ed), 6th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Villiard, P. 1969. Reptiles as Pets. Doubleday & Company; Garden City.
- Vinegar, A. 1968. Brooding of the eastern glass lizard, Ophisaurus ventralis. Bull. South. California Acad. Sci. 67: 65-68.
- Vinegar, A. 1973. The effects of temperature on the growth and development of embryos of the Indian python, Python molurus (Reptilia: Serpentes: Boidae). Copeia 1973: 171-173.
- Vinegar, A. 1974. Evolutionary implications of temperature induced anomalies of development in snake embryos. Herpetologica 30: 72-74.
- Vinegar, A., V.H. Hutchison, and H.G. Dowling. 1970. Metabolism, energetics, and thermoregulation during brooding of snakes of the genus Python (Reptilia, Boidae). Zoologica 55: 19-48.
- Visser, J. 1971. Unusually large skaapesteker eggs from the Cape peninsula. J. Herpetol. Assoc. Africa (7): 9.
- Vitt, L.J., and W.E. Cooper, Jr. 1985a. The relationship between reproduction and lipid cycling in the skink Eumeces laticeps with comments on brooding ecology. Herpetologica 41: 419-432.
- Vitt, L.J., and W.E. Cooper, Jr. 1985b. The evolution of sexual dimorphism in the skink Eumeces laticeps: an example of sexual selection. Canad. J. Zool. 63: 995-1002.

- Vitt, L.J., and W.E. Cooper, Jr. 1986. Skink reproduction and sexual dimorphism: Eumeces fasciatus in the southeastern United States, with notes on Eumeces inexpectatus. J. Herp. 20: 65-76.
- Vitt, L.J., and W.E. Cooper, Jr. 1989. Maternal care in skinks (Sauria: Eumeces). J. Herp. 23: 29-34.
- Vogel, Z. [1958?]. Reptile Life. Spring Books; London.
- Vogel, Z. 1964. Reptiles and Amphibians. Their Care and Behaviour. English Language Edition. Studio Vista; London.
- Vogt, R.C. 1981. Natural History of Amphibians and Reptiles in Wisconsin. Milwaukee Public Museum; Milwaukee.
- Wagner, E.P. 1973. Breeding Python molurus bivittatus. HISS News-J. 1: 113.
- Wagner, E.[P.] 1976. Breeding the Burmese python Python molurus bivittatus at Seattle Zoo. Int. Zoo Yearb. 16: 83-85.
- Waitkus, D. 1983. Reproductive behavior of Ptychozoon lionatum at the Reptile Breeding Foundation. Notes NOAH 11(3): 16.
- Wall, F. 1907. Hatching of the dhaman (Zamensis mucosus) eggs, and observations on the egg tooth. J. Bombay Nat. Hist. Soc. 17: 1033-1035.
- Wall, F. 1908. Notes on the incubation and brood of the Indo-Burmese snake-lizard or slow worm (Ophisaurus gracilis). J. Bombay Nat. Hist. Soc. 18: 503-504.
- Wall, F. 1911. A popular treatise on the common Indian snakes [part XIV with plate XIV]. J. Bombay Nat. Hist. Soc. 20: 603-633.
- Wall, F. 1912. A popular treatise on the common Indian snakes [part XVII with plate XVII]. J. Bombay Nat. Hist. Soc. 21: 447-475.
- Wall, F. 1921. Ophidia Taprobanica or the Snakes of Ceylon. H. R. Cottle, Government Printer; Colombo.
- Wall, F. 1924. The hamadryad or king cobra, Naia hannah (Cantor). J. Bombay Nat. Hist. Soc. 30: 189-195.
- Walsh, T. 1977. Husbandry and breeding of Chondropython viridis. Natl. Assoc. Sound Wildlife Prog. 1(2): 10-22.
- Walsh, T. [1980]. Further notes on the husbandry, breeding, and behavior of Chondropython viridis. [3]:102-111. In R.A. Hahn (ed), 3rd Annual Reptile Symposium on Captive Propagation and Husbandry. [Zoological Consortium]; Thurmont, Maryland.
- Walsh, T., and B. Davis. 1984. Husbandry and breeding of the Brazilian rainbow boa Epicrates cenchira at the National Zoological Park. [7]:108-114. In P.J. Tolson (ed), 7th Annual Reptile Symposium on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- [Wang, B.] 1966. [Studies on the ecology of four species of lizards in Hangchow. II. Breeding.] [in Chinese]. Acta Zool. Sinica 19: 170-186.
- Wasey, G.K. 1892. A nest of king cobra's eggs. J. Bombay Nat. Hist. Soc. 7: 257.
- Wehekind, L. 1960. Trinidad snakes. J. British Guiana Mus. Zoo Royal Agr. Commer. Soc. (27): 71-76.
- Wells, E. 1981. A diurnal variation's effect on a captive breeding of Boa constrictor constrictor. [4]:11-16. In R.A. Hahn (ed), 4th Annual Reptile Symposium on Captive Propagation & Husbandry. Zoological Consortium; Thurmont, Maryland.
- Wells, K.D. 1981. Parental behavior of male and female frogs. Pp. 184-197. In R.D. Alexander and D.W. Tinkle (eds), Natural Selection and Social Behavior. Recent Research and New Theory. Chiron Press; New York.
- Welty, J.C. 1982. The Life of Birds. Third Edition. CBS College Publishing; Philadelphia/New York.

- Wendt, H. 1965. The Sex Life of the Animals. English Translation. Simon and Schuster; New York.
- Werler, J.E. 1951. Miscellaneous notes on the eggs and young of Texan and Mexican reptiles. *Zoologica* 36: 37-48.
- Werner, D.I. 1982. Social organization and ecology of land iguanas, Conolophus subcristatus, on Isla Fernandina, Galápagos. Pp. 342-365. In G.M. Burghardt and A.S. Rand (eds), *Iguanas of the World. Their Behavior, Ecology, and Conservation*. Noyes Publications; Park Ridge, New Jersey.
- Wexo, J.B. 1987. Snakes. *ZOOBOOKS* 4(10): 1-16.
- Wharton, C.H. 1960. Birth and behavior of a brood of cottonmouths, Agkistrodon piscivorus piscivorus, with notes on tail-luring. *Herpetologica* 16: 125-129.
- Wharton, C.H. 1966. Reproduction and growth in the cottonmouth, Agkistrodon piscivorus Lacépède, of Cedar Keys, Florida. *Copeia* 1966: 149-161.
- Whitaker, R. 1977. Two reptile nests. *Hornbill* 1977 (April-June): 7-12.
- Whitaker, R. 1978. *Common Indian Snakes*. Macmillan India; Dehli.
- Whitaker, Z., and R. Whitaker. 1986. *The Snakes Around Us*. Director, National Book Trust; New Dehli.
- White, G. 1789. *The Natural History and Antiquities of Selborne, in the County of Southampton*. B. White and Son; London.
- White, T.H. 1954. *The Bestiary. A Book of Beasts*. G. P. Putnam's Sons; New York.
- Whitfield, P. 1983. *Reptiles and Amphibians*. Longman Group; Essex.
- Wiewandt, T.A. 1977. Ecology, behavior, and management of the Mona Island ground iguana, Cyclura stejnegeri. Unpublished Ph.D. Dissertation; Cornell University, [Ithaca].
- Wiewandt, T.A. 1979. La gran iguana de Mona. *Nat. Hist.* 88(10): 56-65.
- Wiewandt, T.A. 1982. Evolution of nesting patterns of iguanine lizards. Pp. 119-141. In G.M. Burghardt and A.S. Rand (eds), *Iguanas of the World. Their Behavior, Ecology, and Conservation*. Noyes Publications; Park Ridge, New Jersey.
- Williams, G.C. 1966. *Adaptation and Natural Selection. A Critique of Some Current Evolutionary Thought*. Princeton University Press; Princeton.
- Williamson, M.A. 1986. *Trail of the Snake. From Big Bend to Baja*. Sunstone Press; Santa Fe, New Mexico.
- Wittenberger, J.F. 1981. *Animal Social Behavior*. Duxbury Press; Boston.
- Woodbury, A.M. 1931. A descriptive catalog of the reptiles of Utah. *Bull. Univ. Utah* 21(5); *Biol. Ser.* 1(4): 1-129.
- Woodward, S.F. 1933. A few notes on the persistence of active spermatozoa in the African night-adder, Causus rhombeatus. *Proc. Zool. Soc. London* 1933: 189-190.
- Wray, G.O. 1862. [Extract of a letter from Wray, G.O.] *Proc. Zool. Soc. London* 1862: 107-108.
- Wright, A.H., and A.A. Wright. 1957. *Handbook of Snakes of the United States and Canada. Vols. I & II*. Comstock Publishing Company; Ithaca and London.
- Yadav, R.N. 1967. A note on the breeding of Indian pythons Python molurus at Jaipur Zoo. *Int. Zoo Yearb.* 7: 182-183.
- Yahya, H.S.A. 1985. Observations on the banded krait, Bungarus fasciatus. *J. Bombay Nat. Hist. Soc.* 82: 219.
- York, D.S., and G.M. Burghardt. 1988. Brooding in the Malayan pit viper, Calloselasma rhodostoma: temperature, relative humidity, and defensive behaviour. *Herpetol. J.* 6: 210-214.

- Zaworski, J.P. 1987a. Egg molding behavior by male Gekko gecko and Gekko petricolus. Bull. Chicago Herpetol. Soc. 22: 77.
- Zaworski, J.P. 1987b. The captive maintenance and propagation of Gekko petricolus Taylor. Bull. Chicago Herpetol. Soc. 22: 129-130.
- Zaworski, J.P. 1987c. Egg guarding behavior by a male Gekko gecko. Bull. Chicago Herpetol. Soc. 22: 193.
- Zaworski, J.P. 1988. Captive husbandry of the tokay, Gekko gecko Linnaeus. Bull. Chicago Herpetol. Soc. 23: 104-105.
- Zimmermann, E. 1986. Breeding Terrarium Animals. English-language Edition. T.F.H. Publications; Neptune City, New Jersey.
- Zulich, A.W. 1983. Notes on the maternal incubation of Chondropython viridis eggs. J. North. Ohio Herpetol. Soc. 9(1): 13-21.
- Zulich, A.W. 1985. Captive breeding, husbandry, and neonatal care of the green tree python Chondropython viridis with emphasis on maternal incubation of eggs. [8]:114-118. In S. McKeown and F. Caporaso (eds), 8th Annual International Herpetological Symposium on Captive Propagation & Husbandry. Zoological Consortium; Thurmont, Maryland.
- Zweifel, R.G. 1962. Notes on the distribution and reproduction of the lizard Eumeces callicephalus. Herpetologica 18: 63-65.











6  
7

AMPHIBIANS AND REPTILES AS PREY  
OF  
BIRDS  
IN  
SOUTHWESTERN EUROPE



JOSÉ MARTÍN  
&  
PILAR LÓPEZ

Museo Nacional de Ciencias Naturales de Madrid



SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
NO. 82

1990

**SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE**

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with the Division of Amphibians and Reptiles.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions on preparation and submission. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington DC 20560 USA. Please include a self-addressed mailing label with requests.

## INTRODUCTION

This review surveys species of amphibians and reptiles that are reported prey of birds in southwestern Europe. These ectothermic animals are important in Mediterranean and temperate ecosystems, because they are the prey for many specialist birds (eg, the short-toed eagle, Circaetus gallicus, a snake eater). However, European herpetologists know the identity of few amphibian and reptilian predators. In the recently published "Handbuch der Reptilien und Amphibien Europas", references to predation are scarce and even ignored. However, the ornithological literature contains numerous papers on bird diets, and amphibians and reptiles are frequently cited as bird preys.

The Cramp's "Handbook of Birds of the Western Palearctic" was the first book examined; however, most of the predation records were obtained from the main European ornithological journals from Southwestern Europe. Including the Iberian Peninsula (Spain and Portugal), France, Belgium, The Netherlands, and the British Isles -approximately between 10°W and 10°E longitude.

References are listed and numbered alphabetically and can be accessed through either predator or prey. Species are listed systematically.

All scientific names have been updated. Some report provide only general identifications i.e., frog, toad, lizard or snake. In such cases, we have tried to identify the prey precisely as possible. We had problems with the "green frog" group, which includes Rana ridibunda, R. perezi, R. lessonae, and R. esculenta. Since these species are taxonomically complex, we cited them here as green frogs. The same identification problems occurred with the genera Discoglossus, Hyla, and Podarcis.

The number of references is not a good index for predation rate, it only means that one species is more frequently cited. Although some preys occur only occasionally in bird diet, other species apparently experience heavy predation. This aspect is often ignored in ecological studies.

Please remembered that this survey was limited to regional journals and books and complete only for 1950 through 1988 for these. The journals are: Belgium (Le Gerfaut), France (Acta Biologica Montana, Alauda, Aves, Le Bièvre, Le Cormoran, L'Oiseau et la Revue Francaise

d'Ornithologie, Nos Oiseaux), Great Britain (Bird Study, British Birds, The Ibis, Scottish Birds), Portugal (Cyanopica), Spain (Alytes, Ardeola, Boletín de la Estación Central de Ecología, Boletín de la Real Sociedad de Historia Natural, Cuadernos de Ciencias Biológicas, Doñana Acta Vertebrata, Mediterránea, Miscellanea Zoologica, Monografías del I.C.O.N.A., Munibe, Naturalia Hispanica, Publicaciones del Centro Pirenaico de Biología Experimental).

**Acknowledgments:** The Sociedad Española de Ornitología library allowed access to these journals. E. Moreno helped us with the English translation.

## BIRDS THAT EAT AMPHIBIANS AND REPTILES

| <u>PREDATOR</u>         | <u>PREY</u>           | <u>REFERENCES</u> |
|-------------------------|-----------------------|-------------------|
| <b>GAVIIFORMES</b>      |                       |                   |
| GAVIA ARCTICA           | Rana sp.              | 49                |
| <b>PODICIPEDIFORMES</b> |                       |                   |
| TACHYBAPTUS RUFICOLLIS  | Amphibia              | 49                |
|                         | Triturus alpestris    | 171               |
| PODICEPS CRISTATUS      | Natrix natrix         | 49                |
| <b>PELECANIFORMES</b>   |                       |                   |
| PHALACROCORAX CARBO     | Rana temporaria       | 49, 158           |
| <b>CICONIIFORMES</b>    |                       |                   |
| ARDEIDAE                | Anura                 | 96, 151           |
|                         | "Green frogs"         | 190               |
|                         | Tarentola mauritanica | 153               |
|                         | Lacertidae            | 153               |
| BOTAURUS STELLARIS      | Triturus sp.          | 49                |
|                         | Anura                 | 49                |
|                         | Rana sp.              | 49                |
|                         | "Green frogs"         | 49, 98            |
|                         | Natrix natrix         | 49                |
| IXOBRYCHUS MINUTUS      | Amphibia              | 49                |
|                         | "Green frogs"         | 49                |
| NYCTICORAX NYCTICORAX   | Pleurodeles waltl     | 190               |
|                         | Triturus sp.          | 49                |
|                         | Anura                 | 189, 202          |
|                         | Pelobates cultripes   | 19, 189, 190, 191 |
|                         | Rana sp.              | 49, 189           |
|                         | "Green frogs"         | 49, 189, 202      |
|                         | Lacertidae            | 49, 189           |
|                         | Natrix sp.            | 189               |
|                         | Natrix maura          | 190               |
|                         | Natrix natrix         | 49                |
| ARDEOLA RALLOIDES       | Triturus cristatus    | 189               |
|                         | Pelobates cultripes   | 191               |
|                         | Hyla arborea          | 49, 190           |
|                         | Rana sp.              | 49, 100, 189      |
|                         | "Green frogs"         | 49                |
|                         | Lacertidae            | 49, 189           |

## BULBUCUS IBIS

|                            |                   |
|----------------------------|-------------------|
| Amphibia                   | 138               |
| Pleurodeles waltl          | 138               |
| Hyla arborea               | 19, 191           |
| Rana sp.                   | 19, 49            |
| "Green frogs"              | 19, 110, 138, 166 |
| Reptilia                   | 138               |
| Tarentola mauritanica      | 49, 110           |
| Chalcides chalcides        | 49, 110, 138, 190 |
| Lacertidae                 | 19                |
| Acanthodactylus erythrurus | 138, 189, 191     |
| Lacerta lepida             | 138               |
| Podarcis hispanica         | 166               |
| Psammodromus sp.           | 189               |
| Psammodromus algirus       | 49, 110, 138      |
| Psammodromus hispanicus    | 138               |
| Natrix natrix              | 166               |

## EGRETTA GARZETTA

|                     |              |
|---------------------|--------------|
| Caudata             | 202          |
| Triturus helveticus | 189          |
| Anura               | 38, 202      |
| Pelobates cultripes | 49, 189, 202 |
| Hyla arborea        | 49, 189, 190 |
| Rana sp.            | 49, 100, 189 |
| "Green frogs"       | 19, 189, 202 |
| Reptilia            | 202          |
| Lacertidae          | 49           |

## ARDEA CINEREA

|                     |             |
|---------------------|-------------|
| Triturus sp.        | 149         |
| Pelobates cultripes | 49          |
| Pelobates fuscus    | 49          |
| Bufo bufo           | 90          |
| Rana sp.            | 49, 96, 189 |
| Rana temporaria     | 149         |
| Lacertidae          | 49          |
| Lacerta viridis     | 144         |
| Natrix maura        | 144         |
| Natrix natrix       | 49, 144     |

## ARDEA PURPUREA

|                     |                  |
|---------------------|------------------|
| Pleurodeles waltl   | 2                |
| Triturus sp.        | 49               |
| Pelobates cultripes | 2, 49            |
| Rana sp.            | 49               |
| Lacertidae          | 49               |
| Natrix sp.          | 49               |
| Natrix maura        | 2, 162, 190, 191 |
| Natrix natrix       | 23               |

## CICONIA CICONIA

|                     |         |
|---------------------|---------|
| Caudata             | 164     |
| Pleurodeles waltl   | 124     |
| Discoglossus sp.    | 49, 124 |
| Pelobates cultripes | 124     |
| Bufo sp.            | 49, 164 |





## MERGUS MERGANSER

Rana sp. 74

## FALCONIFORMES

## ELANUS CAERULEUS

Rana sp. 50, 177, 178  
 Lacerta lepida 106

## PERNIS APIVORUS

Bufo bufo 50  
 Hyla arborea 37, 50  
 Rana sp. 77  
 "Green frogs" 22, 50, 85  
 Rana temporaria 50  
 Anguis fragilis 50  
 Lacertidae 50  
 Lacerta lepida 37, 50, 77  
 Lacerta schreiberi 37, 50  
 Lacerta viridis 77  
 Elaphe longissima 50  
 Natrix sp. 50  
 Natrix natrix 22, 50

## MILVUS MILVUS

Caudata 190, 191  
 Pleurodeles waltl 65  
 Triturus marmoratus 190  
 Pelobates cultripes 50, 86  
 Bufo sp. 50  
 Bufo bufo 50, 65, 86, 203  
 Bufo calamita 86  
 Rana sp. 50  
 "Green frogs" 86  
 Rana temporaria 50, 58, 203  
 Testudo graeca 65  
 Blanus cinereus 65  
 Anguis fragilis 50, 86, 203  
 Chalcides bedriagai 25  
 Lacertidae 25, 50  
 Lacerta lepida 7, 25, 50, 65,  
 85, 86, 154, 190,  
 191  
 Podarcis hispanica 25  
 Psammodromus sp. 50  
 Psammodromus algirus 25  
 Psammodromus hispanicus 86  
 Colubridae 25, 190, 191  
 Elaphe scalaris 50, 65, 86  
 Macroprotodon cucullatus 50, 86  
 Malpolon monspessulanus 50, 65, 85, 86  
 Natrix sp. 50  
 Natrix maura 25, 65, 86, 191  
 Natrix natrix 65, 86

## MILVUS MIGRANS

Pleurodeles waltl 50, 61, 190

|                    |                                   |   |
|--------------------|-----------------------------------|---|
|                    | <i>Pelobates cultripes</i>        | 11, 18, 50, 61,<br>80, 86, 190, 191                 |
|                    | <i>Bufo</i> sp.                   | 50  |
|                    | <i>Bufo bufo</i>                  | 5, 7, 80, 86  |
|                    | <i>Rana</i> sp.                   | 50  |
|                    | "Green frogs"                     | 61, 80, 191   |
|                    | <i>Mauremys caspica</i>           | 61, 80  |
|                    | Lacertidae                        | 50  |
|                    | <i>Lacerta lepida</i>             | 9, 11, 50, 61,<br>80, 85, 86, 152,<br>154, 190, 191 |
|                    | Colubridae                        | 191   |
|                    | <i>Coronella girondica</i>        | 11  |
|                    | <i>Elaphe scalaris</i>            | 80  |
|                    | <i>Malpolon monspessulanus</i>    | 85, 190, 191  |
|                    | <i>Natrix</i> sp.                 | 50  |
|                    | <i>Natrix maura</i>               | 61, 80, 190, 191                                    |
|                    | <i>Natrix natrix</i>              | 80  |
| ACCIPITER GENTILIS | <i>Amphibia</i>                   | 75  |
|                    | Lacertidae                        | 85  |
|                    | <i>Lacerta lepida</i>             | 6, 7, 85, 86, 143                                   |
|                    | <i>Psammodromus algirus</i>       | 6   |
| ACCIPITER NISUS    |                                   |   |
|                    | Lacertidae                        | 50  |
| BUTEO BUTEO        | <i>Amphibia</i>                   | 185   |
|                    | <i>Salamandra salamandra</i>      | 33  |
|                    | <i>Pleurodeles waltl</i>          | 191   |
|                    | <i>Anura</i>                      | 33, 131   |
|                    | <i>Discoglossus</i> sp.           | 33  |
|                    | <i>Pelobates cultripes</i>        | 50, 86  |
|                    | <i>Bufo</i> sp.                   | 50  |
|                    | <i>Bufo bufo</i>                  | 5, 33, 77, 85, 86                                   |
|                    | <i>Bufo calamita</i>              | 86  |
|                    | <i>Rana</i> sp.                   | 50  |
|                    | "Green frogs"                     | 21, 86, 191   |
|                    | <i>Rana temporaria</i>            | 134   |
|                    | <i>Blanus cinereus</i>            | 86, 191   |
|                    | Gekkonidae                        | 97  |
|                    | <i>Anguis fragilis</i>            | 33, 50, 131   |
|                    | Lacertidae                        | 33, 50, 185   |
|                    | <i>Acanthodactylus erythrurus</i> | 89  |
|                    | <i>Lacerta lepida</i>             | 6, 7, 18, 33, 85,<br>86, 97, 143, 154,<br>191, 197  |
|                    | <i>Lacerta schreiberi</i>         | 7   |
|                    | <i>Podarcis</i> sp.               | 33  |
|                    | <i>Podarcis hispanica</i>         | 191   |
|                    | <i>Podarcis muralis</i>           | 33  |
|                    | <i>Psammodromus</i> sp.           | 89  |
|                    | <i>Psammodromus algirus</i>       | 6   |
|                    | <i>Psammodromus hispanicus</i>    | 86  |

|                       |                         |  |
|-----------------------|-------------------------|--|
|                       | Colubridae              | 6, 7, 86, 97, 185                                    |
|                       | Coluber hippocrepis     | 85, 86   |
|                       | Coronella girondica     | 6, 33  |
|                       | Elaphe scalaris         | 50, 86, 154  |
|                       | Malpolon monspessulanus | 6, 33, 86  |
|                       | Natrix maura            | 85, 97, 191  |
|                       | Natrix natrix           | 24, 50, 97   |
|                       | Vipera sp.              | 50   |
| HIERAETUS PENNATUS    |                         |  |
|                       | Bufo bufo               | 5  |
|                       | Lacerta lepida          | 7, 12, 18, 50,<br>85, 86, 119, 145,<br>154, 191, 197 |
| HIERAETUS FASCIATUS   |                         |  |
|                       | Bufo bufo               | 173  |
|                       | Lacertidae              | 50, 173  |
|                       | Lacerta lepida          | 6, 8, 28, 41, 50,<br>55, 84, 123, 159,<br>173, 195   |
|                       | Lacerta viridis         | 28, 41, 173, 195                                     |
|                       | Colubridae              | 41   |
|                       | Malpolon monspessulanus | 6, 123   |
| AQUILA HELIACA        |                         |  |
|                       | Reptilia                | 62   |
|                       | Testudo graeca          | 50   |
|                       | Lacerta lepida          | 50, 62, 86, 190,<br>191                              |
|                       | Colubridae              | 190, 191   |
|                       | Coluber hippocrepis     | 62   |
|                       | Elaphe scalaris         | 50, 62   |
|                       | Malpolon monspessulanus | 50, 62, 190, 191                                     |
|                       | Natrix maura            | 62   |
| AQUILA CHRYSAETOS     |                         |  |
|                       | Amphibia                | 43, 44   |
|                       | Reptilia                | 43, 64, 107  |
|                       | Testudo sp.             | 50   |
|                       | Testudo graeca          | 64   |
|                       | Lacertidae              | 50, 182  |
|                       | Lacerta lepida          | 18, 64, 77, 78,<br>86, 123                           |
|                       | Lacerta viridis         | 13   |
|                       | Colubridae              | 13, 43, 44, 64,<br>78, 123, 182                      |
|                       | Elaphe scalaris         | 18, 138  |
|                       | Malpolon monspessulanus | 64   |
|                       | Vipera sp.              | 50   |
| NEOPHRON PERCNOPTERUS |                         |  |
|                       | Amphibia                | 181  |
|                       | Pleurodeles waltl       | 17, 86   |
|                       | Pelobates cultripes     | 191  |
|                       | Bufo sp.                | 50, 73   |
|                       | Rana sp.                | 50   |
|                       | Reptilia                | 181  |

|                    |                         |  |
|--------------------|-------------------------|--|
|                    | Testudines              | 50   |
|                    | Testudo sp.             | 17   |
|                    | Emys orbicularis        | 17   |
|                    | Mauremys caspica        | 17, 86, 154                                  |
|                    | Lacertidae              | 50   |
|                    | Lacerta lepida          | 6, 17, 50, 73,<br>85, 86, 127, 154<br>163    |
|                    | Lacerta viridis         | 50   |
|                    | Colubridae              | 6, 17, 50                                    |
|                    | Elaphe scalaris         | 17, 73, 127                                  |
|                    | Malpolon monspessulanus | 17, 50, 73, 85,<br>127, 163                  |
|                    | Natrix maura            | 17, 154                                      |
|                    | Natrix natrix           | 17, 86, 127                                  |
| GYPAETUS BARBATUS  |                         |  |
|                    | Testudo sp.             | 50, 113                                      |
|                    | Testudo graeca          | 50   |
|                    | Testudo hermanni        | 50   |
|                    | Lacerta lepida          | 113  |
|                    | Colubridae              | 113  |
| AEGYPIUS MONACHUS  |                         |  |
|                    | Testudines              | 50   |
|                    | Testudo sp.             | 50   |
|                    | Lacertidae              | 50   |
|                    | Lacerta lepida          | 111  |
|                    | Psammodromus algirus    | 111  |
|                    | Malpolon monspessulanus | 111  |
| CIRCAETUS GALLICUS |                         |  |
|                    | Amphibia                | 183  |
|                    | Pelobates cultripipes   | 50, 86                                       |
|                    | Bufo bufo               | 4, 5, 50                                     |
|                    | Bufo viridis            | 50   |
|                    | Rana sp.                | 4  |
|                    | "Green frogs"           | 50   |
|                    | Anguis fragilis         | 42, 50, 183                                  |
|                    | Chalcides chalcides     | 86   |
|                    | Lacertidae              | 4  |
|                    | Lacerta agilis          | 50   |
|                    | Lacerta lepida          | 4, 18, 50, 86,<br>120, 183, 190,<br>191, 197 |
|                    | Lacerta viridis         | 42, 50, 183                                  |
|                    | Podarcis muralis        | 183  |
|                    | Psammodromus algirus    | 4, 86  |
|                    | Colubridae              | 197  |
|                    | Coluber hippocrepis     | 4, 86  |
|                    | Coluber viridiflavus    | 50, 183                                      |
|                    | Coronella sp.           | 183  |
|                    | Coronella austriaca     | 50, 86                                       |
|                    | Coronella girondica     | 4  |

|                    |                                   |   |
|--------------------|-----------------------------------|---|
|                    | <i>Elaphe longissima</i>          | 42, 50, 183                             |
|                    | <i>Elaphe scalaris</i>            | 4, 6, 86, 120,<br>152, 154, 190,<br>191 |
|                    | <i>Malpolon monspessulanus</i>    | 4, 50, 85, 86,<br>120, 183, 190,<br>191 |
|                    | <i>Natrix</i> sp.                 | 18                                      |
|                    | <i>Natrix maura</i>               | 4, 50, 183, 190,<br>191                 |
|                    | <i>Natrix natrix</i>              | 4, 42, 50, 183                          |
|                    | <i>Vipera aspis</i>               | 42, 50                                  |
|                    | <i>Vipera latastei</i>            | 4                                       |
|                    | <i>Vipera berus</i>               | 50                                      |
| CIRCUS CYANEUS     |                                   |   |
|                    | Lacertide                         | 169                                     |
|                    | <i>Lacerta vivipara</i>           | 157                                     |
| CIRCUS PYGARGUS    |                                   |   |
|                    | Anura                             | 169                                     |
|                    | <i>Bufo bufo</i>                  | 50                                      |
|                    | <i>Rana</i> sp.                   | 155                                     |
|                    | "Green frogs"                     | 50                                      |
|                    | <i>Rana temporaria</i>            | 50                                      |
|                    | Reptilia                          | 114                                     |
|                    | <i>Blanus cinereus</i>            | 114                                     |
|                    | <i>Tarentola mauritanica</i>      | 86                                      |
|                    | <i>Chalcides bedriagai</i>        | 114                                     |
|                    | <i>Chalcides chalcides</i>        | 114, 154                                |
|                    | Lacertidae                        | 18, 114, 155, 169                       |
|                    | <i>Acanthodactylus erythrurus</i> | 154, 191                                |
|                    | <i>Lacerta lepida</i>             | 114, 154, 155                           |
|                    | <i>Psammodromus algirus</i>       | 50, 114                                 |
|                    | <i>Psammodromus hispanicus</i>    | 86                                      |
|                    | Colubridae                        | 18, 114                                 |
|                    | <i>Natrix natrix</i>              | 50                                      |
|                    | <i>Vipera berus</i>               | 50                                      |
| CIRCUS AERUGINOSUS |                                   |   |
|                    | Amphibia                          | 185                                     |
|                    | Anura                             | 169                                     |
|                    | <i>Rana</i> sp.                   | 50                                      |
|                    | "Green frogs"                     | 190                                     |
|                    | Lacertidae                        | 169                                     |
|                    | Colubridae                        | 169                                     |
|                    | <i>Natrix maura</i>               | 50                                      |
|                    | <i>Natrix natrix</i>              | 50                                      |
| FALCO PEREGRINUS   |                                   |   |
|                    | <i>Bufo</i> sp.                   | 50                                      |
|                    | <i>Rana</i> sp.                   | 50                                      |
|                    | Lacertidae                        | 50                                      |
|                    | <i>Lacerta viridis</i>            | 165                                     |

|                     |                            |                               |
|---------------------|----------------------------|-------------------------------|
| FALCO SUBBUTEO      |                            |                               |
|                     | Reptilia                   | 50                            |
| FALCO ELEONORAE     |                            |                               |
|                     | Lacertidae                 | 50                            |
|                     | Podarcis lilfordi          | 10, 167                       |
| FALCO COLUMBARIUS   |                            |                               |
|                     | Rana temporaria            | 160                           |
|                     | Lacerta vivipara           | 20                            |
| FALCO NAUMANNI      |                            |                               |
|                     | Pelodytes punctatus        | 50, 83                        |
|                     | Blanus cinereus            | 50, 82, 83                    |
|                     | Tarentola mauritanica      | 50, 82, 191                   |
|                     | Chalcides sp.              | 50, 83                        |
|                     | Chalcides chalcides        | 82, 83                        |
|                     | Lacertidae                 | 18, 26, 83                    |
|                     | Lacerta lepida             | 50, 82, 83                    |
|                     | Podarcis hispanica         | 50, 83                        |
|                     | Psammodromus sp.           | 18, 50, 85                    |
|                     | Psammodromus algirus       | 82, 83                        |
|                     | Psammodromus hispanicus    | 83                            |
|                     | Malpolon monspessulanus    | 82                            |
|                     | Natrix maura               | 50, 83                        |
| FALCO TINNUNCULUS   |                            |                               |
|                     | Amphibia                   | 185                           |
|                     | Anura                      | 29, 191                       |
|                     | Rana temporaria            | 50, 59, 188, 200              |
|                     | Reptilia                   | 180                           |
|                     | Blanus cinereus            | 191                           |
|                     | Anguis fragilis            | 50, 59, 77                    |
|                     | Lacertidae                 | 6, 7, 29, 77, 85,<br>172, 185 |
|                     | Acanthodactylus erythrurus | 85, 191                       |
|                     | Lacerta lepida             | 7, 18, 191                    |
|                     | Lacerta vivipara           | 50, 59, 200, 205              |
|                     | Podarcis hispanica         | 148, 191                      |
|                     | Psammodromus algirus       | 6, 18, 191                    |
|                     | Psammodromus hispanicus    | 6, 191                        |
|                     | Colubridae                 | 77, 185                       |
| <b>GALLIFORMES</b>  |                            |                               |
| PHASIANUS COLCHICUS |                            |                               |
|                     | Rana sp.                   | 50                            |
|                     | Lacerta agilis             | 50                            |
|                     | Natrix natrix              | 50                            |
|                     | Vipera berus               | 50                            |
| GALLUS GALLUS       |                            |                               |
|                     | Salamandra salamandra      | 116                           |
| <b>GRUIFORMES</b>   |                            |                               |
| GRUS GRUS           |                            |                               |
|                     | Rana sp.                   | 50                            |
|                     | Anguis fragilis            | 50                            |
|                     | Lacertidae                 | 50                            |

|                        |                      |              |
|------------------------|----------------------|--------------|
|                        | Colubridae           | 50           |
| RALLUS AQUATICUS       |                      |              |
|                        | Triturus sp.         | 50           |
|                        | Bufo sp.             | 50           |
|                        | Rana sp.             | 50           |
| PORZANA PUSILLA        |                      |              |
|                        | Lacertidae           | 50           |
| GALLINULA CHLOROPUS    |                      |              |
|                        | Amphibia             | 50           |
| PORPHYRIO PORPHYRIO    |                      |              |
|                        | Natrix maura         | 50           |
| FULICA ATRA            |                      |              |
|                        | Rana sp.             | 50           |
| OTIS TARDA             |                      |              |
|                        | Rana sp.             | 50           |
|                        | Lacertidae           | 50           |
| OTIS TETRAX            |                      |              |
|                        | Rana sp.             | 50           |
| <b>CHARADRIIFORMES</b> |                      |              |
| SCOLOPAX RUSTICOLA     |                      |              |
|                        | Chalcides chalcides  | 60           |
| HIMANTOPUS HIMANTOPUS  |                      |              |
|                        | Amphibia             | 51           |
|                        | Pleurodeles waltl    | 199          |
| BURHINUS OEDICNEMUS    |                      |              |
|                        | Amphibia             | 1, 51        |
|                        | Pelobates fuscus     | 51           |
|                        | Psammodromus algirus | 191          |
|                        | Colubridae           | 191          |
| TRINGA NEBULARIA       |                      |              |
|                        | Rana sp.             | 179          |
| LARUS RIDIBUNDUS       |                      |              |
|                        | Pelobates cultripes  | 51           |
|                        | Reptilia             | 51           |
| LARUS CANUS            |                      |              |
|                        | Rana temporaria      | 161          |
| LARUS ARGENTATUS       |                      |              |
|                        | Amphibia             | 51           |
|                        | Bufo bufo            | 103          |
|                        | Rana temporaria      | 103          |
|                        | Reptilia             | 51           |
|                        | Lacertidae           | 132          |
|                        | Vipera berus         | 130          |
| LARUS MARINUS          |                      |              |
|                        | Bufo bufo            | 103          |
|                        | Rana temporaria      | 103          |
| CHLIDONIAS HYBRIDA     |                      |              |
|                        | Rana sp.             | 190          |
| GELOCHELIDON NILOTICA  |                      |              |
|                        | Pelobates cultripes  | 48, 192, 193 |
|                        | "Green frogs"        | 48, 126      |
|                        | Lacerta lepida       | 193          |

## CUCULIFORMES

## CLAMATOR GLANDARIUS

Lacertidae 52

## CUCULUS CANORUS

Bufo bufo 140

Rana temporaria 140

## STRIGIFORMES

## TYTO ALBA

Amphibia 3, 32, 34, 76,  
95, 176

Anura 31, 57

Discoglossus sp. 32, 47, 52, 196

Pelobates cultripes 32, 36, 52, 108,  
129, 137, 152,  
190, 191

Pelodytes punctatus 52, 108, 137

Bufo bufo 52, 92, 185

Rana sp. 15, 32, 95

Rana dalmatina 52

"Green frogs" 35, 36, 47, 52,  
95, 108, 137, 185

Rana temporaria 52, 66, 92, 174,  
185, 204

Reptilia 176

Chamaeleo chamaeleon 52

Hemidactylus turcicus 137, 198

Tarentola mauritanica 52, 137, 146,  
194, 198

Anguis fragilis 52

Lacertidae 32, 57, 76, 92,  
185

Acanthodactylus erythrurus 35, 52, 190, 191

Lacerta lepida 52, 108, 109

Lacerta viridis 52

Podarcis hispanica 35, 52, 108, 137,  
198

Podarcis muralis 52

Psammodromus sp. 35, 108

Psammodromus algirus 52, 190, 191

Psammodromus hispanicus 137, 198

Colubridae 185

Natrix natrix 52, 185

## BUBO BUBO

Amphibia 14, 54, 184

Anura 27, 147

Pelobates cultripes 52, 112, 156

Bufo sp. 147

Bufo bufo 5, 27, 52, 112

"Green frogs" 52, 112

Rana temporaria 52, 69, 72

Reptilia 14, 54

Mauremys caspica 52, 112



|               |                            |                               |
|---------------|----------------------------|-------------------------------|
|               | Tarentola mauritanica      | 147                           |
|               | Anguis fragilis            | 52                            |
|               | Lacertidae                 | 27, 112, 184                  |
|               | Lacerta agilis             | 69                            |
|               | Lacerta lepida             | 27, 52, 112, 147,<br>156, 198 |
|               | Lacerta viridis            | 147                           |
|               | Podarcis hispanica         | 52, 112                       |
|               | Psammodromus algirus       | 52, 112                       |
|               | Colubridae                 | 6, 112, 184                   |
|               | Coluber viridiflavus       | 99                            |
|               | Natrix natrix              | 52                            |
| ASIO OTUS     |                            |                               |
|               | Amphibia                   | 185                           |
|               | Pelobates fuscus           | 52                            |
|               | Rana sp.                   | 175                           |
|               | Rana temporaria            | 52, 94, 175, 204              |
|               | Tarentola mauritanica      | 196                           |
|               | Anguis fragilis            | 52                            |
|               | Lacertidae                 | 46, 52, 175, 196,<br>197      |
| ASIO FLAMMEUS |                            |                               |
|               | Rana temporaria            | 52, 91, 93                    |
|               | Lacerta vivipara           | 52                            |
|               | Podarcis muralis           | 52                            |
| OTUS SCOPS    |                            |                               |
|               | Hyla arborea               | 52                            |
|               | Lacertidae                 | 52                            |
|               | Psammodromus algirus       | 6                             |
| ATHENE NOCTUA |                            |                               |
|               | Amphibia                   | 128, 136, 185                 |
|               | Pleurodeles waltl          | 52, 135, 136                  |
|               | Triturus boscai            | 52, 141                       |
|               | Discoglossus sp.           | 52, 136                       |
|               | Pelobates cultripes        | 135, 136, 190,<br>191         |
|               | Pelobates fuscus           | 52                            |
|               | Hyla arborea               | 52                            |
|               | Rana sp.                   | 128                           |
|               | "Green frogs"              | 52, 136                       |
|               | Rana temporaria            | 52, 136, 185                  |
|               | Reptilia                   | 135, 136, 185                 |
|               | Anguis fragilis            | 52                            |
|               | Chalcides bedriagai        | 52, 136                       |
|               | Lacertidae                 | 63, 136                       |
|               | Acanthodactylus erythrurus | 52, 191                       |
|               | Lacerta agilis             | 52                            |
|               | Lacerta vivipara           | 52                            |
|               | Podarcis sp.               | 136                           |
|               | Podarcis hispanica         | 136                           |
|               | Podarcis muralis           | 52                            |
|               | Psammodromus sp.           | 136                           |
|               | Psammodromus algirus       | 52, 136                       |



## PICIFORMES

JYNX TORQUILLA

Rana temporaria 52

PICUS VIRIDIS

Vipera berus 52

## PASSERIFORMES

MOTACILLA ALBA

Rana temporaria 187

LANIUS SENATOR

Amphibia 88

Lacertidae 88

LANIUS EXCUBITOR

Rana temporaria 102

Blanus cinereus 190, 191

Chalcides bedriagai 190, 191

Lacertidae 190

Acanthodactylus erythrurus 190

Podarcis hispanica 191

Psammodromus algirus 190, 191

LANIUS COLLURIO

Anura 125

Rana sp. 125

Lacertidae 125

SAXICOLA TORQUATA

Lacertidae 191

MONTICOLA SAXATILIS

Rana sp. 53

Lacertidae 53, 133

MONTICOLA SOLITARIUS

Anura 53

Gekkonidae 53

Lacertidae 53

Colubridae 53

ERITHACUS RUBECULA

Lacerta agilis 104

Lacerta vivipara 104

TURDUS TORQUATUS

Salamandra salamandra 53

Lacertidae 121

Lacerta vivipara 53, 121

TURDUS MERULA

Triturus sp. 53

Anura 53

Anguis fragilis 81

Lacertidae 53

Colubridae 170

Natrix sp. 53

TURDUS PHILOMELOS

Anguis fragilis 52, 117

Lacertidae 53

Lacerta vivipara 39, 53

|                     |                         |          |
|---------------------|-------------------------|----------|
| STURNUS UNICOLOR    | Pelobates cultripes     | 190, 191 |
|                     | Colubridae              | 190      |
| GARRULUS GLANDARIUS | Reptilia                | 115, 168 |
|                     | Natrix sp.              | 115      |
| PICA PICA           | Amphibia                | 115      |
|                     | Reptilia                | 115      |
|                     | Podarcis hispanica      | 70       |
|                     | Malpolon monspessulanus | 70       |
| CORVUS MONEDULA     | Amphibia                | 115      |
| CORVUS CORONE       | Amphibia                | 115, 122 |
|                     | Anura                   | 122      |
|                     | Reptilia                | 122      |
|                     | Lacertidae              | 122      |
| CORVUS CORAX        | Amphibia                | 115      |
|                     | Pelobates cultripes     | 191      |
|                     | Bufo sp.                | 56       |
|                     | Rana sp.                | 56, 191  |
|                     | Reptilia                | 115      |
|                     | Colubridae              | 56       |

#### AMPHIBIANS AND REPTILES EATEN BY BIRDS

| <u>PREY</u>    | <u>PREDATOR</u>        | <u>REFERENCE</u> |
|----------------|------------------------|------------------|
| ** AMPHIBIA ** | Tachybaptus ruficollis | 49               |
|                | Ixobrychus minutus     | 49               |
|                | Bulbucus ibis          | 138              |
|                | Anas platyrynchos      | 49               |
|                | Anas strepera          | 49               |
|                | Anas acuta             | 49               |
|                | Anas clypeata          | 49               |
|                | Netta rufina           | 49               |
|                | Accipiter gentilis     | 75               |
|                | Buteo buteo            | 185              |
|                | Aquila chrysaetos      | 43, 44           |
|                | Neophron percnopterus  | 181              |
|                | Circus gallicus        | 183              |
|                | Circus aeruginosus     | 185              |
|                | Falco tinnunculus      | 185              |
|                | Gallinula chloropus    | 50               |
|                | Himantopus himantopus  | 51               |

|                       |                        |                           |
|-----------------------|------------------------|---------------------------|
|                       | Burhinus oedicephalus  | 1, 51                     |
|                       | Larus argentatus       | 51                        |
|                       | Tyto alba              | 3, 32, 34, 76,<br>95, 176 |
|                       | Bubo bubo              | 14, 54, 184               |
|                       | Asio otus              | 185                       |
|                       | Athene noctua          | 128, 136, 185             |
|                       | Alcedo atthis          | 101                       |
|                       | Lanius senator         | 89                        |
|                       | Pica pica              | 115                       |
|                       | Corvus monedula        | 115                       |
|                       | Corvus corone          | 115, 122                  |
|                       | Corvus corax           | 115                       |
| CAUDATA               |                        |                           |
|                       | Egretta garzetta       | 202                       |
|                       | Ciconia ciconia        | 164                       |
|                       | Ciconia nigra          | 49                        |
|                       | Milvus milvus          | 190, 191                  |
| SALAMANDRA SALAMANDRA |                        |                           |
|                       | Buteo buteo            | 33                        |
|                       | Gallus gallus          | 116                       |
|                       | Strix aluco            | 148                       |
|                       | Turdus torquatus       | 53                        |
| PLEURODELES WALTJ     |                        |                           |
|                       | Nycticorax nycticorax  | 190                       |
|                       | Bulbucus ibis          | 138                       |
|                       | Ardea purpurea         | 2                         |
|                       | Ciconia ciconia        | 124                       |
|                       | Ciconia nigra          | 71, 87                    |
|                       | Milvus milvus          | 65                        |
|                       | Milvus migrans         | 50, 61, 190               |
|                       | Buteo buteo            | 191                       |
|                       | Neophron percnopterus  | 17, 86                    |
|                       | Himantopus himantopus  | 199                       |
|                       | Athene noctua          | 52, 135, 136              |
| TRITURUS SP.          |                        |                           |
|                       | Botaurus stellaris     | 49                        |
|                       | Nycticorax nycticorax  | 49                        |
|                       | Ardea cinerea          | 149                       |
|                       | Ardea purpurea         | 49                        |
|                       | Rallus aquaticus       | 50                        |
|                       | Turdus merula          | 53                        |
| TRITURUS ALPESTRIS    |                        |                           |
|                       | Tachybaptus ruficollis | 171                       |
|                       | Strix aluco            | 16                        |
| TRITURUS BOSCAI       |                        |                           |
|                       | Athene noctua          | 52, 141                   |
| TRITURUS MARMORATUS   |                        |                           |
|                       | Ciconia nigra          | 71                        |
|                       | Milvus milvus          | 190                       |
| TRITURUS HELVETICUS   |                        |                           |
|                       | Egretta garzetta       | 189                       |

## TRITURUS CRISTATUS

|                   |     |
|-------------------|-----|
| Ardeola ralloides | 189 |
| Strix aluco       | 52  |

## ANURA

|                       |          |
|-----------------------|----------|
| Ardeidae              | 96, 151  |
| Botaurus stellaris    | 49       |
| Nycticorax nycticorax | 189, 202 |
| Egretta garzetta      | 38, 202  |
| Buteo buteo           | 33, 131  |
| Circus pygargus       | 169      |
| Circus aeruginosus    | 169      |
| Falco tinnunculus     | 29, 191  |
| Tyto alba             | 31, 57   |
| Bubo bubo             | 27, 147  |
| Strix aluco           | 16, 175  |
| Alcedo atthis         | 118      |
| Upupa epops           | 52       |
| Lanius collurio       | 125      |
| Monticola solitarius  | 53       |
| Turdus merula         | 53       |
| Corvus corone         | 122      |

## DISCOGLOSSIDAE

## ALYTES OBSTETRICANS

|             |         |
|-------------|---------|
| Strix aluco | 52, 201 |
|-------------|---------|

## DISCOGLOSSUS SP.

|                 |                 |
|-----------------|-----------------|
| Ciconia ciconia | 49, 124         |
| Ciconia nigra   | 71              |
| Buteo buteo     | 33              |
| Tyto alba       | 32, 47, 52, 196 |
| Athene noctua   | 52, 136         |
| Strix aluco     | 201             |

## PELOBATIDAE

## PELOBATES CULTRIPES

|                       |                                     |
|-----------------------|-------------------------------------|
| Nycticorax nycticorax | 19, 189, 190, 191                   |
| Ardeola ralloides     | 191                                 |
| Egretta garzetta      | 49, 189, 202                        |
| Ardea cinerea         | 49                                  |
| Ardea purpurea        | 2, 49                               |
| Ciconia ciconia       | 124                                 |
| Ciconia nigra         | 71, 87                              |
| Milvus milvus         | 50, 86                              |
| Milvus migrans        | 11, 18, 50, 61,<br>80, 86, 190, 191 |
| Buteo buteo           | 50, 86                              |
| Neophron percnopterus | 191                                 |
| Circaetus gallicus    | 50, 86                              |
| Larus ridibundus      | 51                                  |
| Gelochelidon nilotica | 48, 192, 193                        |

|                     |                       |  |
|---------------------|-----------------------|--|
|                     | Tyto alba             | 32, 36, 52, 108,<br>129, 137, 152,<br>190, 191 |
|                     | Bubo bubo             | 52, 112, 156                                   |
|                     | Athene noctua         | 135, 136, 190,<br>191                          |
|                     | Strix aluco           | 52, 129, 201                                   |
|                     | Sturnus unicolor      | 190, 191                                       |
|                     | Corvus corax          | 191  |
| PELOBATES FUSCUS    |                       |  |
|                     | Ardea cinerea         | 49   |
|                     | Burhinus oedicephalus | 51   |
|                     | Asio otus             | 52   |
|                     | Athene noctua         | 52   |
|                     | Strix aluco           | 52   |
| <b>PELODYTIDAE</b>  |                       |  |
| PELODYTES PUNCTATUS |                       |  |
|                     | Falco naumanni        | 50, 83   |
|                     | Tyto alba             | 52, 108, 137                                   |
| <b>BUFONIDAE</b>    |                       |  |
|                     | Ciconia ciconia       | 49, 164  |
|                     | Ciconia nigra         | 73   |
|                     | Cygnus olor           | 49   |
|                     | Milvus milvus         | 50   |
|                     | Milvus migrans        | 50   |
|                     | Buteo buteo           | 50   |
|                     | Neophron percnopterus | 50, 71   |
|                     | Falco peregrinus      | 50   |
|                     | Rallus aquaticus      | 50   |
|                     | Bubo bubo             | 147  |
|                     | Corvus corax          | 56   |
| BUFO BUFO           |                       |  |
|                     | Ardea cinerea         | 90   |
|                     | Ciconia ciconia       | 186  |
|                     | Pernis apivorus       | 50   |
|                     | Milvus milvus         | 50, 65, 86, 203                                |
|                     | Milvus migrans        | 5, 7, 80, 86                                   |
|                     | Buteo buteo           | 5, 33, 77, 85, 86                              |
|                     | Hieraetus pennatus    | 5  |
|                     | Hieraetus fasciatus   | 173  |
|                     | Circaetus gallicus    | 4, 5, 50                                       |
|                     | Circus pygargus       | 50   |
|                     | Larus argentatus      | 103  |
|                     | Larus marinus         | 103  |
|                     | Cuculus canorus       | 140  |
|                     | Tyto alba             | 52, 92, 185                                    |
|                     | Bubo bubo             | 5, 27, 52, 112                                 |
|                     | Strix aluco           | 52   |
| BUFO CALAMITA       |                       |  |
|                     | Milvus milvus         | 86   |
|                     | Buteo buteo           | 86   |

## BUFO VIRIDIS

*Circaetus gallicus* 50

## HYLIDAE

## HYLA ARBOREA (INC. H. MERIDIONALIS)

*Ardeola ralloides* 49, 190  
*Bulbucus ibis* 19, 191  
*Egretta garzetta* 49, 189, 190  
*Pernis apivorus* 37, 50  
*Otus scops* 52  
*Athene noctua* 52  
*Strix aluco* 52

## RANIDAE

*Gavia arctica* 49  
*Botaurus stellaris* 49  
*Nycticorax nycticorax* 49, 189  
*Ardeola ralloides* 49, 100, 189  
*Bulbucus ibis* 19, 49  
*Egretta garzetta* 49, 100, 189  
*Ardea cinerea* 49, 96, 189  
*Ardea purpurea* 49  
*Ciconia ciconia* 45, 49, 150, 164  
*Ciconia nigra* 49  
*Cygnus olor* 49  
*Aythya ferina* 49  
*Mergus albellus* 49  
*Mergus merganser* 74  
*Elanus caeruleus* 50, 177, 178  
*Pernis apivorus* 77  
*Milvus milvus* 50  
*Milvus migrans* 50  
*Buteo buteo* 50  
*Neophron percnopterus* 50  
*Circaetus gallicus* 4  
*Circus pygargus* 155  
*Circus aeruginosus* 50  
*Falco peregrinus* 50  
*Phasianus colchicus* 50  
*Grus grus* 50  
*Rallus aquaticus* 50  
*Fulica atra* 50  
*Otis tarda* 50  
*Otis tetrax* 50  
*Tringa nebularia* 179  
*Chlidonias hybrida* 190  
*Tyto alba* 15, 32, 95  
*Asio otus* 175  
*Athene noctua* 128  
*Alcedo atthis* 52, 118  
*Coracias garrulus* 52  
*Lanius collurio* 125  
*Monticola saxatilis* 53



|                 |                              |                                      |
|-----------------|------------------------------|--------------------------------------|
|                 | <i>Corvus corax</i>          | 56, 191                              |
| RANA DALMATINA  | <i>Tyto alba</i>             | 52                                   |
|                 | <i>Strix aluco</i>           | 105                                  |
| "GREEN FROGS"   | <i>Ardeidae</i>              | 190                                  |
|                 | <i>Botaurus stellaris</i>    | 49, 98                               |
|                 | <i>Ixobrychus minutus</i>    | 49                                   |
|                 | <i>Nycticorax nycticorax</i> | 49, 189, 202                         |
|                 | <i>Ardeola ralloides</i>     | 49                                   |
|                 | <i>Bulbucus ibis</i>         | 19, 110, 138, 166                    |
|                 | <i>Egretta garzetta</i>      | 19, 189, 202                         |
|                 | <i>Ciconia ciconia</i>       | 49, 124                              |
|                 | <i>Ciconia nigra</i>         | 73                                   |
|                 | <i>Pernis apivorus</i>       | 22, 50, 85                           |
|                 | <i>Milvus milvus</i>         | 86                                   |
|                 | <i>Milvus migrans</i>        | 61, 80, 191                          |
|                 | <i>Buteo buteo</i>           | 21, 86, 191                          |
|                 | <i>Circaetus gallicus</i>    | 50                                   |
|                 | <i>Circus pygargus</i>       | 50                                   |
|                 | <i>Circus aeruginosus</i>    | 190                                  |
|                 | <i>Gelochelidon nilotica</i> | 48, 126                              |
|                 | <i>Tyto alba</i>             | 35, 36, 47, 52,<br>95, 108, 137, 185 |
|                 | <i>Bubo bubo</i>             | 52, 112                              |
|                 | <i>Athene noctua</i>         | 52, 136                              |
|                 | <i>Strix aluco</i>           | 52, 175                              |
|                 | <i>Alcedo atthis</i>         | 52, 118                              |
|                 | <i>Coracias garrulus</i>     | 52                                   |
| RANA TEMPORARIA | <i>Phalacrocorax carbo</i>   | 49, 158                              |
|                 | <i>Ardea cinerea</i>         | 149                                  |
|                 | <i>Ciconia ciconia</i>       | 49, 79                               |
|                 | <i>Pernis apivorus</i>       | 50                                   |
|                 | <i>Milvus milvus</i>         | 50, 58, 203                          |
|                 | <i>Buteo buteo</i>           | 134                                  |
|                 | <i>Circus pygargus</i>       | 50                                   |
|                 | <i>Falco columbarius</i>     | 160                                  |
|                 | <i>Falco tinnunculus</i>     | 50, 59, 188, 202                     |
|                 | <i>Larus canus</i>           | 161                                  |
|                 | <i>Larus argentatus</i>      | 103                                  |
|                 | <i>Larus marinus</i>         | 103                                  |
|                 | <i>Cuculus canorus</i>       | 140                                  |
|                 | <i>Tyto alba</i>             | 52, 66, 92, 174,<br>185, 204         |
|                 | <i>Bubo bubo</i>             | 52, 69, 70                           |
|                 | <i>Asio otus</i>             | 52, 94, 175, 204                     |
|                 | <i>Asio flammeus</i>         | 52, 91, 93                           |
|                 | <i>Athene noctua</i>         | 52, 136, 185                         |
|                 | <i>Strix aluco</i>           | 52, 175, 185,<br>201, 204            |
|                 | <i>Coracias garrulus</i>     | 52                                   |
|                 | <i>Jynx torquilla</i>        | 52                                   |

|                         |                       |               |
|-------------------------|-----------------------|---------------|
|                         | Motacilla alba        | 187           |
|                         | Lanius excubitor      | 102           |
| <b>** REPTILIA **</b>   |                       |               |
|                         | Bulbucus ibis         | 138           |
|                         | Egretta garzetta      | 202           |
|                         | Aquila heliaca        | 62            |
|                         | Aquila chrysaetos     | 43, 64, 107   |
|                         | Neophron percnopterus | 181           |
|                         | Circus pygargus       | 114           |
|                         | Falco subbuteo        | 50            |
|                         | Falco tinnunculus     | 180           |
|                         | Larus ridibundus      | 51            |
|                         | Larus argentatus      | 51            |
|                         | Tyto alba             | 176           |
|                         | Bubo bubo             | 14, 54        |
|                         | Athene noctua         | 135, 136, 185 |
|                         | Strix aluco           | 185           |
|                         | Garrulus glandarius   | 115, 168      |
|                         | Pica pica             | 115           |
|                         | Corvus corone         | 122           |
|                         | Corvus corax          | 115           |
| <b>TESTUDINES</b>       |                       |               |
|                         | Neophron percnopterus | 50            |
|                         | Aegypius monachus     | 50            |
| <b>TESTUDINIDAE</b>     |                       |               |
|                         | Aquila chrysaetos     | 50            |
|                         | Neophron percnopterus | 17            |
|                         | Gypaetus barbatus     | 50, 113       |
|                         | Aegypius monachus     | 50            |
| <b>TESTUDO GRAECA</b>   |                       |               |
|                         | Milvus milvus         | 65            |
|                         | Aquila heliaca        | 50            |
|                         | Aquila chrysaetos     | 64            |
|                         | Gypaetus barbatus     | 50            |
| <b>TESTUDO HERMANNI</b> |                       |               |
|                         | Gypaetus barbatus     | 50            |
| <b>EMYDIDAE</b>         |                       |               |
|                         | Ciconia ciconia       | 124           |
| <b>EMYS ORBICULARIS</b> |                       |               |
|                         | Neophron percnopterus | 17            |
| <b>MAUREMYS CASPICA</b> |                       |               |
|                         | Ciconia ciconia       | 49            |
|                         | Milvus migrans        | 61, 80        |
|                         | Neophron percnopterus | 17, 86, 154   |
|                         | Bubo bubo             | 52, 112       |
| <b>AMPHISBAENIA</b>     |                       |               |
| <b>BLANUS CINEREUS</b>  |                       |               |
|                         | Milvus milvus         | 65            |
|                         | Buteo buteo           | 86, 191       |
|                         | Circus pygargus       | 114           |

|                              |                      |                          |
|------------------------------|----------------------|--------------------------|
|                              | Falco naumanni       | 50, 82, 83               |
|                              | Falco tinnunculus    | 191                      |
|                              | Lanius excubitor     | 190, 191                 |
| <b>LACERTILIA</b>            |                      |                          |
| <b>CHAMAELEONTIDAE</b>       |                      |                          |
| <b>CHAMAELEO CHAMAELEON</b>  |                      |                          |
|                              | Tyto alba            | 52                       |
| <b>GEKKONIDAE</b>            |                      |                          |
|                              | Buteo buteo          | 97                       |
|                              | Strix aluco          | 52                       |
|                              | Monticola solitarius | 53                       |
| <b>HEMIDACTYLUS TURCICUS</b> |                      |                          |
|                              | Tyto alba            | 137, 198                 |
| <b>TARENTOLA MAURITANICA</b> |                      |                          |
|                              | Ardeidae             | 153                      |
|                              | Bulbucus ibis        | 49, 110                  |
|                              | Circus pygargus      | 86                       |
|                              | Falco naumanni       | 50, 82, 191              |
|                              | Tyto alba            | 52, 137, 146, 194<br>198 |
|                              | Bubo bubo            | 147                      |
|                              | Asio otus            | 196                      |
|                              | Strix aluco          | 40, 201                  |
| <b>ANGUIDAE</b>              |                      |                          |
| <b>ANGUIS FRAGILIS</b>       |                      |                          |
|                              | Ciconia ciconia      | 49, 150                  |
|                              | Peris apivorus       | 50                       |
|                              | Milvus milvus        | 50, 86, 203              |
|                              | Buteo buteo          | 33, 50, 131              |
|                              | Ciracetus gallicus   | 42, 50, 183              |
|                              | Falco tinnunculus    | 50, 59, 77               |
|                              | Grus grus            | 50                       |
|                              | Tyto alba            | 52                       |
|                              | Bubo bubo            | 52                       |
|                              | Asio otus            | 52                       |
|                              | Athene noctua        | 52                       |
|                              | Strix aluco          | 52                       |
|                              | Coracias garrulus    | 52                       |
|                              | Upupa epops          | 52                       |
|                              | Turdus merula        | 81                       |
|                              | Turdus philomelos    | 52, 117                  |
| <b>SCINCIDAE</b>             |                      |                          |
|                              | Ciconia ciconia      | 124                      |
|                              | Falco naumanni       | 50, 83                   |
| <b>CHALCIDES BEDRIAGAI</b>   |                      |                          |
|                              | Milvus milvus        | 25                       |
|                              | Circus pygargus      | 114                      |
|                              | Athene noctua        | 52, 136                  |
|                              | Lanius excubitor     | 190, 191                 |
| <b>CHALCIDES CHALCIDES</b>   |                      |                          |
|                              | Bulbucus ibis        | 49, 110, 138, 190        |
|                              | Circaetus gallicus   | 86                       |

|                   |                              |                               |
|-------------------|------------------------------|-------------------------------|
|                   | <i>Circus pygargus</i>       | 114, 154                      |
|                   | <i>Falco naumanni</i>        | 82, 83                        |
|                   | <i>Scolopax rusticola</i>    | 60                            |
| <b>LACERTIDAE</b> |                              |                               |
|                   | <i>Ardeidae</i>              | 153                           |
|                   | <i>Nycticorax nycticorax</i> | 49, 189                       |
|                   | <i>Ardeola ralloides</i>     | 49, 189                       |
|                   | <i>Bulbucus ibis</i>         | 19                            |
|                   | <i>Egretta garzetta</i>      | 49                            |
|                   | <i>Ardea cinerea</i>         | 49                            |
|                   | <i>Ardea purpurea</i>        | 49                            |
|                   | <i>Ciconia ciconia</i>       | 45, 49, 124, 150,<br>164      |
|                   | <i>Ciconia nigra</i>         | 49                            |
|                   | <i>Pernis apivorus</i>       | 50                            |
|                   | <i>Milvus milvus</i>         | 25, 50                        |
|                   | <i>Milvus migrans</i>        | 50                            |
|                   | <i>Accipiter gentilis</i>    | 85                            |
|                   | <i>Accipiter nisus</i>       | 50                            |
|                   | <i>Buteo buteo</i>           | 33, 50, 185                   |
|                   | <i>Hieraetus fasciatus</i>   | 50, 173                       |
|                   | <i>Aquila chrysaetos</i>     | 50, 182                       |
|                   | <i>Neophron percnopterus</i> | 50                            |
|                   | <i>Aegyptius monachus</i>    | 50                            |
|                   | <i>Circaetus gallicus</i>    | 4                             |
|                   | <i>Circus cyaneus</i>        | 169                           |
|                   | <i>Circus pygargus</i>       | 18, 114, 155, 169             |
|                   | <i>Circus aeruginosus</i>    | 169                           |
|                   | <i>Falco peregrinus</i>      | 50                            |
|                   | <i>Falco eleonora</i>        | 50                            |
|                   | <i>Falco naumanni</i>        | 18, 26, 83                    |
|                   | <i>Falco tinnunculus</i>     | 6, 7, 29, 77, 85,<br>172, 185 |
|                   | <i>Grus grus</i>             | 50                            |
|                   | <i>Porzana pusilla</i>       | 50                            |
|                   | <i>Otis tarda</i>            | 50                            |
|                   | <i>Larus argentatus</i>      | 132                           |
|                   | <i>Clamator glandarius</i>   | 52                            |
|                   | <i>Tyto alba</i>             | 32, 57, 76, 92,<br>185        |
|                   | <i>Bubo bubo</i>             | 27, 112, 184                  |
|                   | <i>Asio otus</i>             | 46, 52, 175, 196,<br>197      |
|                   | <i>Otus scops</i>            | 52                            |
|                   | <i>Athene noctua</i>         | 63, 136                       |
|                   | <i>Coracias garrulus</i>     | 52                            |
|                   | <i>Lanius senator</i>        | 88                            |
|                   | <i>Lanius excubitor</i>      | 190                           |
|                   | <i>Lanius collurio</i>       | 125                           |
|                   | <i>Saxicola torquata</i>     | 191                           |
|                   | <i>Monticola saxatilis</i>   | 53, 133                       |
|                   | <i>Monticola solitarius</i>  | 53                            |
|                   | <i>Turdus torquatus</i>      | 121                           |

|                            |                       |  |
|----------------------------|-----------------------|--|
|                            | Turdus merula         | 53   |
|                            | Turdus philomelos     | 53   |
|                            | Corvus corone         | 122  |
| ACANTHODACTYLUS ERYTHRURUS |                       |  |
|                            | Bulbucus ibis         | 138, 189, 191  |
|                            | Buteo buteo           | 89   |
|                            | Circus pygargus       | 154, 191   |
|                            | Falco tinnunculus     | 85, 191  |
|                            | Tyto alba             | 35, 52, 190, 191                                     |
|                            | Athene noctua         | 52, 191  |
|                            | Upupa epops           | 190  |
|                            | Lanius excubitor      | 190  |
| LACERTA AGILIS             |                       |  |
|                            | Circaetus gallicus    | 50   |
|                            | Phasianus colchicus   | 50   |
|                            | Bubo bubo             | 69   |
|                            | Athene noctua         | 52   |
|                            | Strix aluco           | 52   |
|                            | Coracias garrulus     | 52   |
|                            | Upupa epops           | 52   |
|                            | Erithacus rubecula    | 104  |
| LACERTA LEPIDA             |                       |  |
|                            | Bulbucus ibis         | 138  |
|                            | Ciconia ciconia       | 124  |
|                            | Ciconia nigra         | 62   |
|                            | Elanus caeruleus      | 106  |
|                            | Pernis apivorus       | 37, 50, 77   |
|                            | Milvus milvus         | 7, 25, 50, 65,<br>85, 86, 154, 190,<br>191           |
|                            | Milvus migrans        | 9, 11, 50, 61,<br>80, 85, 86, 152,<br>154, 190, 191  |
|                            | Accipiter gentilis    | 6, 7, 85, 86, 143                                    |
|                            | Buteo buteo           | 6, 7, 18, 33, 85,<br>86, 97, 143, 154,<br>191, 197   |
|                            | Hieraetus pennatus    | 7, 12, 18, 50,<br>85, 86, 119, 145,<br>154, 191, 197 |
|                            | Hieraetus fasciatus   | 6, 8, 28, 41, 50,<br>55, 84, 123, 159,<br>173, 195   |
|                            | Aquila heliaca        | 50, 62, 86, 190,<br>191                              |
|                            | Aquila chrysaetos     | 18, 64, 77, 78,<br>86, 123                           |
|                            | Neophron percnopterus | 6, 17, 50, 71,<br>85, 86, 127, 154,<br>163           |
|                            | Gypaetus barbatus     | 113  |
|                            | Aegypius monachus     | 111  |

|                    |                              |  |
|--------------------|------------------------------|--|
|                    | <i>Circaetus gallicus</i>    | 4, 18, 50, 86,<br>120, 183, 190,<br>191, 197 |
|                    | <i>Circus pygargus</i>       | 114, 154, 155                                |
|                    | <i>Falco naumanni</i>        | 50, 82, 83                                   |
|                    | <i>Falco tinnunculus</i>     | 7, 18, 191                                   |
|                    | <i>Gelochelidon nilotica</i> | 193  |
|                    | <i>Tyto alba</i>             | 52, 108, 109                                 |
|                    | <i>Bubo bubo</i>             | 27, 52, 112, 147,<br>156, 198                |
| LACERTA SCHREIBERI |                              |  |
|                    | <i>Ciconia ciconia</i>       | 142  |
|                    | <i>Pernis apivorus</i>       | 37, 50                                       |
|                    | <i>Buteo buteo</i>           | 7  |
| LACERTA VIRIDIS    |                              |  |
|                    | <i>Ardea cinerea</i>         | 144  |
|                    | <i>Pernis apivorus</i>       | 77   |
|                    | <i>Hieraetus fasciatus</i>   | 28, 41, 173, 195                             |
|                    | <i>Aquila chrysaetos</i>     | 13   |
|                    | <i>Neophron percnopterus</i> | 50   |
|                    | <i>Circaetus gallicus</i>    | 42, 50, 183                                  |
|                    | <i>Falco peregrinus</i>      | 165  |
|                    | <i>Tyto alba</i>             | 52   |
|                    | <i>Bubo bubo</i>             | 147  |
|                    | <i>Strix aluco</i>           | 40, 52                                       |
| LACERTA VIVIPARA   |                              |  |
|                    | <i>Circus cyaneus</i>        | 157  |
|                    | <i>Falco columbarius</i>     | 20   |
|                    | <i>Falco tinnunculus</i>     | 50, 59, 200, 205                             |
|                    | <i>Asio flammeus</i>         | 52   |
|                    | <i>Athene noctua</i>         | 52   |
|                    | <i>Coracias garrulus</i>     | 52   |
|                    | <i>Erithacus rubecula</i>    | 104  |
|                    | <i>Turdus torquatus</i>      | 53, 121                                      |
|                    | <i>Turdus philomelos</i>     | 39, 53                                       |
| PODARCIS SP.       |                              |  |
|                    | <i>Buteo buteo</i>           | 33   |
|                    | <i>Athene noctua</i>         | 136  |
|                    | <i>Coracias garrulus</i>     | 52   |
| PODARCIS HISPANICA |                              |  |
|                    | <i>Bulbucus ibis</i>         | 166  |
|                    | <i>Ciconia ciconia</i>       | 124  |
|                    | <i>Milvus milvus</i>         | 25   |
|                    | <i>Buteo buteo</i>           | 191  |
|                    | <i>Falco naumanni</i>        | 50, 83                                       |
|                    | <i>Falco tinnunculus</i>     | 148, 191                                     |
|                    | <i>Tyto alba</i>             | 35, 52, 108, 137,<br>198                     |
|                    | <i>Bubo bubo</i>             | 52, 112                                      |
|                    | <i>Athene noctua</i>         | 136  |
|                    | <i>Strix aluco</i>           | 201  |
|                    | <i>Lanius excubitor</i>      | 191  |
|                    | <i>Pica pica</i>             | 71   |

## PODARCIS MURALIS

|                    |     |
|--------------------|-----|
| Buteo buteo        | 33  |
| Circaetus gallicus | 183 |
| Tyto alba          | 52  |
| Asio flammeus      | 52  |
| Athene noctua      | 52  |
| Strix aluco        | 52  |

## PODARCIS LILFORDI

|                 |         |
|-----------------|---------|
| Falco eleonorae | 10, 167 |
|-----------------|---------|

## PSAMMODROMUS SP.

|                |            |
|----------------|------------|
| Bulbucus ibis  | 189        |
| Milvus milvus  | 50         |
| Buteo buteo    | 89         |
| Falco naumanni | 18, 50, 85 |
| Tyto alba      | 35, 108    |
| Athene noctua  | 136        |

## PSAMMODROMUS ALGIRUS

|                       |              |
|-----------------------|--------------|
| Bulbucus ibis         | 49, 110, 138 |
| Ciconia ciconia       | 124          |
| Milvus milvus         | 25           |
| Accipiter gentilis    | 6            |
| Buteo buteo           | 6            |
| Aegyptius monachus    | 111          |
| Circaetus gallicus    | 4, 86        |
| Circus pygargus       | 50, 114      |
| Falco naumanni        | 82, 83       |
| Falco tinnunculus     | 6, 18, 191   |
| Burhinus oedicephalus | 191          |
| Tyto alba             | 52, 190, 191 |
| Bubo bubo             | 52, 112      |
| Otus scops            | 6            |
| Athene noctua         | 52, 136      |
| Upupa epops           | 190, 191     |
| Lanius excubitor      | 190, 191     |

## PSAMMODROMUS HISPANICUS

|                   |          |
|-------------------|----------|
| Bulbucus ibis     | 138      |
| Milvus milvus     | 86       |
| Buteo buteo       | 86       |
| Circus pygargus   | 86       |
| Falco naumanni    | 83       |
| Falco tinnunculus | 6, 191   |
| Tyto alba         | 137, 198 |

## SERPENTES

## COLUBRIDAE

|                     |                                 |
|---------------------|---------------------------------|
| Ciconia ciconia     | 124, 164                        |
| Ciconia nigra       | 49                              |
| Milvus milvus       | 25, 190, 191                    |
| Milvus migrans      | 191                             |
| Buteo buteo         | 6, 7, 86, 97, 185               |
| Hieraetus fasciatus | 41                              |
| Aquila heliaca      | 190, 191                        |
| Aquila chrysaetos   | 13, 43, 44, 64,<br>78, 123, 182 |

|                          |                       |   |
|--------------------------|-----------------------|---|
|                          | Neophron percnopterus | 6, 17, 50                               |
|                          | Gypaetus barbatus     | 113                                     |
|                          | Circaetus gallicus    | 197                                     |
|                          | Circus pygargus       | 18, 114                                 |
|                          | Circus aeruginosus    | 169                                     |
|                          | Falco tinnunculus     | 77, 185                                 |
|                          | Grus grus             | 50                                      |
|                          | Burhinus oedicephalus | 191                                     |
|                          | Tyto alba             | 185                                     |
|                          | Bubo bubo             | 6, 112, 184                             |
|                          | Monticola solitarius  | 53                                      |
|                          | Turdus merula         | 170                                     |
|                          | Sturnus unicolor      | 190                                     |
|                          | Corvus corax          | 56                                      |
| COLUBER HIPPOCREPIS      |                       |   |
|                          | Buteo buteo           | 85, 86                                  |
|                          | Aquila heliaca        | 62                                      |
|                          | Circaetus gallicus    | 4, 86                                   |
| COLUBER VIRIDIFLAVUS     |                       |   |
|                          | Circaetus gallicus    | 50, 183                                 |
|                          | Bubo bubo             | 99                                      |
| CORONELLA SP.            |                       |   |
|                          | Ciconia ciconia       | 49                                      |
|                          | Circaetus gallicus    | 183                                     |
| CORONELLA AUSTRIACA      |                       |   |
|                          | Ciconia ciconia       | 150                                     |
|                          | Circaetus gallicus    | 50, 86                                  |
| CORONELLA GIRONDICA      |                       |   |
|                          | Milvus migrans        | 11                                      |
|                          | Buteo buteo           | 6, 33                                   |
|                          | Circaetus gallicus    | 4                                       |
| ELAPHE LONGISSIMA        |                       |   |
|                          | Pernis apivorus       | 50                                      |
|                          | Circaetus gallicus    | 42, 50, 183                             |
| ELAPHE SCALARIS          |                       |   |
|                          | Milvus milvus         | 50, 65, 86                              |
|                          | Milvus migrans        | 80                                      |
|                          | Buteo buteo           | 50, 86, 154                             |
|                          | Aquila heliaca        | 50, 62                                  |
|                          | Aquila chrysaetos     | 18, 139                                 |
|                          | Neophron percnopterus | 17, 73, 127                             |
|                          | Circaetus gallicus    | 4, 50, 85, 86,<br>120, 183, 190,<br>191 |
|                          | Strix aluco           | 40                                      |
| MACROPROTODON CUCULLATUS |                       |   |
|                          | Milvus milvus         | 50, 86                                  |
| MALPOLON MONSPESSULANUS  |                       |   |
|                          | Milvus milvus         | 50, 65, 85, 86                          |
|                          | Milvus migrans        | 85, 190, 191                            |
|                          | Buteo buteo           | 6, 33, 86                               |
|                          | Hieraetus fasciatus   | 6, 123                                  |
|                          | Aquila heliaca        | 50, 62, 190, 191                        |



|               |                              |   |
|---------------|------------------------------|---|
|               | <i>Aquila chrysaetos</i>     | 64                                      |
|               | <i>Neophron percnopterus</i> | 17, 50, 73, 85,<br>127, 163             |
|               | <i>Aegyptius monachus</i>    | 111                                     |
|               | <i>Circaetus gallicus</i>    | 4, 50, 85, 86,<br>120, 183, 190,<br>191 |
|               | <i>Falco naumanni</i>        | 82                                      |
|               | <i>Pica pica</i>             | 72                                      |
| NATRIX SP.    |                              |   |
|               | <i>Nycticorax nycticorax</i> | 189                                     |
|               | <i>Ardea purpurea</i>        | 49                                      |
|               | <i>Ciconia ciconia</i>       | 124                                     |
|               | <i>Pernis apivorus</i>       | 50                                      |
|               | <i>Milvus milvus</i>         | 50                                      |
|               | <i>Milvus migrans</i>        | 50                                      |
|               | <i>Circaetus gallicus</i>    | 18                                      |
|               | <i>Strix aluco</i>           | 52                                      |
|               | <i>Turdus merula</i>         | 53                                      |
|               | <i>Garrulus glandarius</i>   | 115                                     |
| NATRIX MAURA  |                              |   |
|               | <i>Nycticorax nycticorax</i> | 190                                     |
|               | <i>Ardea cinerea</i>         | 144                                     |
|               | <i>Ardea purpurea</i>        | 2, 162, 190, 191                        |
|               | <i>Ciconia nigra</i>         | 73                                      |
|               | <i>Milvus milvus</i>         | 25, 65, 86, 191                         |
|               | <i>Milvus migrans</i>        | 61, 80, 190, 191                        |
|               | <i>Buteo buteo</i>           | 85, 97, 191                             |
|               | <i>Aquila heliaca</i>        | 62                                      |
|               | <i>Neophron percnopterus</i> | 17, 154                                 |
|               | <i>Circaetus gallicus</i>    | 4, 50, 183, 190,<br>191                 |
|               | <i>Circus aeruginosus</i>    | 50                                      |
|               | <i>Falco naumanni</i>        | 50, 83                                  |
|               | <i>Porphyrio porphyrio</i>   | 50                                      |
|               | <i>Athene noctua</i>         | 52                                      |
| NATRIX NATRIX |                              |   |
|               | <i>Podiceps cristatus</i>    | 49                                      |
|               | <i>Botaurus stellaris</i>    | 49                                      |
|               | <i>Nycticorax nycticorax</i> | 49                                      |
|               | <i>Bulbucus ibis</i>         | 166                                     |
|               | <i>Ardea cinerea</i>         | 49, 144                                 |
|               | <i>Ardea purpurea</i>        | 23                                      |
|               | <i>Ciconia ciconia</i>       | 49, 150                                 |
|               | <i>Pernis apivorus</i>       | 22, 50                                  |
|               | <i>Milvus milvus</i>         | 65, 86                                  |
|               | <i>Milvus migrans</i>        | 80                                      |
|               | <i>Buteo buteo</i>           | 24, 50, 97                              |
|               | <i>Neophron percnopterus</i> | 17, 86, 127                             |
|               | <i>Circaetus gallicus</i>    | 4, 42, 50, 183                          |
|               | <i>Circus pygargus</i>       | 50                                      |
|               | <i>Circus aeruginosus</i>    | 50                                      |
|               | <i>Phasianus colchicus</i>   | 50                                      |

|                 |                     |         |
|-----------------|---------------------|---------|
|                 | Tyto alba           | 52, 185 |
|                 | Bubo bubo           | 52      |
|                 | Athene noctua       | 52      |
|                 | Coracias garrulus   | 52      |
| VIPERIDAE       |                     |         |
|                 | Ciconia ciconia     | 30, 150 |
|                 | Buteo buteo         | 50      |
|                 | Aquila chrysaetos   | 50      |
| VIPERA ASPIS    |                     |         |
|                 | Circaetus gallicus  | 42, 50  |
| VIPERA LATASTEI |                     |         |
|                 | Circaetus gallicus  | 4       |
| VIPERA BERUS    |                     |         |
|                 | Circaetus gallicus  | 50      |
|                 | Circus pygargus     | 50      |
|                 | Phasianus colchicus | 50      |
|                 | Larus argentatus    | 130     |
|                 | Picus viridis       | 52      |

## REFERENCES

1. Amat, J.A. 1986. Information on the diet of the stone curlew Burhinus oedicephalus in Doñana, Southern Spain. *Bird Study* 33(2): 71-73.
2. Amat, J.A., and C.M. Herrera. 1977. Alimentación de la garza imperial (Ardea purpurea) en las Marismas del Guadalquivir durante el periodo de nidificación. *Ardeola* 24: 95-104.
3. Amat, J.A., and R.C. Soriguer. 1981. Analyse comparative de régimes alimentaires de L'effraie Tyto alba et du Moyen-Duc Asio otus dans l'ouest de l'Espagne. *Alauda* 49(2): 112-120.
4. Amores, F., and A. Franco. 1981. Alimentation et écologie du Circaète Jean-le-Blanc dans le sud de l'Espagne. *Alauda* 49(1): 59-61.
5. Amores, F., A. Franco, and J. Mellado. 1979. Presencia de Bufo bufo spinosus Daudin en la dieta de Falconiformes y Strigiformes en Sierra Morena Occidental. Doñana, *Acta Vert.* 6(2): 238-239.
6. Amores, F., J. Oria, E. Roy, and F. Torrent. 1984. Estudio faunístico de la zona del "Alto Tajo" comprendida entre el puente de Valtablado del Río y el Puente de San Pedro. Monogr. I.C.O.N.A. 32. Minist. Agricult.; Madrid.
7. Araujo, J. 1973. Falconiformes del Guadarrama Suroccidental. *Ardeola* 19(2): 257-278.
8. Araujo, J., B. Arroyo, and J.M. Bueno. 1974. Un nido de Aguila perdicera (Hieraetus fasciatus) con tres pollos. *Ardeola* 20: 343-345.
9. Araujo, J., M. Fernandez Cruz, and C. Garcia Rua. 1973. Contenido de un nido de Milvus migrans. *Ardeola* 19(1): 10.
10. Araujo, J., J. Muñoz Cobo, and F.J. Purroy. 1977. Las rapaces y aves marinas del archipiélago de Cabrera. *Naturalia Hispanica* 12. I.C.O.N.A.; Madrid.
11. Arroyo, B. 1978. Alimentación del Milano Negro (Milvus migrans) en una localidad de España Central. *Ardeola* 25: 47-58.
12. Arroyo, B., J.M. Bueno, and V. Pérez Mellado. 1976. Biología de reproducción de una pareja de Hieraetus fasciatus en España Central. Doñana, *Acta Vert.* 3(1): 33-45.
13. Austruy, J.C., and J.M. Cugnasse. 1981. L'Aigle royal Aquila chrysaetos, dans le Massif Central. *Nos oiseaux* 36(4): 133-142.
14. Bagle, P., P.H. Orsini, and J. Boutin. 1987. Variations du régime alimentaire du Hibou grand-duc Bubo bubo en période de reproduction en Basse-Provence. *L'oiseau et R.F.O.* 57(1): 23-31.
15. Bauduin, H. 1980. Les surplus de proies au site de nid chez la chouette effraie, Tyto alba. *Nos oiseaux* 35(5): 232-238.

16. Baudin, H., and J.L. Dessolin. 1987. A propos de la Chouette hulotte, Strix aluco: réponses apportées et questions posées par les "nichoirs-photos". Nos oiseaux 39(3): 129-134.
17. Bergier, P., and G. Cheylan. 1980. Statut, succès de reproduction et alimentation du rautour pernoptère Neophron percnopterus en France méditerranéenne. Alauda 48(2-3): 75-97.
18. Bernis, F. 1973. Algunos datos de alimentación y depredación de Falconiformes y Estrigiformes ibéricas. Ardeola 19(2): 225-248.
19. Bernis, F., and J.A. Valverde. 1954. La gran colonia de garzas de Doñana en 1953. Munibe 6: 1-37.
20. Bibby, C.J. 1987. Foods of breeding Merlins Falco columbarius in Wales. Bird Study 34(1): 64-70.
21. Blanc, T. 1956. Une accumulation de proies sur un nid de Buses. Nos oiseaux 23(8): 217-218.
22. Blanc, T. 1957. Quelques notes sur le régime de la Bondrée. Nos oiseaux 24(1): 16.
23. Blanc, T. 1957. Un jeune Héron pourpré dégorge une couleuvre. Nos oiseaux 24(2): 79.
24. Blanc, T. 1960. La voracité d'un couple de Buses Buteo buteo. Nos oiseaux 25(9): 254.
25. Blanco, J.C., F. Hiraldo, B. Heredia, and L. García. 1987. Alimentación invernal del Milano Real, Milvus milvus (L. 1758) en el Parque Nacional de Doñana. Bol. Est. Centr. Ecol. 16(31): 93-97.
26. Blondel, J. 1964. Notes sur la biologie et le régime alimentaire du Faucon crécerellette Falco naumanni. Nos oiseaux 27(10-11): 329-332.
27. Blondel, J., and O. Badan. 1976. La biologie du Hibou grand-duc en Provence. Nos oiseaux 33(5): 189-219.
28. Blondel, J., L. Coulon, B. Gired, and M. Hortigue. 1969. Deux cents heures d'observation auprès de l'aire de l'Aigle de Bonelli. Nos oiseaux 30(2-3): 37-60.
29. Bonin, B., and L. Strenna. 1986. Sur la biologie du faucon crécerelle Falco tinnunculus en Auxois. Alauda 54(4): 241-262.
30. Bouet, G. 1956. L'alimentation des jeunes cigognes blanches au nid. Rôle des insects. L'oiseau et R.F.O. 26: 59-61.
31. Brown, D.J. 1981. Seasonal variation in the prey of some Barn owls in Gwynedd. Bird Study 28(2): 139-146.
32. Brunet-Leconte, P., and M. Delibes. 1984. Alimentación de la lechuza común Tyto alba en la Cuenca del Duero, España. Doñana, Acta Vert. 11(2): 213-229.
33. Bustamante, J.M. 1985. Alimentación del ratonero común (Buteo buteo, L. 1758) en el Norte de España. Doñana, Acta Vert. 12(1): 51-62.
34. Busser-Stalder, M. 1984. Régime alimentaire de la chouette effraie Tyto alba, durant les périodes de nidification 1978 et 1979, en Anjoie (Jura Suisse). Nos oiseaux 37(8): 392-394.

35. Camacho, I. 1975. La alimentación de Tyto alba (Scop.) en la Vega de Granada. Cuad. Cienc. Biol. (Granada) 1975(4.2): 111-124.
36. Campos, F. 1977. Régimen alimenticio de Tyto alba en las provincias de Salamanca y Zamora. Ardeola 24: 105-119.
37. Castroviejo, J., and S. Fernández. 1968. Nidificación del halcón abejero (Pernis apivorus) en el Bierzo (León). Ardeola 14: 131-135.
38. Catley, G.P. 1983. Unusual feeding action of great white egret. British Birds 76(2): 82-83.
39. Chater, A.O. 1965. Song thrush taking lizard. British Birds 58(12): 513.
40. Cheylan, G. 1971. Le régime de la chouette hulotte (Strix aluco) a Salernes (Var). Alauda 39(2): 150-155.
41. Cheylan, G. 1977. La place trophique de l'aigle de Bonelli Hieraaetus fasciatus dans le biocenoses mediterraneennes. Alauda 45(1): 1-15.
42. Choussy, D. 1973. Observations sur le circaète Jean-le-Blanc. Nos oiseaux 32(4): 83-89.
43. Clouet, M. 1981. L'Aigle Royal (Aquila chrysaetos) dans le Pyrénées Françaises. Resultats de 5 ans d'observations. L'oiseau et R.F.O.: 51(2): 89-100.
44. Clouet, M. 1982. L'Aigle Royal dans les Pyrénées Françaises. Acta Biol. Montana 1: 269-280.
45. Collin, A. 1973. Nidification de la cigogne blanche (Ciconia ciconia) en 1972 a Hachy (Lorraine Belge). Aves 10(2): 29-69.
46. Corral, J.F., J.A. Cortés, and J.M. Gil. 1979. Contribución al estudio de la alimentación de Asio otus en el sur de España. Doñana, Acta Vert. 6(2): 179-190.
47. Cortés, J.A. 1988. Sobre diferencias individuales en la alimentación de Tyto alba. Doñana, Acta Vert. 15(1): 99-109.
48. Costa, L. 1984. Alimetación de la pagaza piconegra (Gelochelidon nilotica) en las Marismas del Guadalquivir. Doñana, Acta Vert. 11(2): 185-195.
49. Cramp, S. (ed). 1977. Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. Vol.1, Ostrich to Ducks. Oxford Univ. Press.
50. Cramp, S. (ed). 1980. Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. Vol.2, Hawks to Bustards. Oxford Univ. Press.
51. Cramp, S. (ed). 1983. Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. Vol.3, Wader to Gulls. Oxford Univ. Press.
52. Cramp, S. (ed). 1985. Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. Vol.4, Terns to Woodpeckers. Oxford Univ. Press.
53. Cramp, S. (ed). 1988. Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. Vol.5, Tyrant Flycatchers to Trushes. Oxford Univ. Press.

54. Cugnasse, J.M. 1983. Contribution à L'étude du Hibou grand-duc, Bubo bubo, dans le Sud du Massif Central. Nos oiseaux 37(3): 117-128.
55. Cugnasse, J.M. 1984. L'Aigle de Bonelli, Hieraetetus fasciatus, en Languedoc-Rousillon. Nos oiseaux 37(5): 223-232.
56. Cugnasse, J.M., and C. Riols. 1987. Note sur le régime alimentaire du grand corbeau, Corvus corax, dans le Sud du Massif Central. Nos oiseaux 39(2): 57-65.
57. Cuisin, J., and M. Cuisin. 1979. Le régime alimentaire du la chouette effraie (Tyto alba (Scopoli)) dans le canton des Ricys (Aube) et ses environs immédiats. L'oiseau et R.F.O. 49(2): 81-89.
58. Davis, P.E., and J.E. Davis. 1981. The food of the red kite in Wales. Bird Study 28(1): 33-40.
59. Davis, T.A.W. 1975. The food of the kestrel in winter and early spring. Bird Study 22(2): 85-91.
60. Debussche, M., and P. Isenmann. 1986. Consommation exceptionnelle d'un vertébré (Chalcides chalcides) par une bécasse (Scolopax rusticola). L'oiseau et R.F.O. 56(2): 212.
61. Delibes, M. 1975. Alimentación del Milano Negro (Milvus migrans) en Doñana (Huelva, España). Ardeola 21(1): 183-207.
62. Delibes, M. 1978. Ecología alimentaria del Aguila Imperial Ibérica (Aquila adalberti) en el Coto de Doñana durante la crianza de los pollos. Doñana, Acta Vert. 5(1-2): 35-60.
63. Delibes, M., P. Brunet-Lecomte, and M. Mañez. 1983. Datos sobre la alimentación de la lechuza común (Tyto alba), el buho chico (Asio otus) y el mochuelo (Athene noctua) en una misma localidad de Castilla la Vieja. Ardeola 30: 57-63.
64. Delibes, M., J. Calderón and F. Hiraldo. 1975. Selección de presas y alimentación en España del Aguila Real (Aquila chrysaetos). Ardeola 21(1): 285-303.
65. Delibes, M., and L. García. 1984. Hábitos alimenticios del Milano Real en Doñana durante el periodo de cría. Ardeola 31: 115-121.
66. Delmee, E. 1985. Régime alimentaire de la chouette effraie (Tyto alba) et evolution du statut des micromammiferes d'un paysage agricole. Aves 22(3): 169-174.
67. Delmee, E., P. Dachy, and P. Simon. 1979. Etude comparative du regimen alimentaire d'une population forestiere de chouettes hulottes (Strix aluco). Le Gerfaut 69(1): 45-77.
68. Delmee, E., P. Dachy, and P. Simon. 1982. Particularites ecologiques des chouettes hulottes (Strix aluco) de la Foret de Belaeil-en-Hainaut. Le Gerfaut 72(3): 287-306.
69. Desfayes, M., and P. Geroudet. 1949. Notes sur le grand-duc Bubo bubo (L.). Nos oiseaux 20(3-4): 49-60.

70. Domínguez, L., C.R. Vigal, E. Lázaro, F. Palacios, and J. Garzón. 1980. Alimentación de la urraca (Pica pica L) en el centro de España. Actas I Reunión Iberoamer. Zool. Vert. La Rábida 1977: 799-805.
71. Domínguez, L., J.L. González, L.M. González, J. Garzón, and C. Llandrés. 1985. Datos sobre la alimentación de la Cigüeña Negra (Ciconia nigra L.) en España Centro occidental. Alytes 3: 51-56.
72. Donazar, J.A., and O. Ceballos. 1988. Sur l'alimentation du Hibou Grand duc (Bubo bubo) dans une localité de la haute montagne pyrénéenne espagnole. Alauda 56(3): 274-276.
73. Donazar, J.A., and O. Ceballos. 1988. Alimentación y tasas reproductoras del alimoche (Neophron percnopterus) en Navarra. Ardeola 35(1): 3-14.
74. Doornbos, G. 1979. Winter food habits of smew (Mergus albellus L.) on Lake Yssel, The Netherlands: species and size selection in relation to the fish stocks. Ardea 67(1-2): 42-48.
75. Doucet, J. 1987. Aspects de la predation de l'autour (Accipiter gentilis) en periode de nidification. Aves, NQesp.: 62-65.
76. Dueñas, M.E., and S.J. Peris. 1985. Análisis de egagrópilas de Tyto alba en la Sierra de Gata (W de España). Alytes 3: 109-144.
77. Elosegui, J. 1973. Informe preliminar sobre alimentación de aves rapaces en Navarra y provincias limítrofes. Ardeola 19(2): 249-256.
78. Elosegui, J. 1975. Notas sobre la Biología del Aguila Real (Aquila chrysaetos, L.) en Navarra. Munibe 27(3-4): 167-172.
79. Fairley, J.S., and F.L. Clark. 1972. Food of barn owls Tyto alba (Scopoli) over one year at roost in Co. Galway. Irish Natur. J. 17(7): 219-222.
80. Fernández Cruz, M. 1973. Sobre la alimentación del Milano Negro (Milvus migrans) en la estación ornitológica de "El Borbollón" (Cáceres). Ardeola 19(2): 337-343.
81. Foote, R.J. 1978. Blackbird eating slow-worm's tail. British Birds 71(3): 131.
82. Franco, A. 1980. Biología de caza en Falco naumanni. Doñana, Acta Vert. 7(2): 213-227.
83. Franco, A., and J. Andrada. 1976. Alimentación y selección de presa en Falco naumanni. Ardeola 23: 137-187.
84. García, L. 1976. Reproducción del Aguila perdicera, Hieraetus fasciatus, en la Sierra de Cabo de Gata de Almería. Bol. Est. Centr. Ecol. 5: 83-92.
85. Garzón, J. 1968. Las rapaces y otras aves de la Sierra de Gata. Ardeola 14: 97-130.
86. Garzón, J. 1973. Contribución al estudio del status, alimentación y protección de la Falconiformes en España Central. Ardeola 19(2): 279-330.

87. Garzón, J. 1974. Ciconia nigra y Ciconia ciconia criando en nidos contiguos. *Ardeola* 20: 321-324.
88. Gil Lletget, A. 1944. Bases para el estudio científico de la alimentación en las aves y resultado del análisis de 400 estómagos. *Bol. Real. Soc. Hist. Nat.* 52: 533-564.
89. Gil Lletget, A. 1945. Bases para el estudio científico de la alimentación en las aves y resultado del análisis de 400 estómagos. *Bol. Real. Soc. Hist. Nat.* 53: 9-23.
90. Giles, N. 1981. Summer diet of the grey Heron. *Scott. Birds* 11(5): 153-159.
91. Glue, D.E. 1970. Prey taken by short-eared owls at British breeding sites and winter quarters. *Bird Study* 17(1): 39-42.
92. Glue, D.E. 1974. Food of the barn owl in Britain and Ireland. *Bird Study* 21(3): 200-210.
93. Glue, D.E. 1977. Feeding ecology of the short-eared owl in Britain and Ireland. *Bird Study* 24(2): 70-78.
94. Glue D.E., and G.J. Hammond. 1974. Feeding ecology of the long-eared owl in Britain and Ireland. *British Birds* 67(9): 361-369.
95. Godin, J. 1975. Données sur le régime alimentaire de la chouette effraie (Tyto alba) en Belgique et dans le Nord de la France. *Aves* 12(3): 105-126.
96. Gomes Oliveira, N. 1971. A colonia da mata de S. Jacinto (Aveiro) em 1971. *Cyanopica* 1(3): 38-44.
97. Gomez Ruiz, R. 1969. Datos sobre la reproducción del águila ratonera (Buteo buteo) en la provincia de Huelva. *Ardeola* 15: 32-35.
98. González, L.M., and J.L. González. 1984. Sobre la alimentación del avetoro común (Botaurus stellaris) en España. *Doñana, Acta Vert.* 11(1): 140-141.
99. Guichard, G. 1956. Notes sur le biologie du grand duc (Bubo bubo L.). *L'oiseau et R.F.O.* 6(2): 126-134.
100. Hafner, H., V. Boy and G. Gory. 1982. Feeding methods, flock size and feeding success in the little egret Egretta garzetta and the squacco heron Ardeola ralloides in Camargue, Southern France. *Ardea* 70(1): 45-54.
101. Hallet, C., and J. Doucet. 1982. Le martin-pêcheur (Alcedo atthis) en Wallonie: statut des populations et mesures de protection. *Aves* 19(1): 1-12.
102. Halliday, K.C.R. 1970. Notes on a great grey shrike wintering in Lanarkshire. *Scott. Birds* 6: 22-23.
103. Harris, M.P. 1965. The food of some Larus gulls. *The Ibis* 107(1): 43-53.
104. Heath, P.J. 1988. Robin attacking common lizard. *British Birds* 81(5): 238.
105. Henry, C., and A. Perthuis. 1986. Composition et structure du régime alimentaire de la chouette hulotte (Strix aluco L.) dans deux régions forestières du Centre de la France. *Alauda* 54(1): 49-65.



106. Heredia, B. 1983. Alimentación invernal del elanio azul (Elanus caeruleus) en la provincia de Salamanca. *Alytes* 1(1): 255-265.
107. Heredia, R. 1973. Nota sobre la alimentación y reproducción de Aquila chrysaetos en Soria y Toledo. *Ardeola* 19(2): 459-460.
108. Herrera, C.M. 1973. Régimen alimenticio de Tyto alba en España Sudoccidental. *Ardeola* 19(2): 359-394.
109. Herrera, C.M. 1973. La captura de carnívoros por las Strigiformes. *Ardeola* 19(2): 439-444.
110. Herrera, C.M. 1974. Observaciones sobre una colonia de Garcillas bueyeras (Bulbucus ibis L.) en Andalucía. *Ardeola* 20: 287-306.
111. Hiraldo, F. 1976. Diet of the black vulture (Aegypius monachus) in the Iberian Peninsula. *Doñana, Acta Vert.* 3(1) : 19-31.
112. Hiraldo, F., J. Andrada, and F.F. Parreño. 1975. Diet of the eagle owl (Bubo bubo) in Mediterranean Spain. *Doñana, Acta Vert.* 2(2): 161-177.
113. Hiraldo, F., M. Delibes, and J. Calderón. 1979. El quebrantahuesos Gypaetus barbatus (L.). Sistemática, taxonomía, biología, distribución y protección. Monogr. I.C.O.N.A. 22. Minist. Agricult.; Madrid.
114. Hiraldo, F., F. Fernández, and F. Amores. 1975. Diet of the Montagu's Harrier (Circus pygargus) in southwestern Spain. *Doñana, Acta Vert.* 2(1): 25-55.
115. Holyoak, D. 1968. A comparative study of the food of some british Corvidae. *Bird Study* 15(3): 147-154.
116. Horter, M, and H. Greven. 1981. The relative edibility of juvenile fire salamanders Salamandra salamandra. *Amphibia-Reptilia* 2(1): 15-22.
117. Husband, C.I. 1967. Song thrush attacking slow-worm. *British Birds* 60(4): 168-169.
118. Iribarren, I.B., and L. Domínguez. 1982. Contribution a l'étude du régime alimentaire du Martin-pêcheur (Alcedo atthis L. 1758). *Alauda* 50(2): 81-91.
119. Iribarren, J.J. 1975. Biología del Aguila calzada (Hieraetus pennatus) durante el periodo de nidificación en Navarra. *Ardeola* 21(1): 305-320.
120. Iribarren, J.J., and A. Rodríguez-Arbeola. 1973. Observaciones de un nido del Aguila culebrera (Circaetus gallicus), Navarra 1972. *Ardeola* 19(1): 101-106.
121. Isenmann, P. 1985. Nidification rupestre et prédation sur un lacertidé du Merle à plastron (Turdus torquatus alpestris) en Savoie. *Alauda* 53(3): 231-232.
122. Jollet, A. 1984. Variations saisonnières du régime alimentaire de la Cornielle noire (Corvus corone L.) dans le bocage limousin. *L'oiseau et R.F.O.* 54(2): 109-130.
123. Jordano, P. 1981. Relaciones interespecíficas y coexistencia entre el Aguila Real (Aquila chrysaetos) y el Aguila Perdicera (Hieraaetus fasciatus) en Sierra Morena Central. *Ardeola* 28: 67-88.

124. Lázaro, E. 1984. Contribución al estudio de la alimentación de la Cigüeña Blanca (Ciconia c. ciconia) (L.) en España. Tesis Doctoral. Univ. Complutense Madrid.
125. Lefranc, N. 1979. Contribution à l'écologie de la Pie-grièche écorcheur Lanius collurio L. dans les Vosges moyennes. L'oiseau et R.F.O. 49(4): 245-298.
126. Leveque, R. 1956. Une colonie de sternes hansel en Camarge. Nos oiseaux 23(9): 233-246.
127. Leveque, R. 1964. A propos d'un nid provençal du Percnoptère. Nos oiseaux 27(12): 329-332.
128. Libois, R. 1977. Contribution a l'étude du régime alimentaire de la chouette cheveche (Athene noctua) en Belgique. Aves 14(3): 165-177.
129. López-Gordo, J.L., E. Lázaro, and M. Fernández Cruz. 1976. Comparación de las dietas de Strix aluco, Asio otus y Tyto alba en un mismo biotopo de la provincia de Madrid. Ardeola 23: 189-221.
130. Lowe, W.P. 1940. Herring-gull and snake. The Ibis 4: 728.
131. Mac Nally, L. 1962. Food at a buzzards' nest. Scott. Birds 2(3): 196-198.
132. Madon, P. 1935. Contribution a l'étude du régime des oiseaux aquatics. Alauda 7: 382-401.
133. Magnenat, D. 1969. Proies volumineuses du Merle de roche Monticola saxatilis. Nos oiseaux 30(2-3): 69.
134. Maguire, E.J. 1979. Notes on the breeding of the buzzard in Kintyre. Western Natur. 8: 3-13.
135. Máñez, M. 1983. Espectro alimenticio del Mochuelo común (Athene noctua) en España. Alytes 1(1): 275-290.
136. Máñez, M. 1983. Variaciones geográficas y estacionales en la dieta del mochuelo común (Athene noctua) en España. Actas XV Congr. Int. Fauna Cinegética y Silvestre. Trujillo 1981: 617-634.
137. Martín, J., and J.R. Vericad. 1977. Datos sobre la alimentación de la lechuza (Tyto alba) en Valencia. Mediterránea 2: 35-47.
138. Mateos, A., and E. Lázaro. 1986. Contribución al estudio de la alimentación de la garcilla bueyera (Bulbucus ibis ibis L.) en Extremadura. Alytes 4: 49-68.
139. Mathieu, R., and J.P. Choisy. 1982. L'Aigle royal (Aquila chrysaetos) dans les Alpes Méridionales françaises de 1964 à 1980. Bièvre 4(1): 1-32.
140. Mc. Dougall, D. 1983. Cuckoo eating young amphibians. Scott. Birds 12(5): 162-163.
141. Melendro, J., and J. Gisbert. 1977. Notas ornitológicas breves. Mochuelo. Ardeola 24: 261.
142. Melendro, J., J. Gisbert, and A. Rodriguez Valverde. 1977. Datos sobre la alimentación de Ciconia ciconia. Ardeola 24: 207-209.
143. Morillo, C., and J. Lalanda. 1972. Primeros datos sobre la ecología de las Falconiformes en los Montes de Toledo. Bol. Est. Centr. Ecología 1(2): 57-67.

144. Moser, M.E. 1986. Prey profitability for adult grey herons Ardea cinerea and the constraints on prey size when feeding young nestlings. *The Ibis* 128: 392-405.
145. Nevado, J.C., L. García, and J.A. Oña. 1988. Sobre la alimentación del Aguila calzada (Hieraaetus pennatus) en las Sierras del Norte de Almería en la época de reproducción. *Ardeola* 35(1): 147-150.
146. Nos, M.R. 1961. Estudio de los ovillos regurgitados por una pareja de Tyto alba en la comarca de la Maresma (Provincia de Barcelona). *Misc. Zool.* 1: 139-146.
147. Orsini, P. 1985. Le régime alimentaire du hibou grand-duc Bubo bubo en Provence. *Alauda* 53(1): 11-28.
148. Otero, C., E. Castán, R. Senosiain, and F. Portillo. 1978. Fauna de Cazorla. Vertebrados. Monogr. I.C.O.N.A. 19. Minist. Agricult.; Madrid.
149. Owen, D.F. 1955. The food of the heron Ardea cinerea in the breeding season. *The Ibis* 97(2): 276-295.
150. Parent, G.H. 1973. La signification ecologique de la nidification de la Cigogne blanche (Ciconia ciconia L.) en Lorraine Belge, en 1972. *Aves* 10(2): 70-112.
151. Pechuán, L. 1971. Sobre la colonia de garzas del lago de La Albufera de Valencia. *Ardeola* vol.esp.: 397-404.
152. Pérez-Chiscano, J.L. 1967. Estudio de una comunidad de aves de presa (primavera-verano, 1968). *Ardeola* 13(1): 177-189.
153. Pérez-Chiscano, J.L. 1970. Notas sobre nuevas colonias de Ardeidas y otros datos de aves de Extremadura. *Ardeola* 16: 246-250.
154. Pérez-Chiscano, J.L. 1973. Sumario informe sobre la alimentación de rapaces en el nor-este de la provincia de Badajoz. *Ardeola* 19(2): 331-336.
155. Pérez-Chiscano, J.L., and M. Fernández Cruz. 1971. Sobre Grus grus y Circus pygargus en Extremadura. *Ardeola* vol esp.: 509-574.
156. Pérez Mellado, V. 1978. Alimentación del Buho Real (Bubo bubo L.) en España Central. *Ardeola* 25: 98-112.
157. Picozzi, N. 1978. Dispersion, breeding and prey of the hen harrier Circus cyaneus in Glen Dye, Kincardineshire. *The Ibis* 120(4): 498-509.
158. Ponting, E.D. 1967. Cormorant eating frog. *British Birds* 60(11): 482.
159. Real i Orti, J. 1982. Premieres donnees sur une population d'aigle de Bonelli dans une partie de Cordillere pre-littorale catalane. *Acta Biol. Montana* 1: 293-301.
160. Rebecca, G.W. 1983. Cuckoo eating young amphibians. *Scott. Birds* 12(5): 162-163.
161. Ritchie, J. 1926. A curious food suply of the Common gull. Common gull feeding upon frogs. *Scotland Natur.* 1926: 22-23.

162. Rodríguez de los Santos, M., and J.P. Canavate. 1985. Sélection des proies par le Héron pourpré Ardea purpurea pendant la période de reproduction dans les Marais du Guadalquivir (Espagne). *L'oiseau et R.F.O.* 55: 195-204.
163. Rodríguez Jiménez, F.L., and E. Balcells. 1968. Notas biológicas sobre el alimoche, Neophron percnopterus, en el Alto Aragón. *Publ. Cent. pir. Biol. Exp.* 2: 159-187.
164. Rooth, J. 1957. Over het voedsel, de terreinkeus en de achteruitgang van de ooievaar, Ciconia ciconia L., in Nederland. *Ardea* 45(3-4): 93-116.
165. Rosnoblet, R. 1974. Le faucon pèlerin prédateur du lézard vert. *Nos oiseaux* 32(7): 202.
166. Ruiz, X. 1985. An analysis of the diet of cattle egrets in the Ebro Delta, Spain. *Ardea* 73: 49-60.
167. Salvador, A. 1980. Interaction between the balearic lizard (Podarcis lilfordi) and Eleonora's falcon (Falco eleonorae). *J. Herpetol.* 14(1): 101.
168. San Miguel Ayanz, A. 1983. Contribución al estudio de la alimentación del arrendajo (Garrulus glandarius L. 1758) en España. *Bol. Est. Centr. Ecología* 12(23): 77-81.
169. Schipper, W.J.A. 1973. A comparison of prey selection in sympatric harriers (Circus) in Western Europe. *Le Gerfaut* 63: 17-120.
170. Schlee, M.A. 1984. Blackbirds eating large food items. *British Birds* 77(2): 74-75.
171. Schmitz, L. 1985. Consommation de tritons alpestres (Triturus alpestris) par le grebe castagneux (Tachybaptus ruficollis). *Aves* 22(3): 213.
172. Schrubbs, M. 1980. Farming influences on the food and hunting of the kestrels. *Bird Study* 27(2): 109-115.
173. Simeon, D., and J.L. Wilhelm. 1988. Essai sur l'alimentation annuelle de l'aigle de Bonelli Hieraaetus fasciatus en Provence. *Alauda* 56(3): 226-237.
174. Smal, C.M. 1987. The diet of the barn owl Tyto alba in Southern Ireland, with reference to a recently introduced prey species - the bank vole Clethrionomys glareolus. *Bird Study* 34(2): 113-125.
175. Smeenk, C. 1972. Ökologische vergleiche zwischen waldkauz Strix aluco und waldohreule Asio otus. *Ardea* 60(1-2): 1-71.
176. Straeten, E. Van der, and R. Asselberg. 1973. Het voedsel van de Kerkuil Tyto alba in België. *Le Gerfaut* 63(1-2): 149-159.
177. Suetens, W., and P. Van Groenendael. 1975. Observaciones en 1975 en un par de nidos extremeños de Elanio azul. *Ardeola* 22: 113-123.
178. Suetens, W., and P. Van Groenendael. 1977. Nidification de l'elanion blanc (Elanus caeruleus) en Espagne. *Le Gerfaut* 67(1-2): 54-72.
179. Taylor, D.W. 1976. Greenshank eating large frog. *British Birds* 69(10): 409.

180. Thiollay, J.M. 1963. Notes sur le régime alimentaire du Faucon crécerelle Falco tinnunculus en hiver. Nos oiseaux 27(2): 71-73.
181. Thiollay, J.M. 1966. Essai sur les rapaces du midi de La France. Distribution-Ecologie. Tentative de dénombrement. Alauda 34(3): 210-227.
182. Thiollay, J.M. 1967. Essai sur les rapaces du midi de La France. Distribution-Ecologie. Tentative de dénombrement. Aigle Royal, Aquila chrysaetos chrysaetos (L.). Alauda 35(2): 140-150.
183. Thiollay, J.M. 1968. Essai sur les rapaces du midi de La France. Distribution-Ecologie. Tentative de dénombrement. Circaète Jean le Blanc, Circaetus gallicus gallicus (Gmelin). Alauda 36(3): 179-189.
184. Thiollay, J.M. 1968. Essai sur les rapaces du midi de La France. Distribution-Ecologie. Tentative de dénombrement. Hibou grand duc, Bubo bubo bubo L. Alauda 37(1): 15-27.
185. Thiollay, J.M. 1968. Le régime alimentaire de nos rapaces: quelques analyses françaises. Nos oiseaux 29(10): 249-269.
186. Triplet, P. 1982. Note sur le cigogne blanche Ciconia ciconia en hiver. L'oiseau et R.F.O. 52(4): 372-373.
187. Turrian, F. 1987. La Bergeronnette grise, Motacilla alba, prédatrice de la grenouille rousse, Rana temporaria. Nos oiseaux 39(3): 137.
188. Uttendoerfen, O. 1952. Documents sur le Faucon crécerelle Falco tinnunculus L. Nos oiseaux 21(9): 217-223.
189. Valverde, J. A. 1956. Essai sur l'aigrette garzette (Egretta g. garzetta) en France. Alauda 24(1): 1-36.
190. Valverde, J. A. 1960. Vertebrados de las Marismas del Guadalquivir. (Introducción a su estudio ecológico). Archiv. Inst. Aclimatación 9: 1-168.
191. Valverde, J.A. 1967. Estructura de una comunidad de Vertebrados terrestres. Monogr. Est. Biol. Doñana 1. C.S.I.C.; Madrid.
192. Vargas, J.M., A. Antúnez, and M. Blasco. 1977. Comportamiento reproductivo y alimentario de la pagaza piconegra (Gelochelidon nilotica) en la laguna de Fuentepiedra de Málaga. Ardeola 24: 227-231.
193. Vargas, J.M., A. Antúnez, and M. Blasco. 1983. Los Vertebrados de la laguna de Fuentepiedra (Málaga). Monogr. I.C.O.N.A. 28. Minist. Agricult.; Madrid.
194. Vargas, J.M., E. Miguel, and M. Blasco. 1980. Estudio estacional y comparativo del régimen alimenticio de Tyto alba Scopoli en Fuentepiedra de Málaga y el Padul de Granada (España). Misc. Zool. 6: 95-102.
195. Vaucher, C.A. 1971. Notes sur l'éthologie de L'Aigle de Bonelli, Hieraetus fasciatus. Nos oiseaux 31(5-6): 101-111.

196. Veiga, J.P. 1978. Alimentación y relaciones tróficas entre la lechuza común (Tyto alba) y el buho chico (Asio otus) en la Sierra de Guadarrama (España). *Ardeola* 25: 113-142.
197. Veiga, J.P. 1985. Ecología de las rapaces de un ecosistema mediterráneo de montaña. Aproximación a su estructura comunitaria. Tesis Doctoral. Univ. Complutense Madrid.
198. Vericad, J.R., A. Escarré, and E. Rodríguez. 1976. Datos sobre la dieta de Tyto alba y Bubo bubo (SE de Iberia). *Mediterránea* 1: 47-59.
199. Vermot, M. 1980. Capture et ingurgitation d'un vertebre par une enchasse blanche, Himantopus himantopus. *Nos oiseaux* 35(6): 289.
200. Village, A. 1982. The diet of kestrels in relation to vole abundance. *Bird Study* 29(2): 129-138.
201. Villarán, A., and L.T. Medina. 1983. Alimentación del cárabo (Strix aluco L. 1758) en España. *Alytes* 1(1): 291-306.
202. Voisin, C. 1978. Utilisation des zones humides du Delta Rhodanien par les Ardeides. *L'oiseau et R.F.O.* 48(4): 329-380.
203. Walters Davies, P., and P.E. Davies. 1973. The ecology and conservation of the red kite in Wales. *British Birds* 66(6): 241-270.
204. Yalden, D.W. 1985. Dietary separation of owls in the Peak District. *Bird Study* 32(2): 122-131.
205. Yalden, D.W., and A.B. Warburton. 1979. The diet of the kestrel in the Lake District. *Bird Study* 26(3): 163-170.










0  
66  
P7

SEX DETERMINATION IN REPTILES:  
SUMMARY OF EFFECTS OF CONSTANT TEMPERATURES OF  
INCUBATION ON SEX RATIOS OF OFFSPRING




GARY L. PAUKSTIS

&

FREDRIC J. JANZEN

Department of Ecology and Evolution  
University of Chicago



SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
NO. 83

1990

**SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE**

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with the Division of Amphibians and Reptiles.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions on preparation and submission. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington DC 20560 USA. Please include a self-addressed mailing label with requests.

## INTRODUCTION

The phenomenon of environmental sex determination (ESD) in reptiles has been highly publicized in recent years. However, the underlying mechanism(s) that control this process are still poorly known. Additionally, the distribution of ESD within the Reptilia is poorly known, with only 93 of the approximately 6500 species of extant reptiles having been examined for the presence of ESD (Janzen and Paukstis, *Quart. Rev. Biol.*).

This paper provides a summary of much of the research conducted on ESD in reptiles from laboratory studies that have employed constant temperatures of incubation (Table 1). Table 1 is an extension of and appendix to a review article on ESD in reptiles (Janzen and Paukstis, *Quart. Rev. Biol.*). As a result of the recent proliferation of publications concerning various aspects of ESD in reptiles, this table originated as a tool to provide an overview of laboratory results that were currently available. As the data accumulated, we were impressed not by their consistency, but by the amount of variability that existed among different studies. Although each of these studies individually provided important new data on ESD, when many papers were viewed simultaneously it became very difficult to compare results.

The potential sources of this variability in sex ratios among different studies are many but, in general, fall into two broad categories--biological and artifactual. Among biological sources of variability are such factors as inter- and intrapopulational genetic differences, nongenetic maternal influences, and different regulatory mechanisms that may vary taxonomically within the Reptilia. Artifacts (nonbiological and experimental) include differences in experimental design and implementation (e.g., how closely temperature was monitored or regulated, randomization of eggs across experimental treatments, small sample sizes) and different techniques in sexing hatchlings (e.g., presence/absence of oviducts, histological confirmation).

Another factor that may reduce the value of results from some studies is the manner in which the results are presented (e.g., actual sample sizes as opposed to just the sex ratios of hatchlings, data on embryonic mortality). An additional problem is inconsistency and lack of definition of terminology. For example, what is the actual relationship between the morphology of a given gonad and the terminology used to describe it in hatchlings from different studies, when these hatchlings have been described as "hermaphrodites", "intersexes", "unsexable", or "unsexed"? Even though the answer to this question may be obvious, unless the terminology in each of these papers has been adequately defined, it is difficult to ascertain precisely how the gonads of these hatchlings may or may not differ. To properly understand ESD in reptiles, it is very important to differentiate between these biological and artificial/experimental sources of variation and to strive to minimize those sources of variation that may mask actual biological effects.

In this paper we provide a compilation of much of the published empirical research on ESD in reptiles. Specifically, we summarize information on incubation temperatures, sex ratios, and sample sizes from laboratory studies that have used constant temperatures of incubation. Comments are provided in those instances where they may lend insight into variability of sex ratios or to denote information that may be of particular interest. The arrangement of the major taxa used in Table 1 follows that of Janzen and Paukstis (*Quart. Rev. Biol.*). Families are listed alphabetically within the major taxa and genera are presented alphabetically within families.

We thank E. D. Brodie, III, L. E. Brown, J. J. Bull, S. O'Steen, P. A. Verrell, and M. J. Wade for support and discussion during the preparation of this manuscript. This work has been supported in part by an NIH Pre-Doctoral Training Grant in Genetics and Regulation (GM-07197) and by an NSF Doctoral Dissertation Improvement Grant (BSR-8914686) to FJJ.

Table 1. SUMMARY OF OFFSPRING SEX RATIOS FROM STUDIES USING CONSTANT TEMPERATURES OF INCUBATION

The number of males and females presented in this table were, in some cases, calculated from percentages provided in the sources indicated. In other cases, percentages were calculated from sample sizes. Question marks (?) denote data that were not presented in the source.

| Taxa                           | Temp (C) | <u>Males</u><br># ♂♂ (%) | <u>Females</u><br># ♀♀ (%) | # eggs/#<br>clutches | Comments  | Source |
|--------------------------------|----------|--------------------------|----------------------------|----------------------|---|--------|
| <b>TESTUDINES</b>              |          |                          |                            |                      |   |        |
| <b>CRYPTODIRA</b>              |          |                          |                            |                      |   |        |
| <b>CARETTOCHELYIDAE</b>        |          |                          |                            |                      |   |        |
| <i>Carettochelys insculpta</i> |          |                          |                            |                      |   |        |
|                                | 28       | 12(100)                  | 0( 0)                      | 12/3                 | Eggs in 2nd group at 32° were collected late in development | 98     |
|                                | 30       | 20(100)                  | 0( 0)                      | 24/5                 |   |        |
|                                | 32       | 0( 0)                    | 9(100)                     | 12/3                 |   |        |
|                                | 32       | 23( 50)                  | 23( 50)                    | 46/5                 |   |        |
| <b>CHELONIIDAE</b>             |          |                          |                            |                      |   |        |
| <i>Caretta caretta</i>         |          |                          |                            |                      |   |        |
|                                | 26       | 20(100)                  | 0( 0)                      | 20/5                 | 89% hatching success  | 38     |
|                                | 32       | 0( 0)                    | 20(100)                    | 20/5                 | 83% hatching success  |        |
|                                | 25       | ?(100)                   | 0( 0)                      | 10/1                 |   | 53     |
|                                | 26       | ?(100)                   | 0( 0)                      | 10/1                 |   |        |
|                                | 27.5     | ?( 80)                   | ?( 20)                     | 10/1                 |   |        |
|                                | 29       | ?( 50)                   | ?( 50)                     | 10/1                 |   |        |
|                                | 30.5     | 0( 0)                    | ?(100)                     | 10/1                 |   |        |
|                                | 32       | 0( 0)                    | ?(100)                     | 10/1                 |   |        |
|                                | 25       | 8(100)                   | 0( 0)                      | 10/1                 | 2 unsexable   |        |
|                                | 25       | 7(100)                   | 0( 0)                      | 10/1                 |   |        |
|                                | 26       | 16(100)                  | 0( 0)                      | 16/2                 |   |        |
|                                | 26       | 6( 67)                   | 3( 33)                     | 10/1                 |   |        |
|                                | 26       | 10(100)                  | 0( 0)                      | 10/1                 |   |        |
|                                | 26       | 7( 88)                   | 1( 12)                     | 10/1                 | 1 unsexable   |        |
|                                | 27       | 8( 80)                   | 2( 20)                     | 10/1                 |   |        |
|                                | 27       | 6( 86)                   | 1( 14)                     | 10/1                 | 2 unsexable   |        |
|                                | 27.5     | 5( 56)                   | 4( 44)                     | 10/1                 |   |        |
|                                | 27.5     | 3( 30)                   | 7( 70)                     | 10/1                 |   |        |
|                                | 27.5     | 8( 80)                   | 2( 20)                     | 10/1                 |   |        |
|                                | 28       | 4( 44)                   | 5( 56)                     | 10/1                 |   |        |
|                                | 28       | 2(100)                   | 0( 0)                      | 10/1                 |   |        |
|                                | 29       | 1( 11)                   | 8( 89)                     | 10/1                 |   |        |

| <u>Taxa</u>           | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>     | <u>Source</u>   |
|-----------------------|-----------------|-----------------|-----------------|--------------------------|---------------------|-----------------|
|                       | 29              | 5( 56)          | 4( 44)          | 10/1                     |                     |                 |
|                       | 29              | 2( 20)          | 8( 80)          | 10/1                     |                     |                 |
|                       | 29              | 5( 50)          | 5( 50)          | 10/1                     |                     |                 |
|                       | 29              | 6( 60)          | 4( 40)          | 10/1                     |                     |                 |
|                       | 30              | 0( 0)           | 7(100)          | 10/1                     |                     |                 |
|                       | 30              | 3( 33)          | 6( 67)          | 10/1                     |                     |                 |
|                       | 30              | 2( 20)          | 8( 80)          | 10/1                     |                     |                 |
|                       | 30.5            | 0( 0)           | 1(100)          | 1/1                      |                     |                 |
|                       | 30.5            | 2( 33)          | 4( 67)          | 7/1                      |                     |                 |
|                       | 30.5            | 0( 0)           | 10(100)         | 10/1                     |                     |                 |
|                       | 31              | 0( 0)           | 8(100)          | 10/1                     | 1 unsexable         |                 |
|                       | 31              | 0( 0)           | 16(100)         | 20/2                     |                     |                 |
|                       | 32              | 0( 0)           | 19(100)         | 20/2                     |                     |                 |
|                       | 32              | 0( 0)           | 6(100)          | 10/1                     | 1 unsexable         |                 |
|                       | 27.5            | 6( 86)          | 1( 14)          | ?/2                      | North Carolina (NC) | 62              |
|                       | 28.0            | 26( 90)         | 3( 10)          | ?/2                      | NC                  |                 |
|                       | 28.5            | 22( 69)         | 10( 31)         | ?/2                      | NC                  |                 |
|                       | 28.5            | 9( 33)          | 18( 67)         | ?/2                      | Georgia (GA)        |                 |
|                       | 28.5            | 25( 74)         | 9( 26)          | ?/2                      | Florida (FL)        |                 |
|                       | 28.8            | 19( 79)         | 5( 21)          | ?/2                      | NC                  |                 |
|                       | 28.8            | 12( 34)         | 23( 66)         | ?/2                      | GA                  |                 |
|                       | 28.8            | 22( 65)         | 12( 35)         | ?/2                      | FL                  |                 |
|                       | 29.2            | 4( 50)          | 4( 50)          | ?/2                      | NC                  |                 |
|                       | 29.5            | 8( 33)          | 16( 67)         | ?/2                      | NC                  |                 |
|                       | 29.5            | 8( 20)          | 31( 78)         | ?/2                      | GA; 1 intersex      |                 |
|                       | 29.5            | 12( 35)         | 25( 65)         | ?/2                      | FL                  |                 |
|                       | 30.0            | 3( 37)          | 5( 63)          | ?/2                      | NC                  |                 |
|                       | 30.4            | 0( 0)           | 25(100)         | ?/2                      | GA                  |                 |
|                       | 30.4            | 2( 5)           | 36( 92)         | ?/2                      | FL; 1 intersex      |                 |
|                       | 30.5            | 0( 0)           | 23(100)         | ?/2                      | NC                  |                 |
|                       | 24              | 11(100)         | 0( 0)           | 23/3                     |                     | 104,105,<br>106 |
|                       | 26              | 24(100)         | 0( 0)           | 26/5                     |                     |                 |
|                       | 28              | 20(100)         | 0( 0)           | 26/5                     |                     |                 |
|                       | 30              | 5( 36)          | 9( 64)          | 15/1                     | Incubated in 1978   |                 |
|                       | 30              | 5( 56)          | 4( 44)          | 15/2                     | Incubated in 1979   |                 |
|                       | 30              | 4( 80)          | 1( 20)          | 6/2                      | Incubated in 1980   |                 |
|                       | 32              | 0( 0)           | 21(100)         | 26/5                     |                     |                 |
|                       | 34              | 0( 0)           | 7(100)          | 26/5                     |                     |                 |
| <i>Chelonia mydas</i> |                 |                 |                 |                          |                     |                 |
|                       | 26              | 16( 84)         | 0( 0)           | 20/1                     | 3 intersexes        | 57              |
|                       | 29              | 0( 0)           | 37( 90)         | 41/1                     | 4 intersexes        |                 |
|                       | 33              | 0( 0)           | 12( 86)         | 20/1                     | 2 intersexes        |                 |
|                       | 27.75           | 19( 68)         | 6( 21)          | 38/3                     | 3 intersexes        | 63              |
|                       | 28.1            | 17( 61)         | 10( 36)         | 37/3                     | 1 intersex          |                 |
|                       | 29.25           | 11( 35)         | 18( 58)         | 38/3                     | 2 intersexes        |                 |
|                       | 30.0            | 9( 43)          | 11( 52)         | 37/3                     | 1 intersex          |                 |

| <u>Taxa</u>                  | <u>Temp (C)</u> | <u>#♂♂ (%)</u> | <u>#♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>   | <u>Source</u> |
|------------------------------|-----------------|----------------|----------------|--------------------------|---|---------------|
|                              | 27.5            | 587(~44)       | 763(~56)       | ??                       | Numbers for each sex are estimates owing to imprecision in sex ratios given | 100           |
|                              | 27.5            | 1178(~27)      | 3181(~73)      | ??                       |   |               |
|                              | 27.5            | 23(~77)        | 7(~23)         | ??                       |   |               |
|                              | 27.5            | 1572(~67)      | 786(~33)       | ??                       |   |               |
|                              | 27.5            | 237(~67)       | 118(~33)       | ??                       |   |               |
|                              | 30.0            | 97(~ 1)        | 13661(~99)     | ??                       |   |               |
| <i>Lepidochelys olivacea</i> |                 |                |                |                          |   |               |
|                              | 26.5            | ?(100)         | 0( 0)          | ?/1                      |   | 24            |
|                              | 29.5            | 0( 0)          | ?(100)         | ?/1                      |   |               |
|                              | 31.5            | 0( 0)          | ?(100)         | ?/1                      |   |               |
|                              | 25              | 23(100)        | 0( 0)          | 50/3                     | 3 unsexable<br>5 unsexable  | 56            |
|                              | 28              | 30( 88)        | 1( 3)          | 50/3                     |   |               |
|                              | 30              | 12( 48)        | 8( 32)         | 50/3                     |   |               |
|                              | 32              | 0( 0)          | 23(100)        | 50/3                     |   |               |
|                              | 26.5            | ?(100)         | 0( 0)          | ?/>1                     |   | 60            |
|                              | 28.0            | ?(100)         | 0( 0)          | ?/>1                     |   |               |
|                              | 29.5            | ?( 40)         | ?( 60)         | ?/>1                     |   |               |
|                              | 30.0            | 0( 0)          | ?(100)         | ?/>1                     |   |               |
|                              | 31.0            | 0( 0)          | ?(100)         | ?/>1                     |   |               |
|                              | 31.5            | 0( 0)          | ?(100)         | ?/>1                     |   |               |
|                              | 27.5            | ?(100)         | 0( 0)          | ??                       | Both sexes produced   | 61            |
|                              | 29.5            | ?( ?)          | ?( ?)          | ??                       |   |               |
|                              | 31.5            | 0( 0)          | ?(100)         | ??                       |   |               |
|                              | 25.7            | 160( 98)       | 1( 1)          | ?/9                      | 3 intersexes  | 84            |
|                              | ~27.4           | 59( 98)        | 1( 2)          | ?/3                      |   |               |
| CHELYDRIDAE                  |                 |                |                |                          |   |               |
| <i>Chelydra serpentina</i>   |                 |                |                |                          |   |               |
|                              | 25              | 10(100)        | 0( 0)          | 10/?                     |   | 21            |
|                              | 30              | 0( 0)          | 11(100)        | 11/?                     |   |               |
|                              | 28.5            | ?( ?)          | ?( ?)          | ??                       | Both sexes produced<br>1 intersex   | 22            |
|                              | 31              | 0( 0)          | 4( 80)         | 5/?                      |   |               |
|                              | 26.0            | 7( 88)         | 1( 12)         | ?/2                      | 44 eggs incubated<br>in total for this<br>experiment                        | 23            |
|                              | 28.5            | 3( 23)         | 10( 77)        | ?/2                      |   |               |
|                              | 31.0            | 0( 0)          | 13(100)        | ?/2                      |   |               |
|                              | 21.5            | 2( 25)         | 6( 75)         | ??                       | Indiana<br>Tennessee<br>Indiana<br>Minnesota                                | 28            |
|                              | 21.5            | 0( 0)          | 3(100)         | ??                       |   |               |
|                              | 22.5            | 5( 50)         | 5( 50)         | ??                       |   |               |
|                              | 22.5            | 32( 94)        | 2( 6)          | ??                       |   |               |
|                              | 25              | 33( 92)        | 3( 8)          | 36/?                     |   | 33            |
|                              | 31              | 0( 0)          | 33(100)        | 33/?                     |   |               |

| <u>Taxa</u>                  | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>   | <u>Source</u>   |
|------------------------------|-----------------|-----------------|-----------------|--------------------------|---|---|
|                              | 26              | 36( 95)         | 2( 5)           | ?/?                      |   | 34  |
|                              | 23              | 54(100)         | 0( 0)           | 54/9                     |   | 42,43   |
|                              | 26              | 50(100)         | 0( 0)           | 50/9                     |   |   |
|                              | 29              | 0( 0)           | 63(100)         | 63/15                    |   | 69  |
|                              | 20              | 0( 0)           | 21(100)         | 85/5                     | Eggs in the two 20 <sup>o</sup> treatments were switched to 26 <sup>o</sup> after 88 days and 83 days, respectively | 102   |
|                              | 20              | 0( 0)           | 37(100)         | 66/2                     |   |   |
|                              | 22              | 19( 90)         | 2( 10)          | 21/3                     |   |   |
|                              | 24              | 18(100)         | 0( 0)           | 18/3                     |   |   |
|                              | 26              | 108(100)        | 0( 0)           | 132/16                   |   |   |
|                              | 26              | 79( 98)         | 2( 2)           | 91/7                     |   |   |
|                              | 28              | 17( 65)         | 9( 35)          | 27/3                     |   |   |
|                              | 30              | 0( 0)           | 5(100)          | 23/5                     |   |   |
|                              | 30              | 0( 0)           | 48(100)         | 72/6                     |   |   |
|                              | 30              | 0( 0)           | 34(100)         | 56/5                     |   |   |
|                              | 20              | 0( 0)           | 149(100)        | 245/?                    |   | Eggs in the 20 <sup>o</sup> treatment completed incubation at 26 <sup>o</sup> |
|                              | 26              | 373( 99)        | 3( 1)           | 431/?                    |   |   |
|                              | 30              | 0( 0)           | 142(100)        | 196/?                    |   |   |
| <i>Macrochelys temmincki</i> |                 |                 |                 |                          |   |   |
|                              | 25              | ?( 60)          | ?( 40)          | ?/?                      |   | 6   |
|                              | 31              | 0( 0)           | ?(100)          | ?/?                      |   |   |
|                              | 22.5            | 2( 11)          | 16( 89)         | ?/?                      |   | 28  |
|                              | 25              | 9( 69)          | 4( 31)          | ?/?                      |   |   |
|                              | 27              | 10( 71)         | 4( 29)          | ?/?                      |   |   |
|                              | 30              | 0( 0)           | 11(100)         | ?/?                      |   |   |
| DERMOCHELYIDAE               |                 |                 |                 |                          |   |   |
| <i>Dermochelys coriacea</i>  |                 |                 |                 |                          |   |   |
|                              | 27.4            | 50(100)         | 0( 0)           | 50/5                     | Temps measured every 2-5 days at 0700 & 1800  | 25  |
|                              | 28.1            | 50(100)         | 0( 0)           | 50/5                     |   |   |
|                              | 27              | ?(100)          | 0( 0)           | ~25/1                    |   | 52  |
|                              | ~28             | ?(100)          | 0( 0)           | ~25/1                    |   |   |
|                              | ~29             | ?(100)          | 0( 0)           | ~25/1                    |   |   |
|                              | 31              | 0( 0)           | 40(100)         | 40/?                     |   |   |
|                              | 27              | 33(100)         | 0( 0)           | 75/2                     |   | 79  |
|                              | 27.25           | 5(100)          | 0( 0)           | 11/1                     |   |   |
|                              | 28              | 4(100)          | 0( 0)           | 38/2                     |   |   |
|                              | 28.25           | 1(100)          | 0( 0)           | 10/1                     |   |   |
|                              | 28.75           | 15(100)         | 0( 0)           | 51/2                     |   |   |
|                              | 29.75           | 0( 0)           | 4(100)          | 11/1                     |   |   |
|                              | 30.5            | 0( 0)           | 18(100)         | 59/3                     |   |   |
|                              | 32              | 0( 0)           | 3(100)          | 35/2                     |   |   |



| <u>Taxa</u>                     | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u> | <u>Source</u> |
|---------------------------------|-----------------|-----------------|-----------------|--------------------------|-----------------|---------------|
|                                 | 29.25           | 9(100)          | 0( 0)           | 140/4                    |                 | 80            |
|                                 | 29.5            | 1( 25)          | 3( 75)          | 16/1                     |                 |               |
|                                 | 29.5            | 12( 86)         | 2( 14)          | 16/1                     |                 |               |
|                                 | 29.75           | 0( 0)           | 32(100)         | 68/2                     |                 |               |
| <b>EMYDIDAE</b>                 |                 |                 |                 |                          |                 |               |
| <b>Batagurinae</b>              |                 |                 |                 |                          |                 |               |
| <i>Chinemys reevesii</i>        | ~25             | 18(100)         | 0( 0)           | 25/?                     |                 | 40            |
|                                 | 32              | 0( 0)           | 12( 92)         | 25/?                     | 1 intersex      |               |
| <i>Mauremys mutica</i>          | 25              | 3( 75)          | 1( 25)          | ?/?                      |                 | 28            |
|                                 | 30              | 0( 0)           | 9(100)          | ?/?                      |                 |               |
| <i>Melanochelys trijuga</i>     | 23.8            | 0( 0)           | 2(100)          | ?/?                      |                 | 28            |
|                                 | 25              | 7( 23)          | 23( 77)         | ?/?                      |                 |               |
|                                 | 27              | 15( 56)         | 12( 44)         | ?/?                      |                 |               |
|                                 | 30              | 1( 3)           | 31( 97)         | ?/?                      |                 |               |
| <i>Rhinoclemmys areolata</i>    | 25              | 6(100)          | 0( 0)           | ?/?                      |                 | 28            |
|                                 | 30              | 0( 0)           | 6(100)          | ?/?                      |                 |               |
| <i>Rhinoclemmys pulcherrima</i> | 25              | 14(100)         | 0( 0)           | ?/?                      |                 | 28            |
|                                 | 30              | 2( 25)          | 6( 75)          | ?/?                      |                 |               |
| <b>Emydinae</b>                 |                 |                 |                 |                          |                 |               |
| <i>Chrysemys picta</i>          | 25              | 81(100)         | 0( 0)           | 102/?                    |                 | 11            |
|                                 | 30.5            | 0( 0)           | 81(100)         | 101/?                    |                 |               |
|                                 | 28.0            | 1( 2)           | 40( 98)         | 41/>10                   | Tennessee (TN)  | 13            |
|                                 | 28.0            | 92( 98)         | 2( 2)           | 94/>25                   | Wisconsin (WI)  |               |
|                                 | 28.3            | 1( 9)           | 10( 91)         | 11/>10                   | TN              |               |
|                                 | 29.0            | 0( 0)           | 12(100)         | 12/>10                   | TN              |               |
|                                 | 29.0            | 24( 63)         | 14( 37)         | 38/>25                   | WI              |               |
|                                 | 29.5            | 0( 0)           | 5(100)          | 5/>10                    | TN              |               |
|                                 | 29.5            | 0( 0)           | 7(100)          | 7/>25                    | WI              |               |
|                                 | 30.0            | 0( 0)           | 16(100)         | 16/>10                   | TN              |               |
|                                 | 30.0            | 0( 0)           | 56(100)         | 56/>25                   | WI              |               |
|                                 | 30.6            | 0( 0)           | 14(100)         | 14/>10                   | TN              |               |
|                                 | 30.6            | 0( 0)           | 22(100)         | 22/>25                   | WI              |               |
|                                 | 28.5            | 0( 0)           | ?(100)          | ?/?                      |                 | 22            |
|                                 | 31              | 0( 0)           | ?(100)          | ?/?                      |                 |               |

| <u>Taxa</u>              | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>       | <u>Source</u> |
|--------------------------|-----------------|-----------------|-----------------|--------------------------|-----------------------|---------------|
|                          | 26.0            | 7(100)          | 0( 0)           | ?/7                      |                       | 23            |
|                          | 28.5            | 0( 0)           | 16(100)         | ?/7                      |                       |               |
|                          | 31.0            | 0( 0)           | 18(100)         | ?/7                      |                       |               |
|                          | 21.5            | 102(100)        | 0( 0)           | ?/?                      | some feminization     | 28            |
|                          | 22.5            | 10(100)         | 0( 0)           | ?/?                      |                       |               |
|                          | 25              | 83(100)         | 0( 0)           | ?/?                      |                       |               |
|                          | 27              | 33(100)         | 0( 0)           | ?/?                      |                       |               |
|                          | 30              | 0( 0)           | 78(100)         | ?/?                      |                       |               |
|                          | 25              | 23(100)         | 0( 0)           | 23/?                     |                       | 33            |
|                          | 31              | 0( 0)           | 28(100)         | 28/?                     |                       |               |
|                          | 26.5            | 19(100)         | 0( 0)           | 19/?                     |                       | 36            |
|                          | 26.5            | 28( 80)         | 7( 20)          | 35/?                     |                       |               |
|                          | 27              | 21(100)         | 0( 0)           | 30/?                     | 9 unsexable           |               |
|                          | 27              | 6( 43)          | 8( 57)          | 18/?                     | 4 unsexable           |               |
|                          | 28.5            | 4( 19)          | 17( 81)         | 21/?                     |                       |               |
|                          | 28.5            | 3( 14)          | 18( 86)         | 21/?                     |                       |               |
|                          | 28.5            | 7( 78)          | 2( 22)          | 9/?                      |                       |               |
|                          | 28.5            | 6( 75)          | 2( 25)          | 8/?                      |                       |               |
|                          | 30.5            | 0( 0)           | 37(100)         | 37/?                     |                       |               |
|                          | 32              | 0( 0)           | 14(100)         | 15/?                     | 1 unsexable           |               |
|                          | 22              | 6( 35)          | 11( 65)         | 40/21                    | 1 unsexable           | 37            |
|                          | 27              | 21(100)         | 0( 0)           | 31/21                    | 9 unsexable           |               |
|                          | 32              | 0( 0)           | 14(100)         | 31/21                    | 1 unsexable           |               |
|                          | 25.7            | 41(100)         | 0( 0)           | 45/24                    | -150 and -1100 kPa    | 68            |
|                          | 26.7            | 38(100)         | 0( 0)           | 45/24                    | -150 and -1100 kPa    |               |
|                          | 27.7            | 29(100)         | 0( 0)           | 45/24                    | -150 and -1100 kPa    |               |
|                          | 28.7            | 13( 68)         | 6( 32)          | 21/12                    | -150 kPa              |               |
|                          | 28.7            | 7( 50)          | 6( 43)          | 24/12                    | -1100 kPa, 1 intersex |               |
|                          | 20              | 3( 50)          | 3( 50)          | 21/19                    |                       | 83            |
|                          | 22              | 14(100)         | 0( 0)           | 20/19                    |                       |               |
|                          | 24              | 17(100)         | 0( 0)           | 21/19                    |                       |               |
|                          | 26              | 18(100)         | 0( 0)           | 21/19                    |                       |               |
|                          | 28              | 3( 19)          | 13( 81)         | 21/19                    |                       |               |
|                          | 30              | 0( 0)           | 19(100)         | 21/19                    |                       |               |
|                          | 32              | 0( 0)           | 17(100)         | 21/19                    |                       |               |
| <i>Clemmys guttata</i>   | 22.5            | 10( 91)         | 1( 9)           | ?/?                      |                       | 28            |
|                          | 25              | 14( 70)         | 6( 30)          | ?/?                      |                       |               |
|                          | 27              | 12( 92)         | 1( 8)           | ?/?                      |                       |               |
|                          | 30              | 0( 0)           | 19(100)         | ?/?                      |                       |               |
| <i>Clemmys insculpta</i> | 25              | 6( 33)          | 12( 67)         | 18/6                     |                       | 10            |
|                          | 30              | 7( 39)          | 11( 61)         | 18/6                     |                       |               |

| <u>Taxa</u>                    | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u> | <u>Source</u> |
|--------------------------------|-----------------|-----------------|-----------------|--------------------------|-----------------|---------------|
|                                | 22.5            | 15( 44)         | 19( 56)         | ???                      |                 | 28            |
|                                | 25              | 19( 44)         | 24( 56)         | ???                      |                 |               |
|                                | 27              | 15( 44)         | 19( 56)         | ???                      |                 |               |
|                                | 30              | 24( 53)         | 21( 47)         | ???                      |                 |               |
| <i>Clemmys muhlenbergii</i>    | 25              | 1( 33)          | 2( 67)          | ???                      |                 | 28            |
| <i>Deirochelys reticularia</i> | 25              | 16(100)         | 0( 0)           | ???                      |                 | 28            |
|                                | 30              | 2( 11)          | 17( 89)         | ???                      |                 |               |
| <i>Emydoidea blandingii</i>    | 22.5            | 40(100)         | 0( 0)           | ???                      |                 | 28            |
|                                | 25              | 57( 97)         | 2( 3)           | ???                      |                 |               |
|                                | 30              | 0( 0)           | 63(100)         | ???                      |                 |               |
|                                | 26.5            | 10(100)         | 0( 0)           | ?/6                      |                 | 35            |
|                                | 31.0            | 0( 0)           | 10(100)         | ?/6                      |                 |               |
| <i>Emys orbicularis</i>        | 30              | 1( 4)           | 23( 96)         | 24/?                     |                 | 71            |
|                                | 25              | 40(100)         | 0( 0)           | 40/?                     |                 | 72            |
|                                | 29.5            | 0( 0)           | 11(100)         | 11/?                     |                 |               |
|                                | 27.5            | 25(100)         | 0( 0)           | 25/?                     |                 | 73            |
|                                | 25              | 76(100)         | 0( 0)           | 76/?                     |                 | 74            |
|                                | 27.5            | 25(100)         | 0( 0)           | 25/?                     |                 |               |
|                                | 29.5            | 0( 0)           | 117(100)        | 117/?                    |                 |               |
|                                | 27.75           | 30(100)         | 0( 0)           | 30/?                     |                 | 75            |
|                                | 28.25           | 19( 95)         | 1( 5)           | 20/?                     |                 |               |
|                                | 28.75           | 8( 42)          | 11( 58)         | 19/?                     |                 |               |
|                                | 29.25           | 1( 3)           | 29( 94)         | 31/?                     | 1 intersex      |               |
|                                | 27.75           | 23( 77)         | 0( 0)           | 30/?                     | 7 intersexes    | 76            |
|                                | 28.25           | 20( 54)         | 4( 11)          | 37/?                     | 13 intersexes   |               |
|                                | 28.75           | 6( 18)          | 13( 39)         | 33/?                     | 14 intersexes   |               |
|                                | 29.25           | 0( 0)           | 29( 94)         | 31/?                     | 2 intersexes    |               |
|                                | 29.75           | 0( 0)           | 54(100)         | 54/?                     |                 |               |
|                                | 18              | 8(100)          | 0( 0)           | 8/?                      |                 | 77            |
|                                | 19.5            | 6(100)          | 0( 0)           | 6/?                      |                 |               |
|                                | 35              | 0( 0)           | 10(100)         | 10/?                     |                 |               |
|                                | 25.5            | 149(100)        | 0( 0)           | 149/?                    |                 | 107           |
|                                | 28.75           | 6( 16)          | 30( 81)         | 37/?                     | 1 intersex      |               |
|                                | 30.25           | 0( 0)           | 127(100)        | 127/?                    |                 |               |

| <u>Taxa</u>                   | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>    | <u>Source</u> |
|-------------------------------|-----------------|-----------------|-----------------|--------------------------|--------------------|---------------|
|                               | 28.5            | 1( 11)          | 8( 89)          | 9/1                      |                    | 108           |
|                               | 28.5            | 0( 0)           | 10(100)         | 10/1                     |                    |               |
|                               | 28.5            | 0( 0)           | 5( 63)          | 8/1                      | 3 intersexes       |               |
|                               | 28.5            | 4( 80)          | 1( 20)          | 6/1                      | 1 embryo died      |               |
|                               | 28.5            | 4( 67)          | 1( 17)          | 6/1                      | 1 intersex         |               |
|                               | 28.5            | 4( 44)          | 1( 11)          | 9/1                      | 4 intersexes       |               |
|                               | 28.5            | 2( 18)          | 9( 82)          | 11/1                     |                    |               |
|                               | 28.5            | 1( 14)          | 6( 86)          | 7/1                      |                    |               |
|                               | 28.5            | 5( 83)          | 1( 17)          | 7/1                      | 1 egg unfertilized |               |
|                               | 28.5            | 1( 10)          | 6( 60)          | 10/1                     | 3 intersexes       |               |
| <i>Graptemys barbouri</i>     |                 |                 |                 |                          |                    |               |
|                               | 25              | 9(100)          | 0( 0)           | ??/?                     |                    | 28            |
|                               | 30              | 0( 0)           | 9(100)          | ??/?                     |                    |               |
| <i>Graptemys geographica</i>  |                 |                 |                 |                          |                    |               |
|                               | 25              | 98(100)         | 0( 0)           | 122/?                    |                    | 11            |
|                               | 30.5            | 0( 0)           | 88(100)         | 119/?                    |                    |               |
|                               | 28.0            | 26(100)         | 0( 0)           | 26/7                     |                    | 13            |
|                               | 29.0            | 2( 33)          | 4( 67)          | 6/7                      |                    |               |
|                               | 30.0            | 0( 0)           | 28(100)         | 28/7                     |                    |               |
|                               | 22.5            | 14(100)         | 0( 0)           | ??/?                     |                    | 28            |
|                               | 25              | 33(100)         | 0( 0)           | ??/?                     |                    |               |
|                               | 27              | 22(100)         | 0( 0)           | ??/?                     |                    |               |
|                               | 30              | 0( 0)           | 44(100)         | ??/?                     |                    |               |
|                               | 33              | 0( 0)           | 3(100)          | ??/?                     |                    |               |
| <i>Graptemys kohyii</i>       |                 |                 |                 |                          |                    |               |
|                               | 25              | 151(100)        | 0( 0)           | ??/?                     |                    | 28            |
|                               | 30              | 0( 0)           | 153(100)        | ??/?                     |                    |               |
| <i>Graptemys nigrinoda</i>    |                 |                 |                 |                          |                    |               |
|                               | 25              | 6(100)          | 0( 0)           | ??/?                     |                    | 28            |
|                               | 30              | 0( 0)           | 7(100)          | ??/?                     |                    |               |
| <i>Graptemys ouachitensis</i> |                 |                 |                 |                          |                    |               |
|                               | 25              | 210(100)        | 0( 0)           | 233/?                    |                    | 11            |
|                               | 30.5            | 0( 0)           | 211(100)        | 237/?                    |                    |               |
|                               | 29.25           | 3( 30)          | 7( 70)          | 10/1                     |                    | 12            |
|                               | 29.25           | 7( 78)          | 2( 22)          | 10/1                     |                    |               |
|                               | 29.25           | 4( 40)          | 6( 60)          | 10/1                     |                    |               |
|                               | 29.25           | 4( 50)          | 3( 38)          | 10/1                     | 1 intersex         |               |
|                               | 29.25           | 2( 22)          | 7( 78)          | 10/1                     |                    |               |
|                               | 29.25           | 5( 50)          | 4( 40)          | 10/1                     | 1 intersex         |               |
|                               | 29.25           | 0( 0)           | 9(100)          | 10/1                     |                    |               |
|                               | 29.25           | 7( 78)          | 2( 22)          | 10/1                     |                    |               |
|                               | 29.25           | 0( 0)           | 10(100)         | 10/1                     |                    |               |
|                               | 29.25           | 1( 10)          | 9( 90)          | 10/1                     |                    |               |

| <u>Taxa</u>                       | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>               | <u>Source</u> |
|-----------------------------------|-----------------|-----------------|-----------------|--------------------------|-------------------------------|---------------|
|                                   | 29.25           | 9( 90)          | 0( 0)           | 10/1                     | 1 intersex                    |               |
|                                   | 29.25           | 2( 22)          | 7( 78)          | 10/1                     |                               |               |
|                                   | 29.25           | 3( 30)          | 7( 70)          | 10/1                     |                               |               |
|                                   | 29.25           | 4( 44)          | 5( 56)          | 10/1                     |                               |               |
|                                   | 29.25           | 10(100)         | 0( 0)           | 10/1                     |                               |               |
|                                   | 29.25           | 4( 40)          | 6( 60)          | 10/1                     |                               |               |
|                                   | 29.25           | 4( 40)          | 5( 50)          | 10/1                     | 1 intersex                    |               |
|                                   | 29.25           | 1( 10)          | 9( 90)          | 10/1                     |                               |               |
|                                   | 29.25           | 3( 30)          | 6( 60)          | 10/1                     | 1 intersex                    |               |
|                                   | 29.25           | 5( 50)          | 5( 50)          | 10/1                     |                               |               |
|                                   | 28.0            | 93(100)         | 0( 0)           | 93/>25                   |                               | 13            |
|                                   | 29.0            | 53( 83)         | 11( 17)         | 64/>25                   |                               |               |
|                                   | 30.0            | 1( 1)           | 88( 99)         | 89/>25                   |                               |               |
|                                   | 25              | 69(100)         | 0( 0)           | ?/?                      |                               | 28            |
|                                   | 30              | 0( 0)           | 64(100)         | ?/?                      |                               |               |
| <i>Gratemys pseudogeographica</i> |                 |                 |                 |                          |                               |               |
|                                   | 25              | 173(100)        | 0( 0)           | 222/?                    |                               | 11            |
|                                   | 30.5            | 4( 3)           | 147( 97)        | 232/?                    |                               |               |
|                                   | 28.0            | 7(100)          | 0( 0)           | 7/7                      | Tennessee (TN)                | 13            |
|                                   | 28.0            | 70(100)         | 0( 0)           | 70/9                     | Wisconsin (WI)                |               |
|                                   | 28.3            | 24( 96)         | 1( 4)           | 25/7                     | TN                            |               |
|                                   | 28.3            | 14(100)         | 0( 0)           | 14/9                     | WI                            |               |
|                                   | 29.0            | 0( 0)           | 5(100)          | 5/7                      | TN                            |               |
|                                   | 29.0            | 22( 92)         | 2( 8)           | 24/9                     | WI                            |               |
|                                   | 29.3            | 13( 28)         | 34( 72)         | 47/7                     | TN                            |               |
|                                   | 29.3            | 33( 58)         | 24( 42)         | 57/9                     | WI                            |               |
|                                   | 29.5            | 4( 16)          | 21( 84)         | 25/7                     | TN                            |               |
|                                   | 29.5            | 5( 33)          | 10( 67)         | 15/9                     | WI                            |               |
|                                   | 30.0            | 0( 0)           | 5(100)          | 5/7                      | TN                            |               |
|                                   | 30.0            | 9( 11)          | 73( 89)         | 82/9                     | WI                            |               |
|                                   | 30.6            | 0( 0)           | 22(100)         | 22/7                     | TN                            |               |
|                                   | 30.6            | 0( 0)           | 17(100)         | 17/9                     | WI                            |               |
|                                   | 22.5            | 11(100)         | 0( 0)           | ?/?                      |                               | 28            |
|                                   | 25              | 16(100)         | 0( 0)           | ?/?                      |                               |               |
|                                   | 30              | 0( 0)           | 14(100)         | ?/?                      |                               |               |
|                                   | 33              | 0( 0)           | 11(100)         | ?/?                      |                               |               |
|                                   | 25              | 54(100)         | 0( 0)           | 66/17                    | Eggs from <i>ouachitensis</i> | 92            |
|                                   | 35              | 0( 0)           | 17(100)         | 70/17                    | and <i>pseudogeographica</i>  |               |
| <i>Gratemys pulchra</i>           |                 |                 |                 |                          |                               |               |
|                                   | 28.0            | 17(100)         | 0( 0)           | 17/>10                   |                               | 13            |
|                                   | 29.0            | 0( 0)           | 4(100)          | 4/>10                    |                               |               |
|                                   | 30.0            | 0( 0)           | 14(100)         | 14/>10                   |                               |               |

| <u>Taxa</u>                | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/#<br/>clutches</u> | <u>Comments</u> | <u>Source</u> |
|----------------------------|-----------------|-----------------|-----------------|------------------------------|-----------------|---------------|
| <i>Malaclemys terrapin</i> |                 |                 |                 |                              |                 |               |
|                            | 24              | 20(100)         | 0( 0)           | ??                           |                 | 28            |
|                            | 30              | 0( 0)           | 34(100)         | ??                           |                 |               |
|                            | 27              | 2( 67)          | 1( 33)          | 7/1                          | 1972            | 82            |
|                            | 27              | 35(100)         | 0( 0)           | 63/9                         | 1973-1975       |               |
|                            | 27              | 7( 88)          | 1( 12)          | 8/1                          | 1977            |               |
|                            | 27              | 8(100)          | 0( 0)           | 9/1                          | 1977            |               |
| <i>Pseudemys concinna</i>  |                 |                 |                 |                              |                 |               |
|                            | 22.5            | 13(100)         | 0( 0)           | ??                           |                 | 28            |
|                            | 25              | 52( 91)         | 5( 9)           | ??                           |                 |               |
|                            | 30              | 0( 0)           | 55(100)         | ??                           |                 |               |
| <i>Pseudemys floridana</i> |                 |                 |                 |                              |                 |               |
|                            | 25              | 4(100)          | 0( 0)           | ??                           |                 | 28            |
|                            | 30              | 0( 0)           | 4(100)          | ??                           |                 |               |
| <i>Terrapene carolina</i>  |                 |                 |                 |                              |                 |               |
|                            | 26.0            | 3( 50)          | 3( 50)          | ?/5                          |                 | 23            |
|                            | 28.5            | 2( 40)          | 3( 60)          | ?/5                          |                 |               |
|                            | 31.0            | 1( 14)          | 6( 86)          | ?/5                          |                 |               |
|                            | 21.5            | 13( 93)         | 1( 7)           | ??                           |                 | 28            |
|                            | 22.5            | 24( 73)         | 9( 27)          | ??                           |                 |               |
|                            | 25              | 73( 96)         | 3( 4)           | ??                           |                 |               |
|                            | 27              | 25( 81)         | 6( 19)          | ??                           |                 |               |
|                            | 30              | 0( 0)           | 84(100)         | ??                           |                 |               |
| <i>Terrapene ornata</i>    |                 |                 |                 |                              |                 |               |
|                            | 21.5            | 1(100)          | 0( 0)           | ??                           |                 | 28            |
|                            | 22.5            | 14(100)         | 0( 0)           | ??                           |                 |               |
|                            | 25              | 8(100)          | 0( 0)           | ??                           |                 |               |
|                            | 29              | 0( 0)           | 28(100)         | 31/9                         | 2 unsexed       | 70            |
| <i>Trachemys scripta</i>   |                 |                 |                 |                              |                 |               |
|                            | 28.0            | 21(100)         | 0( 0)           | 21/>10                       | Alabama (AL)    | 13            |
|                            | 28.3            | 33( 92)         | 3( 8)           | 36/>10                       | Tennessee (TN)  |               |
|                            | 29.0            | 6( 38)          | 10( 62)         | 16/>10                       | AL              |               |
|                            | 29.5            | 12( 30)         | 28( 70)         | 40/>10                       | TN              |               |
|                            | 30.0            | 0( 0)           | 17(100)         | 17/>10                       | AL              |               |
|                            | 30.6            | 2( 5)           | 40( 95)         | 42/>10                       | TN              |               |
|                            | 21.5            | 3(100)          | 0( 0)           | ??                           |                 | 28            |
|                            | 22.5            | 23(100)         | 0( 0)           | ??                           |                 |               |
|                            | 25              | 21(100)         | 0( 0)           | ??                           |                 |               |
|                            | 27              | 3(100)          | 0( 0)           | ??                           |                 |               |
|                            | 30              | 0( 0)           | 20(100)         | ??                           |                 |               |

| <u>Taxa</u>                    | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>       | <u>Source</u> |
|--------------------------------|-----------------|-----------------|-----------------|--------------------------|-----------------------|---------------|
| <b>KINOSTERNIDAE</b>           |                 |                 |                 |                          |                       |               |
| <i>Kinosternon flavescens</i>  |                 |                 |                 |                          |                       |               |
|                                | 25              | 5( 33)          | 10( 67)         | ???                      |                       | 28            |
|                                | 27              | 13( 93)         | 1( 7)           | ???                      |                       |               |
|                                | 30              | 10( 91)         | 1( 9)           | ???                      |                       |               |
|                                | 32              | 0( 0)           | 8(100)          | ???                      |                       |               |
|                                | 25              | 11( 79)         | 3( 21)          | 29/10                    |                       | 94            |
|                                | 31              | 0( 0)           | 16(100)         | 27/10                    |                       |               |
|                                | 25              | ?( 79)          | ?( 21)          | ???                      | <i>K. flavescens?</i> | 6             |
|                                | 31              | 0( 0)           | ?(100)          | ???                      |                       |               |
| <i>Kinosternon leucostomum</i> |                 |                 |                 |                          |                       |               |
|                                | 22.5            | 1 or 2( 19)     | 6 or 7( 81)     | ???                      |                       | 28            |
|                                | 24              | 1(100)          | 0( 0)           | ???                      |                       |               |
|                                | 25              | 3( 75)          | 1( 25)          | ???                      |                       |               |
|                                | 27              | 0( 0)           | 6(100)          | ???                      |                       |               |
|                                | 30              | 0( 0)           | 9(100)          | ???                      |                       |               |
| <i>Kinosternon scorpioides</i> |                 |                 |                 |                          |                       |               |
|                                | 22.5            | 8( 22)          | 14( 78)         | ???                      |                       | 28            |
|                                | 24              | 25( 82)         | 6( 18)          | ???                      |                       |               |
|                                | 25              | 53( 81)         | 12( 19)         | ???                      |                       |               |
|                                | 27              | 23( 70)         | 10( 30)         | ???                      |                       |               |
|                                | 30              | 0( 0)           | 73(100)         | ???                      |                       |               |
| <i>Kinosternon subrubrum</i>   |                 |                 |                 |                          |                       |               |
|                                | 22.5            | 1( 17)          | 5( 83)          | ???                      |                       | 28            |
| <i>Sternotherus carinatus</i>  |                 |                 |                 |                          |                       |               |
|                                | 22.5            | 0( 0)           | 5(100)          | ???                      |                       | 28            |
|                                | 25              | 1( 20)          | 4( 80)          | ???                      |                       |               |
|                                | 27              | 6(100)          | 0( 0)           | ???                      |                       |               |
|                                | 30              | 0( 0)           | 6(100)          | ???                      |                       |               |
| <i>Sternotherus minor</i>      |                 |                 |                 |                          |                       |               |
|                                | 22.5            | 1( 4)           | 27( 96)         | ???                      |                       | 28            |
|                                | 24              | 1( 8)           | 11( 92)         | ???                      |                       |               |
|                                | 25              | 22( 76)         | 7( 24)          | ???                      |                       |               |
|                                | 27              | 1( 6)           | 17( 94)         | ???                      |                       |               |
|                                | 30              | 0( 0)           | 36(100)         | ???                      |                       |               |
|                                | 32              | 0( 0)           | 3(100)          | ???                      |                       |               |
| <i>Sternotherus odoratus</i>   |                 |                 |                 |                          |                       |               |
|                                | 21.5            | 0( 0)           | 14(100)         | ???                      |                       | 28            |
|                                | 22.5            | 0( 0)           | 59(100)         | ???                      |                       |               |
|                                | 23.8            | 8( 31)          | 18( 69)         | ???                      |                       |               |
|                                | 25              | 46( 94)         | 3( 6)           | ???                      |                       |               |
|                                | 27              | 6( 23)          | 20( 77)         | ???                      |                       |               |

| <u>Taxa</u>                    | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u> | <u>Source</u> |
|--------------------------------|-----------------|-----------------|-----------------|--------------------------|-----------------|---------------|
|                                | 30              | 0( 0)           | 51(100)         | ?/?                      |                 |               |
|                                | 23.5            | 6( 19)          | 26( 81)         | 34/20                    |                 | 94            |
|                                | 25              | 27( 82)         | 6( 18)          | 34/20                    |                 |               |
|                                | 28              | 2( 3)           | 68( 97)         | 92/58                    |                 |               |
|                                | 29.5            | 2( 3)           | 56( 97)         | 58/38                    |                 |               |
|                                | 30.5            | 1( 2)           | 40( 98)         | 69/58                    |                 |               |
| <b>STAUROTYPIDAE</b>           |                 |                 |                 |                          |                 |               |
| <i>Staurotypus salvinii</i>    |                 |                 |                 |                          |                 |               |
|                                | 22.5            | 1( 25)          | 4( 75)          | ?/?                      | from 1 female   | 28            |
|                                | 22.5            | 18(100)         | 0( 0)           | ?/?                      | from 3 females  |               |
|                                | 25              | 8( 53)          | 7( 47)          | ?/?                      | from 1 female   |               |
|                                | 25              | 19(100)         | 0( 0)           | ?/?                      | from 3 females  |               |
|                                | 27              | 3( 60)          | 2( 40)          | ?/?                      | from 1 female   |               |
|                                | 27              | 14(100)         | 0( 0)           | ?/?                      | from 3 females  |               |
|                                | 30              | 3( 33)          | 6( 67)          | ?/?                      | from 1 female   |               |
|                                | 30              | 7( 78)          | 2( 22)          | ?/?                      | from 3 females  |               |
| <i>Staurotypus triporcatus</i> |                 |                 |                 |                          |                 |               |
|                                | 22.5            | 11( 44)         | 14( 56)         | ?/?                      |                 | 28            |
|                                | 25              | 17( 55)         | 14( 45)         | ?/?                      |                 |               |
|                                | 27              | 12( 40)         | 18( 60)         | ?/?                      |                 |               |
|                                | 30              | 8( 53)          | 7( 47)          | ?/?                      |                 |               |
| <b>TESTUDINIDAE</b>            |                 |                 |                 |                          |                 |               |
| <i>Testudo graeca</i>          |                 |                 |                 |                          |                 |               |
|                                | 26.5            | 19(100)         | 0( 0)           | 19/?                     |                 | 71            |
|                                | 30              | 22( 96)         | 1( 4)           | 23/?                     |                 |               |
|                                | 31              | 0( 0)           | 20(100)         | 20/?                     |                 | 72            |
|                                | 33              | 0( 0)           | 20(100)         | 20/?                     |                 |               |
|                                | 26.5            | 19(100)         | 0( 0)           | 19/?                     |                 | 74            |
|                                | 29.5            | 37( 97)         | 1( 3)           | 38/?                     |                 |               |
|                                | 31.5            | 0( 0)           | 16(100)         | 16/?                     |                 |               |
| <i>Testudo hermanni</i>        |                 |                 |                 |                          |                 |               |
|                                | ~23.5           | 6(100)          | 0( 0)           | 6/1                      |                 | 26            |
| <b>TRIONYCHIDAE</b>            |                 |                 |                 |                          |                 |               |
| <i>Trionyx muticus</i>         |                 |                 |                 |                          |                 |               |
|                                | 27              | 23( 44)         | 29( 56)         | ?/?                      |                 | 28            |
|                                | 30              | 26( 49)         | 27( 51)         | ?/?                      |                 |               |
|                                | 33              | 30( 54)         | 25( 46)         | ?/?                      |                 |               |
| <i>Trionyx spiniferus</i>      |                 |                 |                 |                          |                 |               |
|                                | 31              | 7( 50)          | 7( 50)          | 51/?                     |                 | 9             |



| <u>Taxa</u>                  | <u>Temp (C)</u> | <u># ♂ (%)</u> | <u># ♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>              | <u>Source</u> |
|------------------------------|-----------------|----------------|----------------|--------------------------|------------------------------|---------------|
|                              | 25              | 33( 49)        | 34( 51)        | 83/?                     |                              | 11            |
|                              | 30.5            | 27( 53)        | 24( 47)        | 86/?                     |                              |               |
|                              | 23              | 7( 41)         | 10( 59)        | 68/?                     | 51 unsexed                   | 93            |
|                              | 25              | 34( 49)        | 35( 51)        | 86/?                     | 17 unsexed                   |               |
|                              | 28              | 29( 52)        | 27( 48)        | 69/?                     | 13 unsexed                   |               |
|                              | 30.5            | 28( 53)        | 25( 47)        | 89/?                     | 36 unsexed                   |               |
|                              | 33              | 21( 41)        | 30( 59)        | 66/?                     | 15 unsexed                   |               |
| PLEURODIRA                   |                 |                |                |                          |                              |               |
| CHELIDAE                     |                 |                |                |                          |                              |               |
| <i>Chelodina longicollis</i> |                 |                |                |                          |                              |               |
|                              | 24              | 7( 35)         | 13( 65)        | 25/15                    | 5 unsexed                    | 31            |
|                              | 26              | 4( 36)         | 7( 64)         | 13/13                    | 2 unsexed                    |               |
|                              | 28              | 4( 31)         | 9( 69)         | 13/13                    |                              |               |
|                              | 30              | 6( 35)         | 11( 65)        | 18/15                    | 1 unsexed                    |               |
|                              | 32              | 14( 64)        | 11( 36)        | 27/15                    | 1 unsexed                    |               |
| <i>Emydura macquarii</i>     |                 |                |                |                          |                              |               |
|                              | 20              | 1( 33)         | 2( 67)         | 6/1                      | Eggs at 20 <sup>o</sup> were | 85,86         |
|                              | ~25             | 13( 65)        | 7( 35)         | 24/8                     | switched to 30 <sup>o</sup>  |               |
|                              | 26              | 7( 37)         | 12( 63)        | 24/8                     | after 91 days                |               |
|                              | 28              | 10( 50)        | 10( 50)        | 24/8                     |                              |               |
|                              | 30              | 18( 78)        | 5( 22)         | 24/8                     |                              |               |
|                              | 30              | 58( 52)        | 53( 48)        | ?/?                      |                              |               |
|                              | 32              | 10( 56)        | 8( 44)         | 24/8                     |                              |               |
| <i>Emydura signata</i>       |                 |                |                |                          |                              |               |
|                              | 25              | 15( 45)        | 18( 55)        | 33/12                    |                              | 10            |
|                              | 28              | 3( 25)         | 9( 75)         | 12/12                    |                              |               |
|                              | 30              | 13( 65)        | 7( 35)         | 20/12                    |                              |               |
| PELOMEDUSIDAE                |                 |                |                |                          |                              |               |
| <i>Pelomedusa subrufa</i>    |                 |                |                |                          |                              |               |
|                              | 24              | 0( 0)          | 2(100)         | ?/?                      |                              | 28            |
|                              | 25              | 0( 0)          | 11(100)        | ?/?                      |                              |               |
|                              | 27              | 0( 0)          | 17(100)        | ?/?                      |                              |               |
|                              | 30              | 12( 71)        | 5( 29)         | ?/?                      |                              |               |
|                              | 33              | 0( 0)          | 9(100)         | ?/?                      |                              |               |
| <i>Pelusios castaneus</i>    |                 |                |                |                          |                              |               |
|                              | 25              | 0( 0)          | 11(100)        | ?/?                      |                              | 28            |
|                              | 27              | 0( 0)          | 3(100)         | ?/?                      |                              |               |
|                              | 30              | 14( 82)        | 3( 18)         | ?/?                      |                              |               |
|                              | 33              | 0( 0)          | 5(100)         | ?/?                      |                              |               |

| <u>Taxa</u>                       | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>     | <u>Source</u> |
|-----------------------------------|-----------------|-----------------|-----------------|--------------------------|---------------------|---------------|
| <b>CROCODYLIA</b>                 |                 |                 |                 |                          |                     |               |
| <b>ALLIGATORIDAE</b>              |                 |                 |                 |                          |                     |               |
| <i>Alligator mississippiensis</i> |                 |                 |                 |                          |                     |               |
|                                   | 32              | 12(100)         | 0( 0)           | 46/?                     |                     | 9             |
|                                   | 26              | 0( 0)           | 10(100)         | 50/13                    |                     | 29,30         |
|                                   | 28              | 0( 0)           | 96(100)         | 100/13                   |                     |               |
|                                   | 30              | 0( 0)           | 97(100)         | 100/13                   |                     |               |
|                                   | 32              | 13( 13)         | 85( 87)         | 100/13                   |                     |               |
|                                   | 34              | 94(100)         | 0( 0)           | 100/13                   |                     |               |
|                                   | 36              | 7(100)          | 0( 0)           | 50/13                    |                     |               |
|                                   | 29.4            | 0( 0)           | 90(100)         | 113/11                   |                     | 44,45         |
|                                   | 30.6            | 13( 41)         | 19( 59)         | 42/11                    |                     |               |
|                                   | 31.7            | 41( 75)         | 14( 25)         | 67/11                    |                     |               |
|                                   | 32.8            | 111( 99)        | 1( 1)           | 135/11                   |                     |               |
|                                   | 30              | 0( 0)           | ?(100)          | ?/?                      |                     | 48            |
| <i>Caiman crocodilus</i>          |                 |                 |                 |                          |                     |               |
|                                   | 28.5            | 0( 0)           | ?(100)          | ?/?                      |                     | 49            |
|                                   | ~28.9           | 0( 0)           | ?(100)          | ?/?                      |                     |               |
|                                   | ~30.1           | 0( 0)           | ?(100)          | ?/?                      |                     |               |
|                                   | ~30.9           | 0( 0)           | ?(100)          | ?/?                      |                     |               |
|                                   | ~31.4           | ?(~60)          | ?(~40)          | ?/?                      |                     |               |
|                                   | ~31.9           | ?(100)          | 0( 0)           | ?/?                      |                     |               |
|                                   | ~32.3           | ?(100)          | 0( 0)           | ?/?                      |                     |               |
|                                   | 33.0            | ?(100)          | 0( 0)           | ?/?                      |                     |               |
|                                   | 33.5            | ?(100)          | 0( 0)           | ?/?                      |                     |               |
| <i>Paleosuchus trigonatus</i>     |                 |                 |                 |                          |                     |               |
|                                   | ≤31             | 0( 0)           | ?(100)          | ?/?                      |                     | 101           |
|                                   | 32              | ?(100)          | 0( 0)           | ?/?                      |                     |               |
| <b>CROCODYLIDAE</b>               |                 |                 |                 |                          |                     |               |
| <i>Crocodylus johnsoni</i>        |                 |                 |                 |                          |                     |               |
|                                   | 28.0            | 0( 0)           | 4(100)          | ?/?                      | Incubation method A | 96            |
|                                   | 29.0            | 0( 0)           | 31(100)         | ?/?                      | A & B               |               |
|                                   | 30.0            | 0( 0)           | 48(100)         | ?/?                      | A & B               |               |
|                                   | 31.0            | 0( 0)           | 9(100)          | ?/?                      | A                   |               |
|                                   | 31.0            | 2( 13)          | 14( 87)         | ?/?                      | B                   |               |
|                                   | 31.5            | 7( 23)          | 24( 77)         | ?/?                      | B                   |               |
|                                   | 31.7            | 5( 25)          | 15( 75)         | ?/?                      | A                   |               |
|                                   | 32.0            | 4( 31)          | 9( 69)          | ?/?                      | A                   |               |
|                                   | 32.0            | 0( 0)           | 14(100)         | ?/?                      | B                   |               |
|                                   | 32.5            | 6( 23)          | 20( 77)         | ?/?                      | A                   |               |
|                                   | 32.5            | 0( 0)           | 6(100)          | ?/?                      | B                   |               |
|                                   | 33.0            | 0( 0)           | 27(100)         | ?/?                      | A & B               |               |

| <u>Taxa</u>                 | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>  | <u>Source</u> |
|-----------------------------|-----------------|-----------------|-----------------|--------------------------|--|---------------|
|                             | 34.0            | 0( 0)           | 9(100)          | ??                       | A  |               |
|                             | 26              | 0( 0)           | 12(100)         | 122/?                    |  | 97            |
|                             | 27.9            | 0( 0)           | 15(100)         | 44/?                     |  |               |
|                             | 29.9            | 0( 0)           | 48(100)         | 70/?                     |  |               |
|                             | 30              | 5( 19)          | 21( 81)         | 41/?                     | 1 clutch gave 12 females, 2 dead                         |               |
|                             | 30              | 1( 1)           | 123( 99)        | 176/?                    | The male was deformed                                    |               |
|                             | 31.1            | 3( 13)          | 20( 87)         | 28/?                     |  |               |
|                             | 31.7            | 0( 0)           | 5(100)          | 6/?                      |  |               |
|                             | 32.0            | 0( 0)           | 13(100)         | 26/?                     |  |               |
|                             | 32.0            | 6( 30)          | 14( 70)         | 33/?                     |  |               |
|                             | 34              | 0( 0)           | 41(100)         | 131/?                    | Most dead  |               |
|                             | 32              | 5( 29)          | 12( 71)         | ??                       |  | 99            |
|                             | 33              | 5( 20)          | 20( 80)         | ??                       |  |               |
| <i>Crocodylus niloticus</i> |                 |                 |                 |                          |  |               |
|                             | 27.83           | 0( 0)           | 82(100)         | 98/9                     |  | 41            |
|                             | 30.96           | 0( 0)           | 94(100)         | 118/10                   |  |               |
|                             | 32.5            | 10( 91)         | 1( 9)           | 18/1                     |  |               |
|                             | 33.83           | 53(100)         | 0( 0)           | 60/6                     |  |               |
|                             | 33.83           | 9( 82)          | 2( 18)          | 18/1                     |  |               |
|                             | 33.83           | 6( 75)          | 2( 25)          | 8/1                      |  |               |
|                             | 33.83           | 11( 85)         | 2( 15)          | 13/1                     |  |               |
|                             | 33.83           | 3( 33)          | 6( 67)          | 13/1                     |  |               |
| <i>Crocodylus palustris</i> |                 |                 |                 |                          |  |               |
|                             | 28              | 0( 0)           | 27(100)         | ?/6                      | >90% hatching in all treatments except for 33.5° and 34° | 49            |
|                             | 28.5            | 0( 0)           | 35(100)         | ?/4                      |  |               |
|                             | 29              | 0( 0)           | 32(100)         | ?/6                      |  |               |
|                             | 29.5            | 0( 0)           | 22(100)         | ?/4                      |  |               |
|                             | 30              | 0( 0)           | 46(100)         | ?/8                      |  |               |
|                             | 30.5            | 0( 0)           | 17(100)         | ?/3                      |  |               |
|                             | 31              | 0( 0)           | 51(100)         | ?/6                      |  |               |
|                             | 31.5            | 2( 22)          | 7( 78)          | ?/1                      |  |               |
|                             | 32              | 7( 70)          | 3( 30)          | ?/1                      |  |               |
|                             | 32              | 2( 25)          | 6( 75)          | ?/1                      |  |               |
|                             | 32              | 9( 69)          | 4( 31)          | ?/1                      |  |               |
|                             | 32              | 1(100)          | 0( 0)           | ?/1                      |  |               |
|                             | 32.5            | 8(100)          | 0( 0)           | ?/1                      |  |               |
|                             | 33              | 3( 21)          | 11( 79)         | ?/1                      |  |               |
|                             | 33              | 3( 75)          | 1( 25)          | ?/1                      |  |               |
|                             | 33              | 3( 27)          | 8( 73)          | ?/1                      |  |               |
|                             | 33.5            | 0( 0)           | 0( 0)           | 19/?                     |  |               |
|                             | 34              | 0( 0)           | 0( 0)           | 4/?                      |  |               |
| <i>Crocodylus porosus</i>   |                 |                 |                 |                          |  |               |
|                             | 30              | 0( 0)           | ?(100)          | ?/?                      |  | 46            |
|                             | 32              | ?(100)          | 0( 0)           | ?/?                      |  |               |
|                             | 28.0            | 0( 0)           | 4(100)          | ?/?                      | Incubation method A                                      | 96            |

| <u>Taxa</u>                   | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u> | <u>Source</u> |
|-------------------------------|-----------------|-----------------|-----------------|--------------------------|-----------------|---------------|
|                               | 29.0            | 0( 0)           | 26(100)         | ??                       | B               |               |
|                               | 30.0            | 0( 0)           | 70(100)         | ??                       | A & B           |               |
|                               | 31.0            | 1( 50)          | 1( 50)          | ??                       | A               |               |
|                               | 31.0            | 2( 12)          | 15( 88)         | ??                       | B               |               |
|                               | 32.0            | 10( 91)         | 1( 9)           | ??                       | A               |               |
|                               | 32.0            | 52( 85)         | 9( 15)          | ??                       | B               |               |
|                               | 33.0            | 4(100)          | 0( 0)           | ??                       | A               |               |
|                               | 33.0            | 1( 4)           | 25( 96)         | ??                       | B               |               |
| <i>Crocodylus siamensis</i>   |                 |                 |                 |                          |                 |               |
|                               | 28              | 0( 0)           | ?(100)          | ??                       |                 | 47            |
|                               | 32.5            | ?(100)          | 0( 0)           | ??                       |                 |               |
|                               | 27.75           | 0( 0)           | 11(100)         | 11/1                     |                 | 48            |
|                               | 33.0            | 14(100)         | 0( 0)           | 14/1                     |                 |               |
| <b>SQUAMATA</b>               |                 |                 |                 |                          |                 |               |
| <b>LACERTILIA</b>             |                 |                 |                 |                          |                 |               |
| <b>AGAMIDAE</b>               |                 |                 |                 |                          |                 |               |
| <i>Agama agama</i>            |                 |                 |                 |                          |                 |               |
|                               | 26.5            | 1( 2)           | 45( 98)         | ??                       |                 | 19            |
|                               | 29              | 30(100)         | 0( 0)           | ??                       |                 |               |
| <i>Agama caucasia</i>         |                 |                 |                 |                          |                 |               |
|                               | 27              | 21( 72)         | 8( 28)          | 44/5                     |                 | 50            |
|                               | 28              | 19( 95)         | 1( 5)           | 20/2                     |                 |               |
| <b>ANGUIDAE</b>               |                 |                 |                 |                          |                 |               |
| <i>Elgaria multicolor</i>     |                 |                 |                 |                          |                 |               |
|                               | 27.5            | ?(>50)          | ?(<50)          | ??                       |                 | 51            |
| <b>GEKKONIDAE</b>             |                 |                 |                 |                          |                 |               |
| <i>Eublepharis macularius</i> |                 |                 |                 |                          |                 |               |
|                               | 26              | 0( 0)           | 20(100)         | 20/?                     |                 | 7,8           |
|                               | 32.5            | 24( 80)         | 6( 20)          | 30/?                     |                 |               |
|                               | 29.5            | ?( 50)          | ?( 50)          | ??                       |                 | 8             |
|                               | 31.5            | 16( 84)         | 3( 16)          | 19/?                     |                 | 9             |
|                               | 31.5            | 13( 93)         | 1( 7)           | 24/?                     |                 |               |
|                               | 26.7            | ?( 0)           | ?(100)          | ??                       |                 | 58            |
|                               | 32.2            | ?(100)          | ?( 0)           | ??                       |                 |               |
|                               | 27              | ?(<<50)         | ?(>>50)         | ??                       |                 | 87            |

| <u>Taxa</u>                      | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u> | <u>Source</u> |
|----------------------------------|-----------------|-----------------|-----------------|--------------------------|-----------------|---------------|
|                                  | 24              | 0( 0)           | 7(100)          | 10/?                     |                 | 95            |
|                                  | 27.85           | 1( 2)           | 44( 98)         | 59/?                     |                 |               |
|                                  | 32.7            | 14( 88)         | 2( 12)          | 18/?                     |                 |               |
| <i>Gekko japonicus</i>           | 20              | 0( 0)           | 0( 0)           | 20/?                     |                 | 88,89,90      |
|                                  | 24              | 1( 7)           | 13( 93)         | 26/?                     |                 |               |
|                                  | 26              | 4( 22)          | 19( 78)         | ?/?                      |                 |               |
|                                  | 28              | 15( 75)         | 5( 25)          | 30/?                     |                 |               |
|                                  | 30              | 4( 22)          | 19( 78)         | ?/?                      |                 |               |
|                                  | 32              | 5( 24)          | 16( 76)         | 35/?                     |                 |               |
| <i>Hemitheconyx caudicinctus</i> | 28.6            | ?( 0)           | ?(100)          | ?/?                      |                 | 1             |
|                                  | 31.7            | ?(100)          | ?( 0)           | ?/?                      |                 |               |
|                                  | 26.7            | ?( 0)           | ?(100)          | ?/?                      |                 | 58            |
|                                  | 32.2            | ?(100)          | ?( 0)           | ?/?                      |                 |               |
|                                  | ??              | 0( 0)           | ?(100)          | ?/?                      |                 | 95            |
| <i>Tarentola boettgeri</i>       | 28.5            | 0( 0)           | 23(100)         | ?/?                      |                 | 65            |
| <i>Tarentola mauritanica</i>     | 28.5            | 0( 0)           | 33(100)         | ?/?                      |                 | 65            |
| IGUANIDAE                        |                 |                 |                 |                          |                 |               |
| <i>Anolis carolinensis</i>       | 24              | ?(~50)          | ?(~50)          | ?/?                      |                 | 91            |
|                                  | 25              | ?(~50)          | ?(~50)          | ?/?                      |                 |               |
|                                  | 27              | ?(~50)          | ?(~50)          | ?/?                      |                 |               |
|                                  | 28              | ?(~50)          | ?(~50)          | ?/?                      |                 |               |
|                                  | 30              | ?(~50)          | ?(~50)          | ?/?                      |                 |               |
|                                  | 32              | ?(~50)          | ?(~50)          | ?/?                      |                 |               |
|                                  | 34              | ?(~50)          | ?(~50)          | ?/?                      |                 |               |
| <i>Dipsosaurus dorsalis</i>      | 28              | 1( 50)          | 1( 50)          | 11/?                     |                 | 64            |
|                                  | 30              | 6( 35)          | 11( 65)         | 22/?                     |                 |               |
|                                  | 32              | 6( 46)          | 7( 54)          | 16/?                     |                 |               |
|                                  | 34              | 12( 75)         | 4( 25)          | 21/?                     |                 |               |
|                                  | 35              | 0( 0)           | 4(100)          | 4/?                      |                 |               |
|                                  | 36              | 38( 49)         | 41( 51)         | 94/?                     |                 |               |
|                                  | 38              | 7( 32)          | 15( 68)         | 22/?                     |                 |               |
|                                  | 40              | 0( 0)           | 3(100)          | 4/?                      |                 |               |
| <i>Sceloporus jarrovi</i>        | 26              | 14( 37)         | 24( 63)         | ?/?                      |                 | 3             |
|                                  | 28              | 28( 70)         | 12( 30)         | ?/?                      |                 |               |

| <u>Taxa</u>                    | <u>Temp (C)</u> | <u>#♂♂ (%)</u> | <u>#♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>          | <u>Source</u> |
|--------------------------------|-----------------|----------------|----------------|--------------------------|--------------------------|---------------|
|                                | 30              | 40( 50)        | 40( 50)        | ?/?                      |                          |               |
|                                | 32              | 51( 49)        | 53( 51)        | ?/?                      |                          |               |
|                                | 34              | 46( 46)        | 54( 54)        | ?/?                      |                          |               |
|                                | 36              | 34( 65)        | 18( 35)        | ?/?                      |                          |               |
| <i>Sceloporus undulatus</i>    |                 |                |                |                          |                          |               |
|                                | 30              | 19( 53)        | 17( 47)        | 37/8                     |                          | 81            |
|                                | 24              | ?(~50)         | ?(~50)         | ?/?                      |                          | 91            |
|                                | 25              | ?(~50)         | ?(~50)         | ?/?                      |                          |               |
|                                | 27              | ?(~50)         | ?(~50)         | ?/?                      |                          |               |
|                                | 28              | ?(~50)         | ?(~50)         | ?/?                      |                          |               |
|                                | 30              | ?(~50)         | ?(~50)         | ?/?                      |                          |               |
|                                | 32              | ?(~50)         | ?(~50)         | ?/?                      |                          |               |
|                                | 34              | ?(~50)         | ?(~50)         | ?/?                      |                          |               |
| LACERTIDAE                     |                 |                |                |                          |                          |               |
| <i>Lacerta viridis</i>         |                 |                |                |                          |                          |               |
|                                | 29              | ?(~50)         | ?(~50)         | ?/?                      |                          | 27            |
|                                | 17.5            | 6( 67)         | 3( 33)         | 23/~3                    | First 5-7 days at 25 C   | 78            |
|                                | 19.5            | 4( 57)         | 3( 43)         | 24/~3                    | First 13 days at 25 C    |               |
|                                | 35.5            | 11( 55)        | 9( 45)         | ?/3                      | First 5-6 days at 25 C   |               |
|                                | 35.5            | 4( 33)         | 8( 67)         | ?/2                      | First 13-14 days at 25 C |               |
| <i>Podarcis pityusensis</i>    |                 |                |                |                          |                          |               |
|                                | 29              | ?( ~8)         | ?(~92)         | ?/?                      | 97% hatch success        | 27            |
| TEIIDAE                        |                 |                |                |                          |                          |               |
| <i>Cnemidophorus inornatus</i> |                 |                |                |                          |                          |               |
|                                | 25              | 12( 55)        | 10( 45)        | 22/?                     |                          | 20            |
|                                | 30              | 10( 50)        | 10( 50)        | 20/?                     |                          |               |
| <i>Cnemidophorus uniparens</i> |                 |                |                |                          |                          |               |
|                                | 25              | 0( 0)          | 78(100)        | 78/?                     | parthenogenetic          | 20            |
|                                | 26              | 0( 0)          | 32(100)        | 32/?                     |                          |               |
|                                | 29              | 0( 0)          | 52(100)        | 52/?                     |                          |               |
|                                | 30              | 0( 0)          | 38(100)        | 38/?                     |                          |               |
|                                | 31              | 0( 0)          | 44(100)        | 44/?                     |                          |               |
|                                | 33              | 0( 0)          | 0( 0)          | 5/?                      |                          |               |
| SERPENTES                      |                 |                |                |                          |                          |               |
| COLUBRIDAE                     |                 |                |                |                          |                          |               |
| <i>Boiga dendrophila</i>       |                 |                |                |                          |                          |               |
|                                | 29.25           | 4( 50)         | 4( 50)         | 9/1                      |                          | 2             |
|                                | 30              | 3( 60)         | 2( 40)         | 7/1                      |                          |               |

| <u>Taxa</u>                    | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u>                                      | <u>Source</u> |
|--------------------------------|-----------------|-----------------|-----------------|--------------------------|--|---------------|
| <i>Cemophora coccinea</i>      | 25              | 5( 71)          | 2( 29)          | 7/1                      |  | 5             |
| <i>Clelia clelia</i>           | 27              | 6( 67)          | 3( 33)          | 10/1                     |  | 55            |
| <i>Coluber constrictor</i>     | 26.5            | 7( 78)          | 2( 22)          | 14/1                     |  | 32            |
| <i>Nerodia fasciata</i>        | 21.65           | 99( 55)         | 81( 45)         | ?/10                     |  | 67            |
|                                | 26.4            | 135( 48)        | 144( 52)        | ?/12                     |  |               |
|                                | 30.0            | 101( 46)        | 119( 54)        | ?/11                     |  |               |
| <i>Pituophis melanoleucus</i>  | 21              | 1( 11)          | 8( 89)          | 19/?                     | Moved to 23° after 70 days<br>Sexed embryos excluded |               |
|                                | 23              | 20( 35)         | 37( 65)         | 73/?                     |  |               |
|                                | 23              | 24( 41)         | 35( 59)         | 66/?                     |  |               |
|                                | 26              | 17( 46)         | 20( 54)         | 43/?                     |  |               |
|                                | 28              | 34( 50)         | 34( 50)         | 73/?                     |  |               |
|                                | 28              | 47( 49)         | 49( 51)         | 97/?                     |  |               |
|                                | 30              | 18( 47)         | 20( 53)         | 39/?                     |  |               |
|                                | 32              | 28( 58)         | 20( 42)         | 62/?                     |  |               |
|                                | 33              | 31( 55)         | 25( 45)         | 64/?                     |  |               |
| <i>Xenocalamus bicolor</i>     | 31              | 2( 50)          | 2( 50)          | 4/1                      |  |               |
| ELAPIDAE                       |                 |                 |                 |                          |  |               |
| <i>Acanthophis antarcticus</i> | 29              | 9( 45)          | 11( 55)         | 20/1                     |  | 39            |
|                                | 29              | 8( 42)          | 11( 58)         | 19/1                     |  |               |
|                                | 29              | 9( 47)          | 10( 53)         | 19/1                     |  |               |
|                                | 29              | 12( 50)         | 12( 50)         | 25/1                     |  |               |
|                                | 29              | 8( 40)          | 12( 60)         | 20/1                     |  |               |
|                                | 29              | 10( 48)         | 11( 52)         | 21/1                     |  |               |
|                                | 29              | 10( 43)         | 13( 57)         | 23/1                     |  |               |
|                                | 29              | 8( 47)          | 9( 53)          | 17/1                     |  |               |
| <i>Pseudechis australis</i>    | 28              | 11( 79)         | 3( 21)          | 15/1                     |  |               |
|                                | 27              | 6( 55)          | 5( 45)          | 15/1                     | Female #1  | 59            |
|                                | 27              | 8( 67)          | 4( 33)          | 14/1                     | "  |               |
|                                | 27              | 9( 90)          | 1( 10)          | 12/1                     | Female #2  |               |
| <i>Pseudechis colletti</i>     | 28              | 13( 81)         | 3( 19)          | 18/1                     |  | 16            |
|                                | 28              | 4( 36)          | 7( 64)          | 12/1                     |  |               |

| <u>Taxa</u>                         | <u>Temp (C)</u> | <u># ♂♂ (%)</u> | <u># ♀♀ (%)</u> | <u># eggs/# clutches</u> | <u>Comments</u> | <u>Source</u> |
|-------------------------------------|-----------------|-----------------|-----------------|--------------------------|-----------------|---------------|
|                                     | 28.5            | 2( 40)          | 3( 60)          | 7/1                      |                 | 18            |
| <i>Pseudechis guttatus</i>          | 28              | 5( 50)          | 5( 50)          | 10/1                     |                 | 16            |
|                                     | 28              | 5( 63)          | 3( 37)          | 8/1                      |                 |               |
| <i>Pseudolaticauda semifasciata</i> | 28              | 51( 61)         | 32( 39)         | 114/29                   |                 | 66            |
| PYTHONIDAE                          |                 |                 |                 |                          |                 |               |
| <i>Aspidites melanocephalus</i>     | 30              | 2( 25)          | 6( 75)          | 8/1                      |                 | 17            |
| <i>Morelia amethystina</i>          | 30              | 3( 43)          | 4( 57)          | 7/1                      |                 | 17            |
| <i>Morelia spilota</i>              | 30              | 11( 52)         | 10( 48)         | 23/1                     |                 | 17            |
|                                     | 30              | 5(100)          | 0( 0)           | 7/1                      |                 |               |
| VIPERIDAE                           |                 |                 |                 |                          |                 |               |
| <i>Crotalus vegrandis</i>           | 28              | 5( 63)          | 3( 37)          | 8/1                      |                 | 15            |

SOURCES--1: Anderson and Oldham (1988), 2: Bakken and Bakken (1988), 3: Beuchat (1983), 4: Branch and Patterson (1976), 5: Braswell and Palmer (1984), 6: Bull (1980), 7: Bull (1987a), 8: Bull (1987b), 9: Bull et al. (1988), 10: Bull et al. (1985), 11: Bull and Vogt (1979), 12: Bull et al. (1982a), 13: Bull et al. (1982b), 14: Burger and Zappalorti (1988), 15: Carl et al. (1982), 16: Charles (1988), 17: Charles et al. (1985), 18: Charles et al. (1983), 19: Charnier (1966), 20: Crews (1989), 21: Crews et al. (1989), 22: Dimond (1979), 23: Dimond (1983), 24: Dimond and Mohanty-Hejmadi (1983), 25: Dutton et al. (1985), 26: Ehrengart (1971), 27: Eichenberger (1981), 28: Ewert and Nelson (in press), 29: Ferguson and Joanen (1982), 30: Ferguson and Joanen (1983), 31: Georges (1988), 32: Gillingham (1976), 33: Gutzke and Bull (1986), 34: Gutzke and Chymiy (1988), 35: Gutzke and Packard (1987), 36: Gutzke and Paukstis (1983), 37: Gutzke and Paukstis (1984), 38: Harry and Limpus (1989), 39: Hay and Magnusson (1986), 40: Hou (1985), 41: Hutton (1987), 42: Janzen (1987), 43: Janzen et al. (in press), 44: Joanen and McNease (1989), 45: Joanen et al. (1987), 46: Joss and Cuff (1987), 47: Lang (1985), 48: Lang (1987), 49: Lang et al. (1989), 50: Langerwerf (1983), 51: Langerwerf (1984), 52: Lescure et al. (1985), 53: Limpus et al. (1983), 54: Limpus et al. (1985), 55: Martinez and Cerdas (1986), 56: McCoy et al. (1983), 57: Miller and Limpus (1981), 58: Miller (1979), 59: Mirtschin (1988), 60: Mohanty-Hejmadi et al. (1985), 61: Mohanty-Hejmadi and Dimond (1986), 62: Mrosovsky (1988), 63: Mrosovsky et al. (1984), 64: Muth and Bull (1981), 65: Nakamoto and Toriba (1986), 66: Nettmann and Rykena (1985), 67: Osgood (1978), 68: Packard et al. (1989), 69: Packard et al. (1984), 70: Packard et al. (1985), 71: Pieau (1971), 72: Pieau (1972), 73: Pieau (1973), 74: Pieau (1975a), 75: Pieau (1975b), 76: Pieau (1976), 77: Pieau (1978), 78: Raynaud and Pieau (1972), 79: Rimblot et al. (1985), 80: Rimblot-Baly et al. (1987), 81: Roggenbuck and Jenssen (1986), 82: Sachsse (1984), 83: Schwarzkopf and Brooks (1985), 84: Standora and Spotila (1985), 85: Thompson (1983), 86: Thompson (1988), 87: Thorogood and Whimster (1979), 88: Tokunaga (1985), 89: Tokunaga (1986), 90: Tokunaga (1989), 91: Viets



(1989), 92: Vogt (1980), 93: Vogt and Bull (1982), 94: Vogt et al. (1982), 95: Wagner (1980), 96: Webb et al. (1987), 97: Webb et al. (1983), 98: Webb et al. (1986), 99: Webb and Smith (1984), 100: Wood and Wood (1982), 101: Yamakoshi et al. (1987), 102: Yntema (1976), 103: Yntema (1981), 104: Yntema and Mrosovsky (1979), 105: Yntema and Mrosovsky (1980), 106: Yntema and Mrosovsky (1982), 107: Zaborski et al. (1982), 108: Zaborski et al. (1988).

## REFERENCES

- Anderson, A. and C. Oldham. 1988. Captive husbandry and propagation of the African fat-tail gecko *Hemitheconyx caudicinctus*. Pp. 75-85. In: K. H. Peterson (ed.), 10th Int. Herpetol. Symp. on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Bakken, D. J. and L. Bakken. 1988. Husbandry and reproduction in *Boiga dendrophila* including a case of delayed fertilization. Pp. 135-140. In: K. H. Peterson (ed.), 10th Int. Herpetol. Symp. on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- Beuchat, C. A. 1988. Temperature effects during gestation in a viviparous lizard. *J. Therm. Biol.* 13: 135-142.
- Branch, W. R. and R. W. Patterson. 1976. Notes on eggs and hatchlings of *Xenocalamus bicolor lineatus* Roux from South Africa. *Herpetol. Rev.* 7: 116-117.
- Braswell, A. L. and W. M. Palmer. 1984. *Cemophora coccinea copei* (northern scarlet snake). Reproduction. *Herpetol. Rev.* 15: 49.
- Bull, J. J. 1980. Sex determination in reptiles. *Quart. Rev. Biol.* 55: 3-21.
- , 1987a. Temperature-sensitive periods of sex determination in a lizard: similarities with turtles and crocodylians. *J. Exp. Zool.* 241: 143-148.
- , 1987b. Temperature-dependent sex determination in reptiles: validity of sex diagnosis in hatchling lizards. *Canadian J. Zool.* 65: 1421-1424.
- , W. H. N. Gutzke and D. Crews. 1988. Sex reversal by estradiol in three reptilian orders. *Gen. Comp. Endocrinol.* 70: 425-428.
- , J. M. Legler and R. C. Vogt. 1985. Non-temperature dependent sex determination in two suborders of turtles. *Copeia* 1985: 784-786.
- and R. C. Vogt. 1979. Temperature-dependent sex determination in turtles. *Science* 206: 1186-1188.
- , ----- and M. G. Bulmer. 1982a. Heritability of sex ratio in turtles with environmental sex determination. *Evolution* 36: 333-341.
- , ----- and C. J. McCoy. 1982b. Sex determining temperatures in turtles: a geographic comparison. *Evolution* 36: 326-332.
- Burger, J. and R. T. Zappalorti. 1988. Effects of incubation temperature on sex ratios in pine snakes: differential vulnerability of males and females. *Amer. Natur.* 132: 492-505.
- Carl, G., K. H. Peterson and R. M. Hubbard. 1982. Reproduction in captive Uracoan rattlesnakes, *Crotalis vegrandis*. *Herpetol. Rev.* 13: 42-43.
- Charles, N. 1988. Captive reproduction of seven species of Australian snakes of the families Elapidae and Boidae. Pp. 158-168. In: K. H. Peterson (ed.), 10th Int. Herpetol. Symp. on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- , R. Field and R. Shine. 1985. Notes on the reproductive biology of Australian pythons, genera *Aspidites*, *Liasis* and *Morelia*. *Herpetol. Rev.* 16: 45-48.
- , A. Watts and R. Shine. 1983. Captive reproduction in an Australian elapid snake *Pseudechis colletti*. *Herpetol. Rev.* 14: 16-18.
- Charnier, M. 1966. Action de la température sur la sex-ratio chez l'embryon d'*Agama agama* (Agamidae, Lacertilien). *Soc. Biol. Ovest Africa* 160: 620-622.
- Crews, D. 1989. Absence of temperature-dependent sex determination in congeneric sexual and parthenogenetic *Cnemidophorus* lizards. *J. Exp. Zool.* 252: 318-320.
- , T. Wibbels and W. H. N. Gutzke. 1989. Action of sex steroid hormones on temperature-induced sex determination in the snapping turtle (*Chelydra serpentina*). *Gen. Comp. Endocrinol.* 76: 159-166.
- Dimond, M. T. 1979. Sex differentiation and incubation temperature in turtles. *Amer. Zool.* 19: 981 (Abstr.).

- , 1983. Sex of turtle hatchlings as related to incubation temperature. Pp. 88-101 *In*: Proc. 6th Reptile Symp. on Captive Propagation and Husbandry. Zoological Consortium; Thurmont, Maryland.
- and P. Mohanty-Hejmadi. 1983. Incubation temperature and sex differentiation in a sea turtle. *Amer. Zool.* 23: 1017 (Abstr.).
- Dutton, P. H., C. P. Whitmore and N. Mrosovsky. 1985. Masculinisation of leatherback turtle *Dermochelys coriacea* hatchlings from eggs incubated in styrofoam boxes. *Biol. Conserv.* 31: 249-264.
- Ehregart, W. 1971. On the breeding and rearing of Greek tortoises. *Salamandra* 7: 71-80.
- Eichenberger, P. 1981. Smaragdhagedissen kweken op bestelling. *Lacerta* 39: 72-77.
- Ewert, M. A. and C. E. Nelson. In press. Sex determination in turtles: diverse patterns and some possible adaptive values. *Copeia*.
- Ferguson, M. W. J. and T. Joanen. 1982. Temperature of egg incubation determines sex in *Alligator mississippiensis*. *Nature* 296: 850-853.
- and -----, 1983. Temperature-dependent sex determination in *Alligator mississippiensis*. *J. Zool.* 200: 143-177.
- Georges, A. 1988. Sex determination is independent of incubation temperature in another chelid turtle, *Chelodina longicollis*. *Copeia* 1988: 248-254.
- Gillingham, J. C. 1976. Early egg deposition by the southern black racer, *Coluber constrictor priapus*. *Herpetol. Rev.* 7: 115.
- Gutzke, W. H. N. and J. J. Bull. 1986. Steroid hormones reverse sex in turtles. *Gen. Comp. Endocrinol.* 64: 368-372.
- and D. B. Chymiy. 1988. Sensitive periods during embryogeny for hormonally induced sex determination in turtles. *Gen. Comp. Endocrinol.* 71: 265-267.
- and G. C. Packard. 1987. The influence of temperature on eggs and hatchlings of Blanding's turtles, *Emydoidea blandingii*. *J. Herpetol.* 21: 161-163.
- and G. L. Paukstis. 1983. Influence of the hydric environment on sexual differentiation of turtles. *J. Exp. Zool.* 226: 467-469.
- and -----, 1984. A low threshold temperature for sexual differentiation in the painted turtle, *Chrysemys picta*. *Copeia* 1984: 546-547.
- Harry, J. L. and C. L. Limpus. 1989. Low-temperature protection of marine turtle eggs during long-distance relocation. *Australian Wildl. Res.* 16: 317-320.
- Hay, M. and W. E. Magnusson. 1986. A captive breeding of the Australian death adder, *Acanthophis antarcticus*. *Herpetol. Rev.* 17: 13-14.
- Hou, L. 1985. Sex determination by temperature for incubation in *Chinemys reevesii*. *Acta Herpetol. Sinica* 4: 130.
- Hutton, J. M. 1987. Incubation temperatures, sex ratios and sex determination in a population of Nile crocodiles (*Crocodylus niloticus*). *J. Zool.* 211: 143-155.
- Janzen, F. J. 1987. Influence of environmental conditions on use of metabolic substrates and growth by embryonic snapping turtles. *Amer. Zool.* 27: 54A (Abstr.).
- , G. C. Packard, M. J. Packard, T. J. Boardman and J. R. zumBrunnen. In press. Mobilization of lipid and protein by embryonic snapping turtles in wet and dry environments. *J. Exp. Zool.*
- and G. L. Paukstis. In press. Environmental sex determination in reptiles: ecology, evolution, and experimental design. *Quart. Rev. Biol.*
- Joanen, T. and L. L. McNease. 1989. Ecology and physiology of nesting and early development of the American alligator. *Amer. Zool.* 29: 987-998.

- , L. McNease and M. W. J. Ferguson. 1987. The effects of egg incubation temperature on post-hatching growth of American alligators. Pp. 533-537 *In*: G. J. W. Webb, S. C. Manolis and P. J. Whitehead (eds.), *Wildlife Management: Crocodiles and Alligators*. Surrey Beatty and Sons; New South Wales, Australia.
- Joss, J. M. P. and A. Cuff. 1987. Steroidogenesis during gonadal differentiation in the salt water crocodile, *Crocodylus porosus*. *Amer. Zool.* 27: 66A (Abstr.).
- Lang, J. W. 1985. Incubation temperature affects thermal selection of hatchling crocodiles. *Amer. Zool.* 25: 18A (Abstr.).
- , 1987. Crocodylian thermal selection. Pp. 301-317 *In*: G. J. W. Webb, S. C. Manolis and P. J. Whitehead (eds.), *Wildlife Management: Crocodiles and Alligators*. Surrey Beatty and Sons; New South Wales, Australia.
- , H. Andrews and R. Whitaker. 1989. Sex determination and sex ratios in *Crocodylus palustris*. *Amer. Zool.* 29: 935-952.
- Langerwerf, B. 1983. Über die Haltung und Zucht von *Agama caucasia* (Eichwald, 1831) (Sauria: Agamidae), nebst Bemerkungen zur erfolgreichen Zucht weiterer palaearktischer Echsen. *Salamandra* 19: 11-20.
- , 1984. Techniques for large-scale breeding of lizards from temperate climates in greenhouse enclosures (breeding many species of lizards in captivity, aiming the maintenance of populations of each species outside their natural habitat). *Acta Zool. Pathol. Antverpiensa* 78: 163-176.
- Lescure J., F. Rimblot, J. Fretey, S. Renous and C. Pieau. 1985. Influence de la température d'incubation des oeufs sur la sex-ratio des nouveaux-nés de la tortue luth, *Dermochelys coriacea*. *Bull. Soc. Zool. France* 110: 355-359.
- Limpus, C. J., P. Reed, and J. D. Miller. 1983. Islands and turtles. The influence of choice of nesting beach on sex ratio. Pp. 397-402 *In*: J. T. Baker, R. M. Carter, P. W. Sammarco and K. P. Stark (eds.), *Proceedings: Inaugural Great Barrier Reef Conference*. James Cook University Press; Townsville.
- , ----- and -----, 1985. Temperature dependent sex determination in Queensland sea turtles: intraspecific variation in *Caretta caretta*. Pp. 343-351 *In*: G. Grigg, R. Shine and H. Ehmann (eds.), *Biology of Australasian Frogs and Reptiles*. Royal Zoological Society; New South Wales.
- Martinez, S. and L. Cerdas. 1986. Captive reproduction of the mussurana, *Clelia clelia* (Daudin) from Costa Rica. *Herpetol. Rev.* 17: 12.
- McCoy, C. J., R. C. Vogt and E. J. Censky. 1983. Temperature-controlled sex determination in the sea turtle *Lepidochelys olivacea*. *J. Herpetol.* 17: 404-406.
- Miller, J. D. and C. J. Limpus. 1981. Incubation period and sexual differentiation in the green turtle *Chelonia mydas* L. Pp. 66-73 *In*: C. B. Banks and A. A. Martin (eds.), *Proc. Melbourne Herpetol. Symp.* Zoological Board of Victoria; Parkville.
- Miller, M. 1979. Preliminary notes on the breeding of geckos in captivity. *Bull. Chicago Herpetol. Soc.* 14: 78-91.
- Mirtschin, P. J. 1988. Captive breeding and oviparity in the king brown snake *Pseudechis australis* (Serpentes Elapidae) from Eyre Peninsula South Australia. Pp. 141-148. *In*: K. H. Peterson (ed.), *10th Int. Herpetol. Symp. on Captive Propagation and Husbandry*. Zoological Consortium; Thurmont, Maryland.
- Mohanty-Hejmadi, P, M. Behra and M. T. Dimond. 1985. Temperature dependent sex differentiation in the olive ridley *Lepidochelys olivacea* and its implications for conservation. Pp. 1-5 *In*: *Symposium on Endangered Marine Animals and Marine Parks, 12-18 January, Paper 25*. Marine Biological Association of India; Cochin.
- and M. T. Dimond. 1986. Temperature dependent sex determination in the olive ridley turtle. Pp. 159-162 *In*: *Progress in Developmental Biology, Part A*. Alan R. Liss; New York.
- Mrosovsky, N. 1988. Pivotal temperatures for loggerhead turtles (*Caretta caretta*) from northern and southern nesting beaches. *Canadian J. Zool.* 66: 661-669.

- , P. H. Dutton and C. P. Whitmore. 1984. Sex ratios of two species of sea turtle nesting in Suriname. *Canadian J. Zool.* 62: 2227-2239.
- Muth, A. and J. J. Bull. 1981. Sex determination in desert iguanas: does incubation temperature make a difference? *Copeia* 1981: 869-870.
- Nakamoto, E. and M. Toriba. 1986. Successful artificial incubation of the eggs of Erabu sea snake, *Pseudolaticauda semifasciata* (Reinwardt). *Snake* 18: 55-56.
- Nettmann, H.-K. and S. Rykena. 1985. Verhaltens- und fortpflanzungsbiologische Notizen über kanarische und nordafrikanische *Tarentola*-Arten. *Bonner Zool. Beitr.* 36: 287-305.
- Osgood, D. W. 1978. Effects of temperature on the development of meristic characters in *Natrix fasciata*. *Copeia* 1978: 33-47.
- Packard, G. C., M. J. Packard and G. F. Birchard. 1989. Sexual differentiation and hatching success by painted turtles incubating in different thermal and hydric environments. *Herpetologica* 45: 385-392.
- , ----- and T. J. Boardman. 1984. Effects of the hydric environment on metabolism of embryonic snapping turtles do not result from altered patterns of sexual differentiation. *Copeia* 1984: 547-550.
- , ----- and W. H. N. Gutzke. 1985. Influence of hydration of the environment on eggs and embryos of the terrestrial turtle *Terrapene ornata*. *Physiol. Zool.* 58: 564-575.
- Pieau, C. 1971. Sur la proportion sexuelle chez les embryons de deux Chéloniens (*Testudo graeca* L. et *Emys orbicularis* L.) issus d'oeufs incubés artificiellement. *C. R. Hebd. Séanc. Acad. Sci., Paris (Ser. D)* 274: 3071-3074.
- 1972. Effets de la température sur le développement des glandes génitales chez les embryons de deux Chéloniens, *Emys orbicularis* L. et *Testudo graeca* L. *C. R. Hebd. Séanc. Acad. Sci., Paris (Ser. D)* 274: 719-722.
- 1973. Nouvelles données expérimentales concernant les effets de la température sur la différenciation sexuelle chez les embryons de Chéloniens. *C. R. Hebd. Séanc. Acad. Sci., Paris (Ser. D)* 277: 2789-2792.
- 1975a. Temperature and sex differentiation in embryos of two chelonians, *Emys orbicularis* L. and *Testudo graeca* L. Pp. 332-339 *In*: R. Reinboth (ed.), *Intersexuality in the Animal Kingdom*. Springer-Verlag; New York.
- 1975b. Action de la température sur le phenotype sexuel des embryons d'un chelonien, *Emys orbicularis* L. *Bull. Soc. Zool. France Suppl.* 1: 12-22.
- 1976. Données récentes sur la différenciation sexuelle en fonction de la température chez les embryons d'*Emys orbicularis* L. (Chélonien). *Bull. Soc. Zool. Fr. Suppl.* 4: 46-53.
- 1978. Effets de températures d'incubation basses et élevées, sur la différenciation sexuelle chez des embryons d'*Emys orbicularis* L. (Chélonien). *C. R. Hebd. Séanc. Acad. Sci., Paris (Ser. D)* 286: 121-124.
- Raynaud, A. and C. Pieau. 1972. Effets de diverses températures d'incubation sur le développement somatique et sexuel des embryons de lézard vert (*Lacerta viridis* Laur.). *C. R. Hebd. Séanc. Acad. Sci., Paris (Ser. D)* 275: 2259-2262.
- Rimblot, F., J. Fretey, N. Mrosovsky, J. Lescure and C. Pieau. 1985. Sexual differentiation as a function of the incubation temperature of eggs in the sea-turtle *Dermochelys coriacea* (Vandelli, 1761). *Amphibia-Reptilia* 6: 83-92.
- Rimblot-Baly, F., J. Lescure, J. Fretey and C. Pieau. 1987. Sensibilité à la température de la différenciation sexuelle chez la Tortue Luth, *Dermochelys coriacea* (Vandelli, 1761); application des données de l'incubation artificielle à l'étude de la sex-ratio dans la nature. *Ann. Sci. Natur. Zool. Paris* 8: 277-290.
- Roggenbuck, M. E. and T. A. Jenssen. 1986. The ontogeny of display behaviour in *Sceloporus undulatus* (Sauria: Iguanidae). *Ethology* 71: 153-165.

- Sachsse, W. 1984. Long term studies of the reproduction of *Malaclemys terrapin centrata*. Acta Zool. Pathol. Antverpiensa 78: 297-308.
- Schwarzkopf, L. and R. J. Brooks. 1985. Sex determination in northern painted turtles: effect of incubation at constant and fluctuating temperatures. Canadian J. Zool. 63: 2543-2547.
- Standora, E. A. and J. R. Spotila. 1985. Temperature dependent sex determination in sea turtles. Copeia 1985: 711-722.
- Thompson, M. B. 1983. The Physiology and Ecology of the Eggs of Pleurodiran Tortoise *Emydura macquarii* (Gray), 1831. Ph.D. Thesis, University of Adelaide, South Australia.
- , 1988. Influence of incubation temperature and water potential on sex determination in *Emydura macquarii* (Testudines: Pleurodira). Herpetologica 44: 86-90.
- Thorogood, J. and I. W. Whimster. 1979. The maintenance and breeding of the leopard gecko *Eublepharis macularius*. Intern. Zoo Yearbk. 19: 74-78.
- Tokunaga, S. 1985. Temperature-dependent sex determination in *Gekko japonicus* (Gekkonidae, Reptilia). Devel. Growth Differ. 27: 117-120.
- , 1986. Ecological significance of temperature-dependent sex determination in reptiles. Acta Herpetol. Sinica 5: 59-60.
- , 1989. Temperature-dependent sex determination and the life history of *Gekko japonicus* (Reptilia: Gekkonidae). (Abstr.) In: First World Congress of Herpetology, 11-19 September 1989, Canterbury.
- Viets, B. E. 1989. The effects of temperature on sex and development in the iguanid lizards *Anolis carolinensis* and *Sceloporus undulatus*. Amer. Zool. 29: 28A (Abstr.).
- Vogt, R. C. 1980. Natural history of the map turtles *Graptemys pseudogeographica* and *G. ouachitensis* in Wisconsin. Tulane Stud. Zool. Bot. 22: 17-48.
- and J. J. Bull. 1982. Genetic sex determination in the spiny softshell *Trionyx spiniferus* (Testudines: Trionychidae)(?). Copeia 1982: 699-700.
- , -----, C. J. McCoy and T. W. Houseal. 1982. Incubation temperature influences sex determination in kinosternid turtles. Copeia 1982: 480-482.
- Wagner, E. 1980. Gecko husbandry and reproduction. Pp. 115-117 In: J. B. Murphy and J. T. Collins (eds.), Reproductive Biology and Diseases of Captive Reptiles. SSAR Contributions to Herpetology No. 1.
- Webb, G. J. W., A. M. Beal, S. C. Manolis and K. E. Dempsey. 1987. The effects of incubation temperature on sex determination and embryonic development rate of *Crocodylus johnstoni* and *C. porosus*. Pp. 507-531 In: G. J. W. Webb, S. C. Manolis and P. J. Whitehead (eds.), Wildlife Management: Crocodiles and Alligators. Surrey Beatty and Sons, New South Wales, Australia.
- , R. Buckworth and S. C. Manolis. 1983. *Crocodylus johnstoni* in the McKinlay River, N. T. VI. Nesting biology. Australian Wildl. Res. 10: 607-637.
- , D. Choquenot and P. J. Whitehead. 1986. Nests, eggs, and embryonic development of *Carettochelys insculpta* (Chelonia: Carettochelidae) from Northern Australia. J. Zool., B 1: 521-550.
- and A. M. A. Smith. 1984. Sex ratio and survivorship in the Australian freshwater crocodile *Crocodylus johnstoni*. Symp. Zool. Soc. London 52: 319-355.
- Wood, F. E. and J. R. Wood. 1982. Sex ratios in captive-reared green turtles, *Chelonia mydas*. Copeia 1982: 482-485.
- Yamakoshi, M., W. E. Magnusson and J. M. Hero. 1987. The nesting biology of *Paleosuchus trigonatus*: sources of heat for nests, survivorship and sex ratios. Amer. Zool. 27: 67A (Abstr.).
- Yntema, C. L. 1976. Effects of incubation temperatures on sexual differentiation in the turtle, *Chelydra serpentina*. J. Morphol. 150: 453-462.
- , 1981. Characteristics of gonads and oviducts in hatchlings and young of *Chelydra serpentina* resulting from three incubation temperatures. J. Morphol. 167: 297-304.

- and N. Mrosovsky. 1979. Incubation temperature and sex ratio in hatchling loggerhead turtles: a preliminary report. *Marine Turtle Newsl.* 11: 9-10.
- and -----, 1980. Sexual differentiation in hatchling loggerheads (*Caretta caretta*) incubated at different controlled temperatures. *Herpetologica* 36: 33-36.
- and -----, 1982. Critical periods and pivotal temperatures for sexual differentiation in loggerhead sea turtles. *Canadian J. Zool.* 60: 1012-1016.
- Zaborski, P., M. Dorizzi and C. Pieau. 1982. H-Y antigen expression in temperature sex-reversed turtles (*Emys orbicularis*). *Differentiation* 22: 73-78.
- , ----- and -----, 1988. Temperature-dependent gonadal differentiation in the turtle *Emys orbicularis*: concordance between sexual phenotype and serological H-Y antigen expression at threshold temperature. *Differentiation* 38: 17-20.








ANNOTATED BIBLIOGRAPHY TO THE HERPETOFAUNA  
OF THE  
PINE-OAK WOODLANDS  
OF THE  
SIERRA MADRE OCCIDENTAL, MEXICO



JAMES R. MCCRANIE  
&  
LARRY DAVID WILSON

Department of Biology  
Miami Dade Community College



SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
NO. 84

**SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE**

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with the Division of Amphibians and Reptiles.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions on preparation and submission. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington DC 20560 USA. Please include a self-addressed mailing label with requests.

## INTRODUCTION

This bibliography is a result of a literature search which helped form the foundation for our study entitled "The biogeography of the herpetofauna of the pine-oak woodlands of the Sierra Madre Occidental of Mexico" Milwaukee Pub. Mus. Contrib. Biol. Geol. (72):1-30, 1987.

The bibliography includes all references known to the authors that contain bonafide records of the occurrence of amphibians and reptiles in the pine-oak woodlands of the Sierra Madre Occidental of Mexico. The reader is referred to that study for a definition of the limits of the study area.

The 86 species included in the study and their literature citations are presented below in alphabetical order within their respective orders. We believe this bibliography (188 references) to be relatively complete through the year 1986. In addition, six later references are included in a addendum.

We wish to extend our sincere gratitude to Joy-Ann Perard for her typing of the manuscript. It was a time-consuming task, and we much appreciate her help.

## SPECIES LIST

## Class Amphibia

## Order Caudata

Ambystoma rosaceum 3, 5, 6, 28, 38, 48, 63, 79, 86, 94, 113, 114, 149, 159  
(also as tigrinum), 160 (as tigrinum), 175, 176, 177, 179, 190.

Ambystoma tigrinum 28, 113, 175.

Pseudoeurycea bellii 85, 161.

## Order Salientia

Bufo compactilis 38, 73, 160, 170, 175.

Bufo microscaphus 3, 22, 24 (as compactilis), 56, 106, 133, 170, 175, 176,  
177, 191.

Bufo occidentalis 38, 56, 57, 73, 93, 94, 106, 152, 160, 175, 176.

Bufo punctatus 160.

Bufo woodhousei 160.

Eleutherodactylus occidentalis 176.

Eleutherodactylus tarahumaraensis 142, 147, 175, 186

Hyla arenicolor 40, 42, 56, 57, 73, 79, 94, 152, 160, 175, 177, 186.

Hyla bistincta 40, 175.

Hyla eximia 3, 40, 42, 56, 63, 69, 73, 79, 131, 133, 160, 175, 176, 177, 179,  
186.

Rana chiricahuensis (all as R. pipiens or synonyms except 62, 102, and 175  
unless otherwise noted) 3, 19, 38, 42, 56 (as forreri), 57, 62, 63, 73, 94,  
102, 109, 111, 114, 152, 160, 175, 176, 177.

Rana magnaocularis 46, 62.

Rana pustulosa 57, 61, 175, 183.

Rana tarahumarae 18, 19, 35, 61, 62, 73, 79, 94, 131, 177, 185, 188.

Scaphiopus multiplicatus 38, 160, 175.

Tomodactylus nitidus 57, 175, 176.

Tomodactylus saxatilis 57, 164, 175, 176, 179.

Class Reptilia

Order Testudines

Kinosternon hirtipes (Because of considerable confusion in the literature on this and the following two species, only recent reviews are included.) 66, 67, 68, 129.

Kinosternon integrum 68, 129.

Kinosternon sonoriense 65, 66, 68, 129, 194.

Order Squamata

Suborder Sauria

Anolis nebulosus 38, 57, 94, 175, 176, 194.

Barisia imbricata 38, 55, 121, 132, 157, 158, 175, 176, 189.

Barisia levicollis 55, 132, 157, 194.

Cnemidophorus costatus 57, 94, 152 (as perplexus), 175, 187.

Cnemidophorus exsanguis 20, 41, 160, 194.

Elgaria kingii 11, 35, 43, 79, 94, 132, 152, 156, 160, 163, 169, 175, 176, 194.

Eumeces brevirostris 13, 34, 38, 42, 56, 82, 107, 132, 137, 146, 167, 175, 176, 184, 194.

Eumeces lynxe 38, 82, 100, 137, 167, 175.

Eumeces multilineatus 2 (as multivirgatus), 7 (as multivirgatus), 82, 95, 136, 194 (also as multivirgatus).

Eumeces parviauriculatus 82, 94, 107, 132, 152, 194.

Eumeces parvulus 57, 107.

Eumeces tetragrammus 79, 82, 83, 94, 107, 142, 146, 152, 194.

Gerrhonotus liocephalus 57, 175, 176, 178.

Phrynosoma douglassii 98, 160.

Phrynosoma orbiculare 12, 38, 42, 56, 64, 93, 97, 98, 105, 118, 121, 132, 144, 152, 160, 175, 194.

Sceloporus bulleri 45, 57, 87 (as torquatus), 166, 175, 176, 184.

Sceloporus clarkii 11, 57, 94, 144, 175, 194.

Sceloporus grammicus 2, 7, 12, 17, 21, 24, 35, 38, 54, 56, 79, 115, 120, 152, 160, 175, 176, 177, 194.

Sceloporus heterolepis 15, 17, 57, 78, 120, 168, 174, 175.

Sceloporus jarrovii 2, 7, 12, 17, 23, 33, 35, 38, 42, 54, 56, 57, 79, 94, 117, 120, 152, 160, 175, 176, 177, 178, 184, 194.

Sceloporus nelsoni 57, 94, 175.

Sceloporus poinsettii 12, 17, 21, 24, 35, 38, 42, 56, 79, 94, 109, 117, 126, 139, 152, 160, 175, 176, 194.

Sceloporus scalaris 4, 27, 35, 38, 54 (as aeneus), 79, 114, 116, 120, 127, 154, 160, 175, 194.

Sceloporus spinosus 94, 194.

Sceloporus torquatus 117.

Sceloporus virgatus 25, 26, 79, 80, 94, 152, 160, 180, 194.

Urosaurus ornatus 35, 160, 194.

#### Suborder Serpentes

Adelophis foxi 109, 110, 175.

Coluber constrictor 125, 175, 181, 182.

Conopsis nasus 14, 38, 56, 57, 72, 140, 142, 152, 153, 175, 176, 178.

Crotalus basiliscus 8, 57, 92, 175.

Crotalus lepidus 8, 16, 38, 53, 57, 58, 60, 76, 79, 88, 124, 142, 143, 151, 160, 175, 184.

Crotalus molossus 8, 35, 38, 53, 76, 77, 103, 124, 142, 152, 160, 175, 176.

Crotalus pricei 1, 8, 24, 32, 35, 38, 42, 53, 76, 77, 91, 118, 124, 142, 152, 160, 175, 176, 177.

Crotalus scutulatus 104, 160.

Crotalus stejnegeri 29, 76, 77.

- Crotalus willardi 8, 35, 53, 59, 60, 75, 76, 124, 142, 152, 160, 175, 193.
- Diadophis punctatus 10, 48, 50, 89, 130, 142, 145, 152, 160, 175.
- Dryadophis cliftoni 57, 175.
- Geophis dugesii 37, 57, 81, 142, 172, 175, 192.
- Hypsiglena torquata 160.
- Lampropeltis mexicana 47, 49, 51, 52, 162, 175.
- Lampropeltis pyromelana 79, 124, 130, 135, 141, 142, 148, 160.
- Leptophis diplotropis 57, 175.
- Masticophis mentovarius 57, 70, 71, 175.
- Masticophis taeniatus 101, 160, 184.
- Pituophis deppei 31, 39, 175.
- Pituophis melanoleucus 74, 134, 142, 160.
- Rhadinaea hesperia 57, 99, 142, 175.
- Rhadinaea laureata 38, 99, 174, 175.
- Salvadora bairdi 57, 94, 175.
- Salvadora grahamiae 142, 160, 175.
- Senticolis triaspis 36, 142, 152.
- Storeria storerioides 2, 56, 57, 94, 142, 145, 175, 176.
- Tantilla bocourti 90, 94, 176.
- Tantilla wilcoxi 84, 94, 123, 130, 42, 152.
- Thamnophis cyrtopsis 30 (part), 38 (as eques), 57, 79, 94, 96, 112 (as eques [part]), 122 (as eques), 124 (as eques), 128 (as eques), 138, 142, 152 (as eques), 160, 165, 173, 175.
- Thamnophis elegans 9, 24 (as cyrtopsis), 30 (as cyrtopsis [part]), 44, 122, 128, 130, 138, 142, 160, 171, 175.
- Thamnophis eques 30, 35, 38 (as macrostemma), 42, 108, 109, 112 (part), 119, 122 (as macrostemma megalops), 124 (as macrostemma megalops), 128 (as macrostemma), 138, 142, 160, 175, 177.
- Thamnophis melanogaster 30, 38, 109, 112, 122, 138, 142, 175.



Thamnophis melanogaster 30, 38, 109, 112, 122, 138, 142, 175.

Thamnophis nigronuchalis 30 (as rufipunctatus [part]), 122 (as rufipunctatus [part]), 142, 155 (also as rufipunctatus [part]), 175.

Thamnophis rufipunctatus 122 (part), 124, 128, 138, 142, 150, 152, 155 (part), 160, 177.

Trimorphodon tau None. Our record based upon LSUMZ 35157 collected in Nayarit at 22.4 mi SW Las Canoas, Durango at 2370 m.

## BIBLIOGRAPHY

1. Amaral, A. 1927. Studies of nearctic Ophidia II. Crotalus pricei Van Denburgh, 1896, a synonym of C. triseriatus (Wagler, 1830). Bull. Antivenin Inst. Amer. 1(2): 48-54.
2. Anderson, J.D. 1960. Storeria storerioides in western Mexico. Herpetologica 16(1): 63-66.
3. Anderson, J.D. 1961. The life history and systematics of Ambystoma rosaceum. Copeia 1961(4): 371-377.
4. Anderson, J.D. 1962. Egg laying and nesting in Sceloporus scalaris slevini. Herpetologica 18(3): 162-164.
5. Anderson, J.D. 1978. Ambystoma rosaceum. Cat. Amer. Amphib. Rept. 206.1-206.2.
6. Anderson, J.D., and R.G. Webb. 1978. Life history aspects of the Mexican salamander Ambystoma rosaceum (Amphibia, Urodela, Ambystomidae). J. Herpetol. 12(1): 89-93.
7. Anderson, J.D., and D.C. Wilhoft. 1959. A new subspecies of Eumeces multivirgatus from Mexico. Copeia 1959(1): 57-60.
8. Armstrong, B.L., and J.B. Murphy. 1979. The natural history of Mexican rattlesnakes. Univ. Kansas Mus. Natur. Hist. Spec. Publ. (5): i-vii + 1-88.
9. Baker, R.H., and R.G. Webb. 1976. Thamnophis elegans captures Sorex emarginatus. Herpetol. Rev. 7(3): 112.
10. Blanchard, F.N. 1942. The ring-neck snakes, genus Diadophis. Bull. Chicago Acad. Sci. 7(1): 1-144.
11. Bogert, C.M., and J.A. Oliver. 1945. A preliminary analysis of the herpetofauna of Sonora. Bull. Amer. Mus. Natur. Hist. 83(6): 297-426.
12. Boulenger, G.A. 1885. Catalogue of the Lizards in the British Museum (Natural History). Second ed. Vol. 2. Taylor and Francis; London.
13. Boulenger, G.A. 1887. Catalogue of the Lizards in the British Museum (Natural History). Second ed. Vol. 3. Taylor and Francis; London.
14. Boulenger, G.A. 1894. Catalogue of the Snakes in the British Museum (Natural History). Vol. 2. Taylor and Francis; London.
15. Boulenger, G.A. 1895 (1894). Second report on additions to the lizard collection in the Natural-History Museum. Proc. Zool. Soc. London 1894: 722-736.
16. Boulenger, G.A. 1896. Catalogue of the Snakes in the British Museum (Natural History). Vol. 3. Taylor and Francis; London.
17. Boulenger, G.A. 1897. A revision of the lizards of the genus Sceloporus. Proc. Zool. Soc. London 1897: 474-522.
18. Boulenger, G.A. 1917. Descriptions of new frogs of the genus Rana. Ann. Mag. Natur. Hist., ser. 8, 20: 413-418.
19. Boulenger, G.A. 1920. A monograph of the American frogs of the genus Rana. Proc. Amer. Acad. Arts Sci. 55: 413-480.
20. Burt, C.E. 1931. A study of the teiid lizards of the genus Cnemidophorus with special reference to their phylogenetic relationships. Bull U.S. Natl. Mus. (154): i-viii, + 1-286.

21. Carpenter, C.C. 1978. Comparative display behavior in the genus Sceloporus (Iguanidae). Milwaukee Pub. Mus. Contrib. Biol. Geol. (18): 1-71.
22. Chrapliwy, P.S. 1956. Extensions of known range of certain amphibians and reptiles of Mexico. *Herpetologica* 12(2): 121-124.
23. Chrapliwy, P.S. 1964. Taxonomy and distribution of the jarrovi complex of lizards of the torquantus group, genus Sceloporus. Ph.D. Thesis, Univ. Illinois.
24. Chrapliwy, P.S., and C.M. Fugler. 1955. Amphibians and reptiles collected in Mexico in the summer of 1953. *Herpetologica* 11(2): 121-128.
25. Cole, C.J. 1963. Variation, distribution, and taxonomic status of the lizard, Sceloporus undulatus virgatus Smith. *Copeia* 1963(2): 413-425.
26. Cole, C.J. 1968. Sceloporus virgatus. *Cat. Amer. Amph. Rept.* 72.1-72.2.
27. Cole, C.J. 1978. Karyotypes and systematics of the lizards in the variabilis, jalapae, and scalaris species groups of the genus Sceloporus. *Amer. Mus. Novitates* (2653): 1-13.
28. Collins, J.P. 1979. Sexually mature larvae of the salamanders Ambystoma rosaceum and A. trigrinum velasci from Chihuahua, Mexico: Taxonomic and ecologic notes. *J. Herpetol.* 13(3): 351-354.
29. Collins, J.T. 1982. Crotalus stejnegeri. *Cat. Amer. Amph. Rept.* 303.1-303.2.
30. Conant, R. 1963. Semiaquatic snakes of the genus Thamnophis from the isolated drainage system of the Rio Nazas and adjacent areas in Mexico. *Copeia* 1963(3): 473-499.
31. Conant, R. 1965. Miscellaneous notes and comments on toads, lizards, and snakes from Mexico. *Amer. Mus. Novitates* (2205): 1-38.
32. Conant, R. 1967. Vacation yields rare rattlesnake. *America's First Zoo* 19(4): 26.
33. Cope, E.D. 1887. Catalogue of batrachians and reptiles of Central America and Mexico. *Bull. U. S. Natl. Mus.* (32): 1-98.
34. Dixon, J.R. 1969. Taxonomic review of the Mexican skinks of the Eumeces brevirostris group. *Los Angeles Co. Mus. Natur. Hist. Contrib. Sci.* (168): 1-30.
35. Dominguez, P., T. Alvarez, and P. Huerta. 1977 (1974). Coleccion de anfibios y reptiles del noroeste de Chihuahua, Mexico. *Rev. Soc. Mexicana Hist. Natur.* 35: 117-142.
36. Dowling, H.G. 1960. A taxonomic study of the ratsnakes, genus Elaphe Fitzinger. VII. The triaspis section. *Zoologica* 45 (6): 53-80.
37. Downs, F.L. 1967. Intrageneric relationships among Colubrid snakes of the genus Geophis Wagler. *Misc. Publ. Mus. Zool. Univ. Michigan* (131): i-iv + 1-193.
38. Drake, J.J. 1958. The brush mouse Peromyscus boylii in southern Durango. *Publ. Michigan St. Mus., Biol. Ser.* 1(3): 97-132.
39. Duellman, W.E. 1960. A taxonomic study of the Middle American snake, Pituophis deppei. *Univ. Kansas Publ. Mus. Natur. Hist.* 10(10): 599-610.
40. Duellman, W.E. 1970. The hylid frogs of Middle America. *Monogr. Mus. Natur. Hist. Univ. Kansas* (1): i-xi + 1-753.

41. Duellman, W.E., and R.G. Zweifel. 1962. A synopsis of the lizards of the sexlineatus group (genus Cnemidophorus). Bull. Amer. Mus. Natur. Hist. 123(3): 155-210.
42. Dunn, E.R. 1936. The amphibians and reptiles of the Mexican expedition of 1934. Proc. Acad. Natur. Sci. Philadelphia 88: 471-477.
43. Fitch, H.S. 1938. A systematic account of the alligator lizards (Gerrhonotus) in the western United States and Lower California. Amer. Midl. Naturalist 20(2): 381-424.
44. Fitch, H.S. 1980. Remarks concerning certain western garter snakes of the Thamnophis elegans complex. Trans. Kansas Acad. Sci. 83(3): 106-113.
45. Fosdick, M.K. 1968. Distributional records from a collection of reptiles from western and central Mexico. Herpeton 3: 1-3.
46. Frost, J.S., and J.T. Bagnara. 1976. A new species of leopard frog (Rana pipiens complex) from northwestern Mexico. Copeia 1976(2): 332-338.
47. Garstka, W.R. 1982. Systematics of the mexicana species group of the colubrid genus Lampropeltis, with an (sic) hypothesis (sic) mimicry. Breviora (466): 1-35.
48. Gehlbach, F.R. 1965. Herpetology of the Zuni Mountains region, northwestern New Mexico. Proc. U. S. Natl. Mus. 116: 243-332.
49. Gehlbach, F.R. 1967. Lampropeltis mexicana. Cat. Amer. Amph. Rept. 55.1-55.2.
50. Gehlbach, F.R. 1974. Evolutionary relations of southwestern ringneck snakes (Diadophis punctatus). Herpetologica 30(2): 140-148.
51. Gehlbach, F.R., and J.K. Baker. 1962. Kingsnakes allied with Lampropeltis mexicana: Taxonomy and natural history. Copeia 1962(2): 291-300.
52. Gehlbach, F.R., and C.J. McCoy, Jr. 1965. Additional observations on variations and distribution of the Gray-banded kingsnake, Lampropeltis mexicana. Herpetologica 21(1): 35-38.
53. Gloyd, H.K. 1940. The rattlesnakes, genera Sistrurus and Crotalus. A study in zoogeography and evolution. Spec. Pub. Chicago Acad. Sci. (4): i-vii + 1-270.
54. Greene, H.W. 1972. Mexican reptiles in the Senckenberg Museum. Carnegie Mus. 15p.
55. Guillette, L.J., Jr., and H.M. Smith. 1982. A review of the Mexican lizard Barisia imbricata, and the description of a new subspecies. Trans. Kansas Acad. Sci. 85(1): 13-33.
56. Gunther, A. 1885-1902. Biologia Centrali-Americana. Reptilia and Batrachia. Porter; London.
57. Hardy, L.M., and R.W. McDiarmid. 1969. The amphibians and reptiles of Sinaloa, Mexico. Univ. Kansas Publ. Mus. Natur. Hist. 18(3): 39-252.
58. Harris, H.S., Jr., and R. S. Simmons. 1972. An April birth record for Crotalus lepidus with a summary of annual broods in rattlesnakes. Bull. Maryland Herpetol. Soc. 8(2): 54-56.
59. Harris, H. S., Jr., and R. S. Simmons. 1976. The paleogeography and evolution of Crotalus willardi, with a formal description of a new subspecies from New Mexico, United States. Bull. Maryland Herpetol. Soc. 12(1): 1-22.

60. Harris, H.S., Jr., and R. S. Simmons. 1978. A preliminary account of the rattlesnakes with the descriptions of four new subspecies. Bull. Maryland Herpetol. Soc. 14(3): 105-211.
61. Hillis, D.M., J.S. Frost, and R.G. Webb. 1984. A new species of frog of the Rana tarahumarae group from southwestern Mexico. Copeia 1984(2): 398-403.
62. Hillis, D.M., J.S. Frost, and D.A. Wright. 1983. Phylogeny and biogeography of the Rana pipiens complex: A biochemical evaluation. Syst. Zool. 32(2): 132-143.
63. Holman, J.A. 1965. A polymorphic deme of Hyla eximia Baird from Durango, Mexico. J. Ohio Herpetol Soc. 5(1): 34.
64. Horowitz, S.B. 1955. An arrangement of the subspecies of the horned toad Phrynosoma orbiculare (Iguanidae). Amer. Midl. Naturalist 54(1): 204-218.
65. Iverson, J.B. 1976. Kinosternon sonoriense. Cat. Amer. Amph. Rept. 176.1-176.2.
66. Iverson, J.B. 1981. Biosystematics of the Kinosternon hirtipes species group (Testudines: Kinosternidae). Tulane Stud. Zool. Bot. 23(1): 1-74.
67. Iverson, J.B. 1985. Kinosternon hirtipes. Cat. Amer. Amph. Rept. 361.1-361.4.
68. Iverson, J.B. 1986. A checklist with Distribution Maps of the Turtles of the World. First ed. Privately Printed; Richmond.
69. Jameson, D.L., J.P. Mackey, and R.C. Richmond. 1966. The systematics of the Pacific tree frog, Hyla regilla. Proc. California Acad. Sci. 33: 551-620.
70. Johnson, J.D. 1977. The taxonomy and distribution of the Neotropical whipsnake Masticophis mentovarius (Reptilia, Serpentes, Colubridae). J. Herpetol. 11(3): 287-309
71. Johnson, J.D. 1982. Masticophis mentovarius. Cat. Amer. Amph. Rept. 295.1-295.4.
72. Johnson, R.M., and E.A. Liner. 1978. Conopsis nasus nasus (Reptilia, Serpentes, Colubridae) in Chihuahua, Mexico. J. Herpetol. 12(1): 108-109.
73. Kellogg, R. 1932. Mexican tailless amphibians in the United States National Museum. Bull. U. S. Natl. Mus. (160): i-iv + 1-224.
74. Klauber, L.M. 1947. Classification and ranges of the gopher snakes of the genus Pituophis in the western United States. Bull. Zool. Soc. San Diego (22): 1-81.
75. Klauber, L.M. 1949. The subspecies of the ridge-nosed rattlesnake, Crotalus willardi. Trans. San Diego Soc. Natur. Hist. 11(8): 121-140.
76. Klauber, L.M., 1952. Taxonomic studies of the rattlesnakes of mainland Mexico. Bull. Zool. Soc. San Diego (26): 1-143.
77. Klauber, L.M. 1972. Rattlesnakes: Their habits, life histories, and influence on mankind. 2 vols. Univ. California Press; Berkeley.

78. Langebartel, D.A. 1959. A new lizard (Sceloporus) from the Sierra Madre Occidental of Mexico. *Herpetologica* 15(1): 25-27.
79. Lara-Gongora, G. 1986. New distributional records for some Mexican reptiles and amphibians. *Bull. Maryland Herpetol. Soc.* 22(2): 62-67.
80. Larsen K.R., and W.W. Tanner. 1974. Numeric analysis of the lizard genus Sceloporus with special reference to cranial osteology. *Great Basin Naturalist* 34(1): 1-41.
81. Legler, J.M. 1959. A new snake of the genus Geophis from Chihuahua, Mexico. *Univ. Kansas Pub. Mus. Natur. Hist.* 11(4): 327-334.
82. Legler, J.M., and R.G. Webb. 1960. Noteworthy records of skinks (genus Eumeces) from northwestern Mexico. *Southwestern Naturalist* 5(1): 16-20.
83. Lieb, C.S. 1985. Systematics and distribution of the skinks allied to Eumeces tetragrammus. *Los Angeles Co. Mus. Natur. Hist. Contrib. Sci.* (327): 1-19.
84. Liner, E.A. 1983. Tantilla wilcoxi. *Cat. Amer. Amph. Rept.* 345.1-345.2.
85. Lowe, C.H., C.J. Jones, and J.W. Wright. 1968. A new plethodontid salamander from Sonora, Mexico. *Los Angeles Co. Mus. Natur. Hist. Contrib. Sci.* (140): 1-11.
86. Maldonado-Koerdell, M. 1964. Algunas problemas de sistematica y distribucion de la familia Amystomidae (Urodela, Amphibia) en la republica Mexicana. *Mems. Rev. Acad. Nac. Cienc. Antonio Alzate* 59(3/4): 379-388.
87. Martin, P.S. 1958. A biogeography of reptiles and amphibians in the Gomez Farias region, Tamaulipas, Mexico. *Misc. Pub. Mus. Zool. Univ. Michigan* (101): 1-102.
88. Martin del Campo, R. 1935. Nota acerca de la distribucion geografica de los reptiles ponzonosas en Mexico. *Ann. Inst. Biol. Univ. Mexico* 6(3/4): 219-300.
89. McCoy, C.J., Jr. 1964. Notes on snakes from northern Mexico. *Southwestern Naturalist* 9(1): 46-48.
90. McCranie, J.R. 1977. First record of Tantilla bocourti (Reptilia: Colubridae) from Zacatecas, Mexico. *Southwestern Naturalist* 22(2): 275.
91. McCranie, J.R. 1981a. Crotalus pricei. *Cat. Amer. Amph. Rept.* 266.1-266.2.
92. McCranie, J.R. 1981b. Crotalus basiliscus. *Cat. Amer. Amph. Rept.* 283.1-283.2.
93. McDiarmid, R.W. 1963. A collection of reptiles and amphibians from the highland faunal assemblage of western Mexico. *Los Angeles Co. Mus. Natur. Hist. Contrib. Sci.* (68): 1-15.
94. McDiarmid, R.W., J.F. Copp, and D.E. Breedlove. 1976. Notes on the herpetofauna of western Mexico: New records from Sinaloa and the Tres Marias Islands. *Los Angeles Co. Mus. Natur. Hist. Contrib. Sci.* (275): 1-17.
95. Mecham, J.S. 1980. Eumeces multivirgatus. *Cat. Amer. Amph. Rept.* 241.1-241.2.
96. Milstead, W.W. 1953. Geographic variation in the garter snake, Thamnophis cyrtopsis. *Texas J. Sci.* 5(3): 348-379.
97. Montanucci, R.R. 1979. Notes on systematics of horned lizards allied to Phrynosoma orbiculare (Lacertilia: Iguanidae). *Herpetologica* 35(2): 116-124.

98. Montanucci, R.R. 1981. Habitat separation between Phrynosoma douglassi and P. orbiculare (Lacertilia: Iguanidae) in Mexico. Copeia 1981(1): 147-153.
99. Myers, C.W. 1974. The systematics of Rhadinaea (Colubridae), a genus of New World snakes. Bull. Amer. Mus. Natur. Hist. 153(1): 1-262.
100. Parker, R.B. 1960. The status of a Mexican Lizard, Eumeces lynxe belli. Copeia 1960(4): 284-286.
101. Parker, W.S. 1982. Masticophis taeniatus. Cat. Amer. Amph. Rept. 304.1-304.4.
102. Platz, J.E., and J.S. Mecham. 1979. Rana chiricahuensis, a new species of leopard frog (Rana pipiens complex) from Arizona. Copeia 1979(3): 383-390.
103. Price, A.H. 1980. Crotalus molossus. Cat. Amer. Amph. Rept. 242.1-242.2.
104. Price, A.H. 1982. Crotalus scutulatus. Cat. Amer. Amph. Rept. 291.1-291.2.
105. Reeve, W.L. 1952. Taxonomy and distribution of the horned lizards genus Phrynosoma. Univ. Kansas Sci. Bull. 34(14): 817-960.
106. Riemer, W.J. 1955. Comments on the distribution of certain Mexican toads. Herpetologica 11(1): 17-23.
107. Robinson, M.D. 1979. Systematics of skinks of the Eumeces brevivirostris species group in western Mexico. Los Angeles Co. Mus. Natur. Hist. Contrib. Sci. (319): 1-13.
108. Rossman, D.A., 1965 (1964). A new subspecies of the common garter snake, Thamnophis sirtalis, from the Florida Gulf Coast. Proc. Louisiana Acad. Sci. 27: 67-73.
109. Rossman, D.A., and R.M. Blaney. 1968. A new natricine snake of the genus Adelophis from western Mexico. Occ. Pap. Mus. Zool. Louisiana St. Univ. (35): 1-12.
110. Rossman, D.A., and V. Wallach. 1987. Adelophis, A. copei, A. foxi. Cat. Amer. Amph. Rept. 408.1-408.2.
111. Ruibal, R. 1957. An altitudinal and latitudinal cline in Rana pipiens. Copeia 1957(3): 212-221.
112. Ruthven, A.G. 1908. Variations and genetic relationships of the garter-snakes. Bull. U.S. Natl. Mus. (61): i + xii + 1-201.
113. Shaffer, H.B. 1983. Biosystematics of Ambystoma rosaceum and A. tigrinum in northwestern Mexico. Copeia 1983(1): 67-78.
114. Shannon, F.A. 1951. Notes on a herpetological collection from Oaxaca and other localities in Mexico. Proc. U.S. Natl. Mus. 101: 465-484.
115. Sites, J.W., Jr., and J.R. Dixon. 1981. A new subspecies of the iguanid lizard, Sceloporus grammicus, from northeastern Mexico, with comments on its evolutionary implications and the status of S. g. disparilis. J. Herpetol. 15(1): 59-69.
116. Smith, H.M. 1937. A synopsis of the scalaris group of the lizard genus Sceloporus. Occ. Pap. Mus. Zool. Univ. Michigan (361): 1-8.
117. Smith, H.M. 1938 (1936). The lizards of the torquatus group of the genus Sceloporus Wiegmann 1828. Univ. Kansas Sci. Bull. 24(21): 539-693.
118. Smith, H.M. 1939a. An annotated list of the Mexican amphibians and reptiles in the Carnegie Museum. Ann. Carnegie Mus. 27: 311-320.

119. Smith, H.M. 1939b. Notes on Mexican reptiles and amphibians. Zool. Ser. Field Mus. Natur. Hist. 24(4): 15-35.
120. Smith, H.M. 1939c. The Mexican and Central American lizards of the genus Sceloporus. Zool. Ser. Field Mus. Natur. Hist. 26: 1-397.
121. Smith, H.M. 1942a. Mexican herpetological miscellany. Proc. U. S. Natl. Mus. 92: 349-395.
122. Smith, H.M. 1942b. The synonymy of the garter snakes (Thamnophis), with notes on Mexican and Central American species. Zoologica 27 (17): 97-123.
123. Smith, H.M. 1942c. A resume of Mexican snakes of the genus Tantilla. Zoologica 27(7): 33-42.
124. Smith, H.M. 1943. Summary of the collections of snakes and crocodilians made in Mexico under the Walter Rathbone Bacon Traveling Scholarship. Proc. U.S. Natl. Mus. 93: 393-504.
125. Smith, H.M. 1971. Distribution of the racer Coluber constrictor in Mexico. J. Herpetol. 5(3/4): 212-214.
126. Smith, H.M., and P.S. Chrapliwy. 1958. New and noteworthy Mexican herptiles from the Lidicker collection. Herpetologica 13(4): 267-271.
127. Smith, H.M., and W.P. Hall. 1974. Contributions to the concepts of reproductive cycles and the systematics of the scalaris group of the lizard genus Sceloporus. Great Basin Naturalist 34(2): 97-104.
128. Smith, H.M., C.W. Nixon, and P.W. Smith. 1950. Mexican and Central American garter snakes (Thamnophis) in the British Museum (Natural History). J. Linnean Soc. Zool. 41: 571-584.
129. Smith, H.M., and R.B. Smith. 1980 (1979). Synopsis of the Herpetofauna of Mexico. VI. Guide to Mexican Turtles. Bibliographic Addendum III. John Johnson; North Bennington.
130. Smith, H.M., and E.H. Taylor. 1945. An annotated checklist and key to the snakes of Mexico. Bull. U. S. Natl. Mus. (187): i-iv + 1-239.
131. Smith, H.M., and E.H. Taylor. 1948. An annotated checklist and key to the amphibia of Mexico. Bull. U. S. Natl. Mus. (194): i-iv + 1-118.
132. Smith, H.M., and E.H. Taylor. 1950. An annotated checklist and key to the reptiles of Mexico exclusive of the snakes. Bull. U. S. Natl. Mus. (199): i-v + 1-253.
133. Stebbins, R.C. 1951. Amphibians of western North America. Univ. California Press; Berkeley.
134. Stull, O.G. 1940. Variations and relationships in the snakes of the genus Pituophis. Bull. U. S. Natl. Mus. (175): i-vi + 1-225.
135. Tanner, W.W. 1953. A study of taxonomy and phylogeny of Lampropeltis pyromelana Cope. Great Basin Naturalist 13(1/2): 47-66.
136. Tanner, W.W. 1957. A new skink of the multivirgatus group from Chihuahua. Great Basin Naturalist 17(3/4): 111-117.
137. Tanner, W.W. 1958. Two new skinks from Durango, Mexico. Great Basin Naturalist 18(2): 57-62.
138. Tanner, W.W. 1959. A new Thamnophis from western Chihuahua with notes on four other species. Herpetologica 15(4): 165-172.
139. Tanner, W.W. 1960. Sceloporus p. poinsetti taken in Sonora. Herpetologica 16(4): 235.
140. Tanner, W.W. 1961. A new subspecies of Conopsis nasus from Chihuahua, Mexico. Herpetologica 17(1): 13-18.



141. Tanner, W.W. 1983. Lampropeltis pyromelana. Cat. Amer. Amph. Rept. 342.1-342.2.
142. Tanner, W.W. 1985. Snakes of western Chihuahua. Great Basin Naturalist 45(4): 615-676.
143. Tanner, W.W., J.R. Dixon, and H.S. Harris, Jr. 1972. A new subspecies of Crotalus lepidus from western Mexico. Great Basin Naturalist 32(1): 16-24.
144. Tanner, W.W. and W.G. Robinson, Jr. 1960a (1959). A collection of herptiles from Urique, Chihuahua. Great Basin Naturalist 19(4): 75-82.
145. Tanner, W.W., and W.G. Robinson, Jr. 1960b. New and unusual serpents from Chihuahua, Mexico. Herpetologica 16(1): 67-70.
146. Taylor, E.H. 1936. A taxonomic study of the cosmopolitan scincoid lizards of the genus Eumeces with an account of the distribution and relationships of its species. Univ. Kansas Sci. Bull. 23: 1-643.
147. Taylor, E.H. 1940a. A new frog from the Tarahumara Mountains of Mexico. Copeia 1940(4): 250-253.
148. Taylor, E.H. 1940b. A new Lampropeltis from western Mexico. Copeia 1940(4): 253-255.
149. Taylor, E.H. 1941a. Two new ambystomid salamanders from Chihuahua. Copeia 1941(3): 143-146.
150. Taylor, E.H. 1941b. Herpetological miscellany, No. II. Univ. Kansas Sci. Bull. 27(7): 105-139.
151. Taylor, E.H. 1944. Two new species of Crotalid snakes from Mexico. Univ. Kansas. Sci. Bull. 30(4): 47-56.
152. Taylor, E.H., and I.W. Knobloch. 1940. Report on an herpetological collection from the Sierra Madre Mountains of Chihuahua. Proc. Biol. Soc. Washington 53: 125-130.
153. Taylor, E.H., and H.M. Smith. 1942. The snake genera Conopsis and Toluca. Univ. Kansas Sci. Bull. 28(15): 325-363.
154. Thomas, R.A., and J.R. Dixon. 1976. A re-evaluation of the Sceloporus scalaris group (Sauria: Iguanidae). Southwestern Naturalist 20(4): 523-536.
155. Thompson, F.G. 1957. A new Mexican gartersnake (genus Thamnophis) with notes on related forms. Occ. Pap. Mus. Zool. Univ. Michigan (584): 1-10.
156. Tihen, J.A. 1948. Two races of Elgaria kingii Gray. Trans. Kansas Acad. Sci. 51(3): 299-301.
157. Tihen, J.A. 1949. A review of the lizard genus Barisia. Univ. Kansas Sci. Bull. 33(3): 217-256.
158. Tihen, J.A. 1954. Gerrhonotine lizards recently added to the American Museum collection, with further revisions of the genus Abronia. Amer. Mus. Novitates (1687): 1-26.
159. Van Devender, T.R. 1973. Populations of Ambystoma tigrinum and A. rosaceum in Chihuahua, Mexico. J. Arizona Acad. Sci. 8(1): 84.
160. Van Devender, T.R., and C.H. Lowe, Jr. 1977. Amphibians and reptiles of Yepomera, Chihuahua, Mexico. J. Herpetol. 11(1): 41-50.
161. Wake, D.B., and J.F. Lynch. 1976. The distribution, ecology, and evolutionary history of Plethodontid salamanders in Tropical America. Los Angeles Co. Mus. Natur. Hist. Sci. Bull. (25): 1-65.

162. Webb, R.G. 1961. A new kingsnake from Mexico, with remarks on the mexicana group of the genus Lampropeltis. Copeia 1961(3): 326-333.
163. Webb, R.G. 1962a. A new alligator lizard (genus Gerrhonotus) from western Mexico. Herpetologica 18(2): 73-79.
164. Webb, R.G. 1962b. A new species of frog (genus Tomodactylus) from western Mexico. Univ. Kansas Pub. Mus. Natur. Hist. 15(3): 175-181.
165. Webb, R.G. 1966. Resurrected names for Mexican populations of black-necked garter snakes Thamnophis cyrtopsis (Kennicott). Tulane Stud. Zool. 13(2): 55-70.
166. Webb, R.G. 1967. Variation and distribution of the iguanid lizard Sceloporus bulleri, and the description of a related new species. Copeia 1967(1): 202-213.
167. Webb, R.G. 1968. The Mexican skink Eumeces lynxe (Squamata, Scincidae). Pub. Michigan St. Mus. Biol. Ser. 4(1): 1-28.
168. Webb, R.G. 1969. Variation, status, and relationship of the iguanid lizard Sceloporus shannonorum. Herpetologica 25(4): 300-307.
169. Webb, R.G. 1970. Gerrhonotus kingii. Cat. Amer. Amph. Rept. 97.1-97.4.
170. Webb, R.G. 1972. Resurrection of Bufo mexicanus Brocchi for a highland toad in western Mexico. Herpetologica 28(1): 1-6.
171. Webb, R.G. 1976. A review of the garter snake Thamnophis elegans in Mexico. Los Angeles Co. Mus. Natur. Hist. Contrib. Sci. (284): 1-13.
172. Webb, R.G. 1977. Comments on snakes of the genus Geophis (Colubridae) from the Mexican states of Durango and Sinaloa. Southwestern Naturalist 21(4): 548-551.
173. Webb, R.G. 1980. Thamnophis cyrtopsis. Cat. Amer. Amph. Rept. 245.1-245.4.
174. Webb, R.G. 1982. Distributional records for Mexican reptiles. Herpetol. Rev. 13(4): 132.
175. Webb, R.G. 1984. Herpetogeography in the Mazatlan-Durango region of the Sierra Madre Occidental, Mexico. Pp. 217-241. In R.A. Seigel, L.E. Hunt, J.L. Knight, L. Malaret, and N.L. Zuschlag (eds.). Vertebrate ecology and systematics. A tribute to Henry S. Fitch. Univ. Kansas Mus. Natur. Hist. Spec. Publ (10): 1-278.
176. Webb, R.G., and R.H. Baker. 1962. Terrestrial vertebrates of the Pueblo Nuevo area of southwestern Durango, Mexico. Amer. Midland Naturalist 68(2): 325-333.
177. Webb, R.G., and R.H. Baker. 1984. Terrestrial vertebrates of the Cerro Mohinora region, Chihuahua, Mexico. Southwestern Naturalist 29(2): 243-246.
178. Webb, R.G., and M. Hensley. 1959. Notes on reptiles from the Mexican state of Durango. Pub. Michigan St. Mus. Biol. Ser. 1(6): 249-258.
179. Welbourn, W.C., Jr., and R.B. Loomis. 1970. Three new species of Hannemania (Acarina, Trombiculidae) from amphibians of western Mexico. Bull. So. California Acad. Sci. 69(2): 65-73
180. Williams, K.L. 1960. Taxonomic notes on Arizona herpetozoa. Southwestern Naturalist 5(1): 25-36.
181. Wilson, L.D. 1966. The range of the Rio Grande racer in Mexico and the status of Coluber oaxaca (Jan). Herpetologica 22(1): 42-47
182. Wilson, L.D. 1978. Coluber constrictor. Cat. Amer. Amph. Rept. 218.1-218.4.

183. Zweifel, R.G. 1954a. A new frog of the genus Rana from western Mexico with a key to the Mexican species of the genus. Bull. So. California Acad. Sci. 53(3): 131-141.
184. Zweifel, R.G. 1954b. Notes on the distribution of some reptiles in western Mexico. Herpetologica 10(3): 145-149.
185. Zweifel, R.G. 1955. Ecology, distribution, and systematics of frogs of the Rana boylei group. Univ. California Pub. Zool. 54(4): 207-292.
186. Zweifel, R.G. 1956. A survey of the frogs of the augusti group, genus Eleutherodactylus. Amer. Mus. Novitates (1813): 1-35.
187. Zweifel, R.G. 1959. Variation in and distribution of lizards of western Mexico related to Cnemidophorus sacki. Bull. Amer. Mus. Natur. Hist. 117(2): 57-116.
188. Zweifel, R.G. 1968. Rana tarahumarae. Cat. Amer. Amph. Rept. 66.1-66.2.

## ADDENDUM


189. Guillette, L.J., Jr., and G. Casas-Andreu. 1987. The reproductive biology of the high elevation Mexican lizard Barisia imbricata. Herpetologica 43(1): 29-38.
190. Jones, T.R., J.P. Collins, T.D. Kocher, and J.B. Mitton. 1988. Systematic status and distribution of Ambystoma tigrinum stebbinsi (Amphibia: Caudata). Copeia 1988(3): 621-635.
191. Price, A.H., and B.K. Sullivan. 1988. Bufo microscaphus. Cat. Amer. Amph. Rept. 415.1-415.3.
192. Restrepo T., J.H., and J.W. Wright. 1987. A new species of the Colubrid snake genus Geophis from Colombia. J. Herpetol. 21(3): 191-196.
193. Stille, B. 1987. Dorsal scale microdermatoglyphics and rattlesnake (Crotalus and Sistrurus) phylogeny (Reptilia: Viperidae: Crotalinae). Herpetologica 43(1): 98-104.
194. Tanner, W.W. 1987. Lizards and turtles of western Chihuahua. Great Basin Naturalist 47(3): 383-421.




514 202SI XL 315  
06/93 198115 2













SMITHSONIAN INSTITUTION LIBRARIES



3 9088 01298 2195